

Sri

MANAKULA VINAYAGAR




Medical College and Hospital

Kalitheerthalkuppam, Madagadipet, Puducherry – 605107.

Department of Pharmacy

DRUG FORMULARY

Edition-II



PREFACE

This Drug Formulary of Sri Manakula Vinayagar Medical College and Hospital is a concise and handy book designed for quick reference of current essential drugs in medical practice. As it is not possible to include all drugs, commonly used drugs in our hospital with their recommended dosages, both in adults and children, adverse drug reaction and cost have been included. However, as the medical knowledge is vast and continuously being upgraded, changes may have occurred and may not be reflected in the incorporated information. Clinical judgment and experience should be used while prescribing the drugs.

Note: Dose of drug varies with indications and severity of disease, the given doses is just for reference, and is for the most common indication. Where feasible, the dose ranges from lowest to maximal permissible doses have been given. Paediatric doses are for children more than 1 month of age i.e. neonatal doses may vary. The price given is for an estimation of the cost of therapy and not the exact selling price, which may vary.

“Let food be thy medicine and medicine be thy food.”

-Hippocrates

CONTENTS

SL. NO.	SECTION	PAGE NO.
1	ANTI-ASTHMA, DECONGESTANTS , OTHER RS	1
2	ANTI-BACTERIALS	5
3	ANTI-CANCER	15
4	ANTI-COAGULANTS AND THROMBOLYTIC	19
5	ANTI-DEPRESSANTS	21
6	ANTI-DIABETIC	22
7	ANTI-EMETIC	23
8	ANTI-EPILEPTICS	24
9	ANTI-FUNGALS	26
10	ANTI-HISTAMINE	28
11	ANTI-HYPERTENSIVE	31
12	ANTI-MALARIAL	34
13	ANTI-PARASITIC	35
14	ANTI-PARKINSONISM	36
15	ANTI-PSYCHOTICS	37
16	ANTI-TUSSIVES & EXPECTORANTS	38
17	ANTI-ULCER	40

18	ANTI-VIRAL	43
19	CNS - OTHERS	43
20	CVS - OTHERS	46
21	HORMONES & THEIR ANTAGONIST	51
22	HYPOLIPIDEMIC DRUGS	57
23	MUSCLE RELAXANTS, ANAESTHETIC & EMERGENCY	58
24	NSAIDS & GOUT	63
25	SEDATIVES AND HYPNOTICS	69
26	VACCINE	70
27	MISCELLANEOUS	70

**“Writing prescriptions is easy, understanding people is
hard”**

- Franz kafker, 1883-1924

ANTI-ASTHMA, DECONGESTANTS , OTHER RS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ACEBROPHYLLIN	AB PHYLLINE - 100ML	Tablets, Capsules, Syrup	100 mg twice daily Children up to 2 years, it's half a teaspoon, twice daily. Age 2 to 5 is half a teaspoon, three times a day	107.00	Restlessness, Tremor, Palpitation, Tachypnoea, Flushing, Hypotension
		AB-PHYLLINE - 100 MG			81.50	
		AB-PHYLLINE SR - 200 MG			141.00	
		ASCOVENT			77.50	
		ASCOVENT-SR			129.00	
2	ACEBROPHYLLIN+N-ACETYLCYSTEINE	AB-PHYLLINE-N-TABLET	Tablet	Acebrophylline 100 mg+N-Acetylcysteine 600 mg OD	149.00	Restlessness, Tremor, Palpitation, Tachypnoea, Flushing, Hypotension
3	AMINOPHYLLINE	AMINOPHYLLINE. I.P. - 10 ML	Injection	Adult: Loading dose IV: 6.3 mg/kg over 30 min Maintenance dose : 0.3 - 0.8 mg/kg/hr Paed dose: Maintenance dose IV 0.8 mg/kg/hr	20.00	GI upset, Rapid IV causes precordial pain, syncope & even sudden death
4	BECLOMETHASONE+LEVOSALBUTAMOL	AEROCORT ROTOCAPS - 30 S	ROTACAPS	1 rotacap 3-4 times daily	35.52	Taste disturbance, Itching of throat & Dryness
5	BUDESONIDE	DERINIDE-0.5 - 2ML	Inhaler, Rotacaps, Respules	Budesonide: 100/200 mcg	24.00	Dryness, Irritation, Smell & taste disturbance
		BUDECORT - 100			285.00	
		BUDECORT - 200			396.00	
		BUDECORT - 100MG			67.00	
		BUDECORT - 200MG			86.00	
		BUDECORT - 0.5 MG			30.00	
6	ETOPHYLLINE AND THEOPHYLLINE	DERIPHYLLIN RETARD - 150 MG	Tablet, Injection	Eto: 115 mg Theo: 35 mg	5.80	Restlessness, tremor, palpitation, tachypnoea, flushing, hypotension
		DERIPHYLLIN - 100 MG			3.55	
		DERIPHYLLIN			4.35	
7	FORMOTEROL FUMARATE + BUDESONIDE	FORACORT 200	Inhaler, Rotacaps, Respules	Formoterol: 6mcg Budesonide: 100/200 mcg	175.00	Nasal irritation, sneezing, itching of throat& dryness
		FORACORT - 0.5MG-2ML			39.00	
	FORACORT - 200 MG	323.00				
	FORACORT - 400 MG	426.00				
	FORACORT 1MG-2ML	44.00				
	FORMONIDE - 200 MG	333.00				
	FORMONIDE - 100 MCG	161.00				
	FORMOFLO -100	143.50				
	FORMONIDE - 200 MG	333.00				
	AIRTEC FB - 200MCG MDI	277.75				
	BUDAMATE - 200	322.50				

		BUDAMATE - 200			159.00	
		BUDAMATE - 400			408.50	
		BUDAMATE -400			205.50	
		FORACORT-100mcg INHALER			235.99	
		FORMOFLO-250mg INHALER			598.65	
		FORMONIDE-0.5mg/2ml RESPULES			43.90	
		FOMTIDE-200mg INHALER			287.43	
		BUDETROL-MDI 200mg			275.94	
8	FORMOTEROL 6 MCG+MOMETASONE FUROATE	FORMOST-200mg(DPT) CAPSULE(30'S)	Respicals	Formoterol: 6mcg 1-2 times/day Mometasone: 200 mcg 1-2 times/day	247.00	Dizziness, headache, dry mouth, mild sore throat, runny or stuffy nose, nausea, stomach pain or upset, nervousness, tiredness tremor and trouble sleeping
9	FORMOTEROL FUMARATE + BUDESONIDE + IPRATROPIUM BROMIDE	TRIMIUM TRANSHALER	Inhaler	Formoterol: 6mcg Budesonide: 100/200 mcg	751.00	Nasal irritation, sneezing, itching of throat& dryness
10	IPRATROPIUM BROMIDE	IPRAVENT (RESP)SOLUTION 15ML	Inhaler, Respules	Adult: Nebulized 500 mcg 3-4 times/day Paed:Nebulized 125-250mcg 3-4 times/day	46.00	Drymouth, Urinary retention,
		IPRAVENT- 2 ML			14.00	Broncho spasm
		DUOLIN INHALER 200 MD			233.00	
11	IPRATROPIUM BROMIDE 500 MCG+LEVOSALBUTAMOL	DUOLIN RESPULES-2.5ml	Respules	One respule 3-4 times/day by nebulization	10.59	Duolin can include fine tremor of skeletal muscles and nervousness and less frequently, tachycardia, dizziness, palpitations or headache, especially in hypersensitive patients.
12	LEVOSALBUTAMOL	LEVOLIN - 0.31 MG - 2.5 ML	Tablet, Inhaler, Respules,	Adult: 0.63 mg TID (Nebulizer) 2 mg (Tablet)	3.37	Headache, Tremor,
		LEVOLIN - 2 MG	Rotacaps, Syrup		7.78	Tachycardia
		LEVOLIN INHALER			120.00	
		LEVOLIN SYRUP - 100ML			16.55	
		SALBAIR TRANSCAP - 30'S			25.70	
		DUOLIN ROTOCAPS			63.00	
		SALBAIR - INHALER			115.50	
		DUOLIN-LD - 2.5ML			10.25	
13	MONTELUKAST SODIUM	MONTAIR - 10 MG	Tablet	Adult: 10 mg	137.00	Dizziness, Fatigue
		TELEKAST- 5 MG		Paed : 5mg	79.50	
	MONTELUKAST SODIUM + LEVOCETRIZINE	MONTEK-LC			114.00	
		MONTGOLD			84.00	
		ROMILAST - L - 10 MG			89.50	
		TELEKAST-L-KID			69.00	
		TELEKAST-L			102.00	

14	MONTELUKAST SODIUM + FEXOFENADINE	MONTI - FX	Tablet	Adult : 10mg + 120mg	118.10	Dizziness, Fatigue
		TELEKAST-F			121.00	
15	MONTELUKAST SODIUM + ACEBROPHYLLIN	TELEKAST-A	Tablet	Adult: 10 mg + 100mg	126.00	Dizziness, Fatigue
16	SALBUTAMOL	ASTHALIN INHALER	Tablet, syrup, Respules, Rotacaps, Inhaler	Oral: 2-4 mg 100- 200 mcg / inhalation	127.13	Palpitation, tremor, restlessness
		ASTHALIN - 2.5 ML RESPULES			4.00	
		ASTHALIN(RESP)SOLUTION-15ML			15.20	
		ASTHALIN ROTOCAPS 30S			16.26	
		ASTHALIN SYRUP			16.33	
		ASTHALIN - 2MG			4.80	
		ASTHALIN - 4			6.00	
17	SALMETEROL + FLUTICASONE PROPIONATE	SEROFLO - 125MG INHALER	Inhaler, Rotacaps	125 - 200 mcg	346.00	Mouth & throat candidiasis throat irritation
		SEROFLO - 250 MG INHALER			485.00	
		SEROFLO - 250 ROTOCAPS			253.00	
		SEROFLO 100 ROTACAPS			130.50	
18	ARFOMOTEROL TARTRATE	FORATEC - RESPULES	Respules	15 mcg inhaled by nebulization twice daily (morning and evening). Maximum Daily Dose:30mcg	24.00	Dyspnea, chest pain, Peripheral edema
19	AZELASTINE HYDROCHLORIDE & FLUTICASONE PROPIONATE	DUONASE NASAL SPRAY	Nasal spray	Nasal Rhinitis 140 mcg into each nostril twice daily.	264.00	Irritation, stinging & itching of the nasal mucosa. Sneezing, nosebleeds, headache; nausea, taste disturbances, somnolence, dry mouth
20	DOXOFYLLINE	DOXOLIN - 400 MG	Tablet	Adult: Up to 1200 mg daily.	54.30	Restlessness, tremor, palpitation, tachypnoea, flushing, hypotension
21	EPHEDRINE	EFIPRES 1ML	Injection	As ephedrine sulfate: 12.5-25 mg.	23.40	Tachycardia, anxiety, nausea, restlessness, insomnia, tremor, dry mouth, impaired circulation to the HTN, headache, cardiac arrhythmia.
22	FLUTICASONE PROPIONATE	FLUTICONE - 12ML	Nasal spray, Ointment, Inhaler, Capsule	100 mcg bd	251.00	Topical: Pruritus, hypertrichosis, dryness, numbness of fingers, burning, eruptions, hypopigmentation, allergic contact dermatitis,
		FLUTIFLO			189.00	
		FLUTIVATE - 20GM			109.15	
		FLUTIVATE-10gm CREAM			133.40	
	FLUTICASONE FUROATE	ENNHALE	275.00			

		FURAMIST			290.00	secondary infection, skin atrophy
		FLUTIFLO-FT			263.50	
	FLUTICASONE AZELASTINE	FURAMIST-AZ			326.50	
		FLOHALE (RESP)0.5MG - 2ML			34.50	
	FLUTICASONE FORMOTEROL	MAXIFLO - 250 ROTO CAPS			244.00	
		MAXIFLO - 100 ROTO CAPS			152.00	
		MAXIFLO - 125			370.00	
		MAXIFLO - 250			490.00	
23	ISOTONIC	SOLSPRE NASAL SPRAY	Nasal spray		254.19	Nasal irritation
24	KETOTIFEN	KETASMA	Tablet	1 mg twice daily, up to 2 mg twice daily	40.00	Sedation, drowsiness, dizziness, dry mouth, weightgain, increasedappetite, CNS stimulation. Rarely, cystitis. Conjunctival injection, headaches and rhinitis (ophthalmic).
25	MONTELUKAST SODIUM	MONTAIR - 10 MG	Tablet	10 mg/day	137.00	Churg-Strauss syndrome; headache, dizziness, asthenia /fatigue, dream abnormalities, hallucinations, agitation
26	ORCIPRENALINE SULFATE	ORCIBEST - 10MG	Tablet	20 mg 3-4 times/day	62.00	Tachycardia, nervousness, increased serum glucose, increased potassium levels, tremor, palpitation
27	PIRFENIDONE	PIRFENEX - 200MG	Tablet	267 - 534 TDS	185.00	Increased ALT, increased AST, increased gamma glutamyl transferase, Photosensitivity, Rash, abdominal pain/ distension/ discomfort, headache,anorexia
28	ROFLUMILAST	ROFADAY	Tablet	500 mcg OD	125.00	GI upset, moderate weight loss, acute pancreatitis, dizziness, headache
29	SERATRODAST	SERADAIR - 80MG	Tablet	80 mg	197.00	GI irritation, Throat dryness
30	PINE BARK EXTRACT	PICNOVIT-50mg TABLET	Tablet	50 mg OD	175.00	Gastrointestinal discomfort, nausea, dizziness, headache, sleepiness, urinary retention, urinary frequency, constipation, and increased perspiration.

31	LEVOCETIRIZINE (2.5MG/5ML)	TELEKAST-L-KID-30ml SYRUP	Syrup	1.25 mg (1/2 teaspoon oral solution) (2.5 mL) once daily in the evening	40.00	Nausea, Diarrhoea, Vomiting, Dryness in mouth, Headache, Skin rash, Flu-like symptoms, Fatigue, Sleepiness.
32	SODIUM 2-MERCAPTOETHANESULPHONATE	MISTABRON-3ml RESPULES	Respules	3 to 6 mL (1 to 2 ampoules) twice a day. In severe cases, this dosage can be increased up to a maximum of 24 mL (8 ampoules) a day	185.83	rash, itching, breathing difficulty, chest tightness, and swelling of the mouth, face, lips, or tongue while taking mesna.
33	PROTINE	LAMINO RESPI-200gm POWDER	POWDER	Add 2 heaped scoop (approx 30 g) in 100 ml lukewarm milk/water OD	449.00	Liver toxicity, Flushing or redness of skin, Stomach upset, Side or lower back pain, Diarrhea, Nausea or vomiting, Stomach cramps, Headache, Flushing, Increased blood sugar level, Blurred vision

ANTI-BACTERIALS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMIKACIN	MIKASTAR - 250MG	Injection	250, 500 mg	31.00	Ototoxicity, Nephrotoxicity and Neuromuscular blockade
		MIKASTAR - 500MG			53.35	
		AMIBIOTIC-500mg INJECTION			92.00	
		AMICIN-500mg INJECTION			84.00	
2	AMOXICILLIN	NOVAMOX - 250 MG	Capsules, Syrup,	Adult: 250, 500 mg Paediatric dose: 20- 40 mg/kg/day	47.86	Diarrhea, Alteration of bacterial flora, rashes, HS reactions
		NOVAMOX 250 REDIUSE 30 ML	Tablet		48.00	
		NOVAMOX 500MG			105.68	
		AXL - 250 MG			74.50	
		BLUMOX-500mg CAPSULE	capsule		62.35	
3	AMOXYCILIN & POTASSIUM CLAVULANATE	MOX - CLAV - 1.2 GM	injection	amoxycillin	147.07	Diarrhea, Alteration of bacterial flora, rashes, HS reactions
		MOXCLAV - 625MG	tablet	Adult: 250, 500 mg	250.55	
		NOVAMOX-CV - 30 ML		Paediatric dose:	64.91	
		NOVAMOX-CV 375MG		20- 40 mg/kg/day	140.00	
		CLAMP KID FORTE		Clavulanic acid:125mg	198.00	
		CLAMP - 300 MG			75.50	
		CLAVAM 125 MG			63.50	
		CLAVAM - 625 MG			222.50	
		AUGMENTIN - I.V - 600 MG			99.41	
		AUGMENTIN - 30ML			64.88	
		AUGMENTIN 625 MG			259.73	
		MOXIKIND CV			48.90	

4	AMOXYCILIN & CLAVUNIC ACID	MOXYNIC-1.2mg INJECTION	Injection, tablet, syrup,	Adult: Tab. Amoxicillin 250 - 500 mg + Clavulanate 125 mg qid Children (Oral) : Based on Amoxicillin dose <40 kg: 20- 60mg/kg tid Adult (inj): Based on amoxicillin dose 500 mg tds by slow inj over 3-4 min Children (inj): Based on amoxicillin dose <3 months: 25mg/kg bd; >3 months: 25mg/kg tds	131.96	Rash, Vomiting, Allergic reaction, Nausea, Diarrhoea
		MOXIKIND-CV-KID-TABLET			6.32	
		ADVENT-625mg TABLET			188.81	
		ADVENT FORTE 457mg-30ML			152.50	
		AUGMENTIN-375mg TABLET			229.00	
		CLAVAM-1.2mg INJECTION			127.62	
5	AMOXYCILIN & DICLOXACILLIN	NOVACLOX	Capsule	Amoxycillin Adult: 250, 500 mg Paediatric dose: 20- 40 mg/kg/day Dicloxacillin Adult 250 mg upto 2 years : 62.5 mg 2 to 10 yrs : 125 mg	53.50	Diarrhea, Alteration of bacterial flora, rashes, HS reactions
6	AMPICILLIN	ROSCILLIN-500MG AMPICILLIN - 500 MG	Capsules, Injection	Adult dose : 500 mg Paediatric dose: 100 - 400mg/kg/ day in 4 divided doses	43.96 15.52	Diarrhea, Alteration of bacterial flora, rashes, HS reactions
7	AMPICILLIN & SULBACTUM	BACTOCILLIN-1.5mg INJECTION	Injection	Adult dose: Ampicillin 1g+sulbactam 0.5g 6 hrly by deep i.m or slow i.v over 10-15 min or i.v infusion over 15-30 min. Max dose; 12 g / day Children: >1year: 300 mg /kg/day (ampicillin 200 mg and sulbacatm 100 mg) in divided doses 6 hrly by i.v infusion	180.00	Nausea, vomiting, stomach pain, bloating, gas ,vaginal itching or discharge ,headache
8	AMPICILLIN + CLOXACILLIN	AMPILOX 500MG	Capsules, Injection	Paediatric dose: upto 2 years : 62.5mg amp & 62.5 mg clox 4-6 hrly 2-10 years : 125mg amp + 125mg clox 4-6 hrly	16.90	Diarrhea, Alteration of bacterial flora, rashes, HS reactions
		AMPILOX 500MG			60.00	
		AMPOXIN - 500MG			15.90	

9	AZITHROMYCIN	AZIBEST- 500 MG	Tablets, Syrup, Injection	Adult dose: 500 mg OD on day 1 then 250 mg OD Paediatric dose: 10mg/kg/day on day 1 then 5mg/kg/day	65.00	Mild GI upset, Headache & dizziness
		AZEE - 500 MG			203.34	
		AZITHRAL 15 ML			26.85	
		AZITHRAL 200 MG			63.40	
		AZIWOK 250MG			71.38	
		AZIWOK 500MG			118.91	
		AZITHROMYCIN-100mg TABLET			120.00	
10	CEFOPERAZONE & SULBACTUM INJ	CEBANEX - 1 GM	Injection	Adult Dose:1 to 4 g bd/ tid upto 12g/ day Paed Dose:100-150mg /kg/day bd/ tid	106.00	GI upset, Skin rash, Perineal pruritis, Depressed Haematopoiesis, HS reactions, Disulfiram like reactions
		LARICEF - 1.5GM			29.29	
		CEBANEX FORTE-1.5gm INJECTIO				
11	CEFAZOLIN SODIUM	REFLIN - 1GM	Injection	Adult: 0.25-0.5g tds deep im or slow iv over 3-5 min, Max dose: 12g/day Children: 25-50 mg/kg/day in 3-4 divided doses by deep im or slow iv over 3-5 min Max dose: 100 mg/kg/day	30.35	GI upset, Skin rash, Perineal pruritis, Depressed Haematopoiesis, HS reactions,
12	CEFEPIME HCL + TAZOBACTAM	AUGEOZ-1GM	Injection	Adult: 0.5 - 2gIV BD Paed: 50 mg/kg IV BD	197.00	GI upset, Skin rash, Perineal pruritis, Depressed Haematopoiesis, HS reactions,
		NOVAPIME - 1 GM			74.77	
13	CEFIXIME	CEFTAS CV - 200 MG	Tablet Syrup	Adult: 200 mg BD Paed: 8mg/kg in 2 divided doses	170.00	GI upset, Skin rash, Perineal pruritis, Depressed Haematopoiesis, HS reactions,
		MAHACEF - 200 MG			37.40	
		MAHACEF			59.00	
		TOPCEF			49.50	
		OMNIX-50MG - 30ML			71.00	
		OMNIX-100MG - 30ML			50.29	
		CEFTAS - 100 MG			50.00	
		CEFTAS - 200 MG			74.80	
14	CEFODROXYL		Tablet	Adult: 500 - 1000 mg OD/BD Paed: 30mg/kg/day in 2 divided doses		GI upset, Skin rash, Perineal pruritis, Depressed Haematopoiesis, HS reactions,
		DROXYL - 500 MG			23.57	
		DROXYL - 250 MG			25.00	
15	CEFOTAXIME SODIUM	TAX-O-BID - 1 GM	Injection	Adult: IM/IV 1-2g BD - QID Paed: IM/IV: 100-150 mg/kg/ day in 2-4 divided doses	16.54	Inflammation at injection site, GI upset, Skin rash, Perineal pruritis, Depressed haematopoiesis, HS reactions
		TAXIM - 250 MG			21.83	
		TAXIM - 500 MG			41.00	
16	CEFPODOXIME PROXETIL	CEFOPRAX - 100MG	Tablet Suspension	Adult:100 -200 mg BD Paed: 5 mg/kg BD	127.00	GI upset, Skin rash, Perineal pruritis, Depressed haematopoiesis, HS reactions
		CEPODEM-100MG/5ML - 30ML			139.00	
		CEPODEM -200MG			139.00	
	CEFPODOXIME +	KEFPOD CV - 200MG	24.40	Thrombocytopenia,		

	POTASSIUM CLAVULANATE	CEPODEM -XP--325 MG			100.00	Eosinophilia, Leukopenia
17	CEFTAZIDIME	ORZID - 1GM	Injection	Adult :IM/IV 1-2 g OD Paed: IM/IV 50-100 mg /kg/day	188.90	GI upset, Skin rash, Perineal pruritis, Depressed haematopoiesis, HS reactions Thrombocytopenia, Eosinophilia, Leukopenia
		FORTUM-1gm INJECTION			212.70	
18	CEFTRIAZONE	CEFOGRAM - 1 GM	Injection	Adult :IM/IV 1-2 g OD Paed: IM/IV 50-100 mg /kg/day	58.00	GI upset, Skin rash, Perineal pruritis, Depressed haematopoiesis, HS reactions Thrombocytopenia, Eosinophilia, Leukopenia
		XONE - 1GM			55.50	
		XONE - 500 MG			139.00	
	CEFTRIAZONE + SULBACTUM INJ	OFRAMAX FORTE - 1.5GM			58.05	
19	CEFUROXIME AXETIL	CEFTUM - 250 MG	Tablet	Adult dose:125-500mg	345.00	GI upset, Skin rash, Perineal pruritis, Depressed haematopoiesis, HS reactions Thrombocytopenia, Eosinophilia, Leukopenia
		CEFTUM - 500 MG			129.00	
20	CEFUROXIME SODIUM	SPIZEF - 750 MG	Injection	Adult dose: 750 mg-1.5g 8 hrly Paed: 5-0- 100 mg/kg/day in 3 divided doses	134.00	GI upset, Skin rash, Perineal pruritis, Depressed haematopoiesis, HS reactions Thrombocytopenia, Eosinophilia, Leukopenia
21	CEPHALEXIN	SPORIDEX - 125 MG 30 ML	Capsule	Adult: 250 -500mg QID Paed: 25 - 100 mg/kg/day in 3 divided doses	134.79	GI upset, Skin rash, Perineal pruritis, Depressed haematopoiesis, HS reactions Thrombocytopenia, Eosinophilia, Leukopenia
		SPORIDEX - 500 MG	Suspension		60.71	
22	CHLORAMPHENECOL	CHLOROMYCETIN - 500MG	Tablet, Capsule, Drops, Ointment	Adult: 250 -500 mg 6 hrly Paed: 25-50mg/kg/day	101.90	Swelling, Dermatitis, pruritis in the ear
		OTOCIN			39.60	
		CANDIBIOTIC EAR			62.00	
		CD - 5GM EYE			52.00	
23	CIPROFLOXACIN	CIPLOX-TZ	Tablet & Injection	Adult dose:250-500mg B.I.D	106.54	Bone marrow depression, HS reaction, Grey baby syndrome, Superinfections
		CIFRAN - 500 MG			42.03	
		ZOXAN - 500 MG			40.50	
		CIPROFLOXACIN I.V - 100 ML			17.85	
		CIFRAN-250mg TABLET			21.16	
		CEBRAN E/E DROPS-5ml			9.90	

24	CLARITHROMYCIN	CLARIMIN-500MG	Tablet, Injection	Adult dose: 250 - 500 mg B.I.D Paed dose: 7.5 - 15 mg/kg every 12 hrs	869.00	GI upset, Taste perversion, Steven Johnson syndrome, anxiety, dizziness, insomnia and confusion
		CLARIBID - 500 MG			482.90	
25	CLINDAMYCIN	CLINDATEC 300MG - 2ML	Injection, gel, capsule	Adult dose: 600 -1800 mg day in four divided doses i.m /i.v: 300 - 2400 mg/day in 2-4 divided doses child <1 mth: 15-20mg/kg /day in 3-4 divided doses child>1mth: 20-40mg/kg/ day in 3-4 divided doses	92.95	GI upset, pseudomembranous colitis, angioneurotic edema, hypotension & cardiac arrest after rapid i.v.
		DALACIN - C - 300 MG CAPSULE			394.00	
		FACECLIN-A - 15GM GEL			173.00	
		FACECLIN - 20 GM GEL			126.00	
		CLINDAC-A-15gm GEL	GEL	137.50		
26	CLINDAMYCIN + CLOTRIMAZOLE	CLINGEN-TABLET(7s)	Suppositories	Vaginal suppository 150-300 mg every 6 hours; for severe infections dose : 300-450 mg every 6 hours	115.00	Vaginal/urethral burning/itching/pain, or lower abdominal crampsmay occur.
27	DOXYCYCLINE	DOXYCYCLINE - 100 MG DOXY - LDR	Capsule	Adult dose: 100 mg OD children > 8 yrs : 2-5mg/kg/ day in 1 to 2 divided doses	35.00	Anorexia, GI upset, photosensitivity, exfoliative dermatitis, superinfections, tooth discolouration, liver damage
28	ERYTHROMYCIN	ERYTHROCIN - 500 MG	Tablet	Adult dose: 250 to 500 mg QID max 4g/day Paed dose: 20 - 40 mg/kg/ day in four divided doses	56.68	Mild GI upset, headache & dizziness
29	GATIFLOXACIN	GATE - 5ML EYE DROPS	Eye drops & eye ointment	Adult and child (above 1 year): Instill one drop 2 hrly into affected eye while awake upto 8 times daily for first 2 days, then reduce to 1 drop qid while	52.00	Eye discharge, Itching eye, Tearing
		GATIBLU - 5ML EYE DROPS			31.00	
		GATILOX - 0.3 % - 5 ML E/DROPS			54.00	
		HI GATI - 5GM EYE OINTMENT			51.00	
30	GENTAMICIN	GENTICYN 80 MG - 2ML	Injection	Adult: 1-1.7 mg/kg every 8 hrs Paed 2-2.5 mg/kg/day in 3 divided doses	4.87	Ototoxicity, nephrotoxicity & neuromuscular blockade
31	IMIPENEM & CILASTATIN	PRIMAPEN - 500 MG	Injection	Adult: 250-500 mgIV infusion 6 hrly Paed: 25 mg/kg IV infusio 8 hrly	866.00	Nausea, vomiting, diarrhea, fever, hypotension, pruritis, utricaria, rashes

32	LEVOFLOXACIN	LEON - 750 MG	Tablet, injection	Adult dose : 500 mg OD	76.00	Dizziness, drowsiness, disorientation, hot & cold flushes, slurring of speech
		LEVOFLOX - 100 ML			118.00	
		LEVOFLOX - 500 MG			103.00	
33	MEROPENEM	EUPEN - 1GM	Injection	Adult: I-3g TID Paed: 60-120 mg/kg/day in 3 doses	1061.00	Diarrhea, nausea, vomiting, GI haemorrhage, fever, chills, tachycardia, hypertension, hypotension, MI, Dyspepsia, Hypochromic anaemia
		MEROMER - 500MG			699.00	
		EUPEN-500gm INJECTION(MEROPENEM)			512.00	
34	MOXIFLOXACIN HCL	MOXAM - 400MG TABLET	Tablets, Injection, Drops, Ointment	Adult: 400mg OD	111.50	QT prolongation, phototoxicity, GI upset
		MOXAM - INJECTION			217.50	
		MOXIBLU - 5ML EYE DROPS			59.00	
		ONEMOX - 3ML DROPS			65.50	
		MOXIBLU-KT-5ML DROPS			72.00	
		APDROPS- 5ML			85.00	
		MOXAP-EYE DROPS			98.00	
		MOXICIP-EYE OINMENT			64.00	
		MOXIGRAM-LX-5ml DROPS			142.00	
35	NADIFLOXACIN	NADIBACT - 10GM CREAM	Cream	Nadifloxacin cream 1 % w/w: Applied to the lesion two times a day till the lesion resolved	63.50	Sweating, Dryness of skin
		NADOXIN CREAM - 10 GM			84.50	
36	NEOMYCIN	NEOSPORIN - H-5GM OINTMENT	Ointment	Adult and children (>2 years): Clean the affected area and apply ointment bd	44.50	Nausea, Diarrhoea, Vomiting
		NEOSPIRIN EYE OINTMENT			40.00	
37	NORFLOXACIN	NORFLOX - 400MG	Tablet	Adult: 400 mg BD	48.50	QT prolongation, phototoxicity, GI upset
		NORFLOX-TZ			66.80	
38	OFLOXACIN	OFLER - 200 MG	Tablet, Injection, Drops, Syrup	Adult: 200-800 mg OD	52.78	Abd pain, Diarrhoea, Nausea, Vomiting
		OFLIN - 100 ML			99.83	
		OFLOX - 100 ML			94.00	
		OFLOX - 200 MG			56.69	
		OFLOX - E/E DROPS			23.50	
		ZENFLOX - 60 ML			40.90	
		OFLACIN-DX - 10ML			11.68	
		OTOCIN-O - 5 ML DROPS			39.60	
		OFLOXACIN +TINIDAZOL			OFLOX - TZ TABLETS	
OFLOXACIN +ORNIDAZOL	ZANOCIN - OZ - 100MLI.V	151.00				
39	PENCILLIN G POTASSIUM	PENTIDS - 400 MG	Tablet	Adult: 400 mg	6.24	Diarrhea, alteration of bacterial flora, rashes, HS reactions

40	PIPERACILLIN & TAZOBACTAM	PIPZO - 2.25MG	Injection	Adult: 2.25- 4.5 g IV 6 hrly	154.50	GI disturbance, hypokalemia, hyponatremia, platelet dysfunction
		TAZAR-1.125GM			98.50	
		ZOPERCIN - 4.5GM			375.00	
41	ROXITHROMYCIN	ROXID - 150 MG	Tablet	Adult: 150-300 mg BD Paed: 2.5 - 5 mg/kg BD	102.32	Mild GI upset, headache & dizziness
42	VANCOMYCIN HCL	VANKING - 1 GM	Injection	Adult: 1g IV 12 hrly	591.00	Allergic reaction, Rash, Itching, Decrease in blood pressure, Breathlessness, Flushing
		VANKING - 500MG			379.00	
43	CLOFAZIMINE	HANSEPRAN - 100MG	Capsule, Tablet	Multibacillary leprosy 300 mg (w/ rifampicin 600 mg) once mthly combined w/50mg (w/ dapsone 100 mg) daily for 12 mth. Erythema nodosum leprosum (Type 2) Treatment depends on severity. 100-200 mg/day	25.72	Red-brownish black discolouration of skin especially areas exposed to sunlight, hair, sweat, sputum, urine, faeces. Rash, pruritus, photosensitivity
		CLOFROS-100mg TABLET			23.52	
44	DOXYCYCLINE	DOXY-1 LDR	Capsule	200 mg on day 1 as a single or in divided doses, followed by 100 mg/day.	65.25	Hypotension, Pericarditis, Angioneurotic oedema, Dyspnoea, Serum sickness, Peripheral oedema, Tachycardia, Urticaria, Haemolytic anaemia, Thrombocytopenia, Neutropenia
45	ETHAMBUTOL HCL & INH&PYRAZINAMIDE	AKT -4	Tablet	Adult: PO Each tab contains rifampicin 120 mg, isoniazid 50 mg and pyrazinamide 300 mg: <40 kg: 3 tab/day; 40-49 kg: 4 tab/day; 50-64 kg: 5 tab/day; ≥65 kg: 6 tab/day.	14.50	GI symptoms, pseudomembranous colitis; flushing and itching w/ or w/o rash, urticaria, pruritus, erythema, rash; hepatitis; thrombocytopenia w/ or w/o purpura, eosinophilia, leucopenia, oedema, muscle weakness, loss of tendon reflexes, paraesthesia, myopathy; vertigo, polyneuritis; occasional menstrual cycle disturbances. Flu-syndrome
		AKT-2-TABLET			67.50	
		AKURIT-2 TABLET			21.70	
		AKURIT-3 TABLET			39.00	
		AKURIT-4 TABLET			51.50	
	ETHAMBUTOL+ISONIAZID	FORECOX-150mg TABLET	36.90			
46	FAROPENEM SODIUM	DUONEM ER - 300MG	Tablet, Capsule	150 – 200 mg, thrice daily	473.50	Diarrhea, nausea, vomiting, GI haemorrhage, fever, chills, tachycardia, hypertension, hypotension, MI, Dyspepsia, Hypochromic anaemia
		FAROBACT - 200 MG			332.50	

47	LINEZOLID	LINOSPAN - 600 MG	Tablet, Syrup, Injection	Adult: PO 400 mg 12 hrly PO/IV600 mg 12 hrly	156.00	Reversible myelosuppression (e.g. leucopenia, thrombocytopenia, anaemia, pancytopenia); headache, fever, convulsions and seizures, rashes, dizziness, nausea and vomiting, metallic taste, insomnia, constipation, diarrhoea, abdominal pain, skin disorder, cough, pharyngitis
		LIZOFORCE - 30ML			108.77	
		LIZOLID 300 ML			382.05	
		LINID-TABLET(LINEZOLID)			347.00	
48	MUPIROCIN	SUPIROCIN - 5GM	Ointment	Adult and children >2 month: 2% ointment applied on affected area tds for 5-10 days	95.00	Burning, stinging, pain, itching, rash, redness, dryness, tenderness, or swelling of the treated skin.
49	MUPIROCIN+METRONIDAZOLE NITROFURANTOIN	CO-MUPIMET-5ml COLLAGENS	Collagens	Adults: Clean the affected area and applied to the lesion three times a day for 10 days(bacterial infection)	450.00	Dry skin, Application site reactions (burning, irritation, itching and redness).
50	NITROFURANTOIN	URIFAST-100MG	Tablet	50-100 mg 4 times/day	123.90	Nausea, vomiting, anorexia, abdominal pain, diarrhoea; headache, drowsiness, vertigo, dizziness, nystagmus, benign intracranial hypertension; rash, urticaria
51	POVIDONE -IODINE	WOKADINE	Ointment, Pessaries, Powder, Gargle, Solution	Topical: Clean affected area. Apply small amount topically to affected area 1—3 times daily. May cover with sterile bandage. Do not use for longer than 1 week. Pessary: one pessary for a period of 7 days. Complete Adult	82.00	Skin irritation, Severe allergic reaction
		BETADINE GARGLE - 50 ML			83.50	
		BETADINE - V - PESSARIES			105.50	
		BETADINE SOLUTION 10% - 100ML			106.05	
		DREZ - S - 10 GM			53.24	
		DREZ - S - 15 GM			77.00	
52	RETAPAMULIN	RETAREL - 5GM	Cream	Adult: Topical As 1% oint: Apply twice daily for 5 days. Total treatment area:≤100cm 2 total body surface area.	84.00	Headache, pyrexia, pruritus, eczema, diarrhoea. Application site irritation, pain, erythema, contact dermatitis.
53	RIFAMPICIN	R-CIN - 600MG	Capsule	PO 8-12 mg/kg/day. <50 kg: 450 mg/day; ≥50 kg: 600 mg/day	21.00	Facial flushing and itching, w/ or w/o a rash, flu-like syndrome characterised by episodes of fever, chills, headache, dizziness, bone pain, shortness of breath
		R-CIN-300MG			34.70	

54	RIFAXIMIN	RIFAGUT - 400MG	Tablet	200 mg orally 3 times a day	239.00	GI upset, periphaeral edema, Pyrexia, Asthenia
55	STREPTOMYCIN	AMBISTRYN - S	Injection	1-2 g/day in divided doses 6-12 hrly.	8.83	Vestibular and cochlear function disturbance, optic nerve dysfunction, peripheral neuritis, arachnoiditis, encenhalopathy
56	MUIPIROCIN+METRONIDAZOLE	MUPIMET-5gm OINTMENT	Ointment	Applied to the affected area three times daily	109.00	Mild burning or stinging
57	BECLOMETHASONE+LIDOCAINE+PHENYLEPHRINE	ANOVATE-20gm CREAM	Cream	Applied to the affected area once or twice daily	91.75	Application site irritation.
58	OFLOXACIN, BECLOMETASONE, CLOTRIMAZOLE.	MYCLIN-O-EAR DROPS	Drops	1 to 2 drops in the affected ear 2 to 3 times a day.	32.90	Irritation.
59	TRICLOSAN	SOAPEX CLASSIC-75gm SOAP	Soapex	Used as bathing soap daily for antibacterial action. Wet the face and body with water Apply the soap and gently work up the lather.Rinse thoroughly	113.20	
60	DERMADEW SOAP SOAP: COCOA BUTTER , COCONUT OIL , PALM OIL , GLYCERIN , ALOE VERA , OLIVE FRUIT EXTRACT.	DERMADEW-75gm SOAP	Soap	Wet the face or body with water Apply Dermadew soap Gently work up the lather Rinse thoroughly	142.00	Drowsiness, dizziness, hypotension or a headache
61	PALM & COCONUT OIL, SHEA BUTTER, ALOE VERA	GLADDEW BABY-75gm SOAP	Soap	Wet the face or body with water Apply the soap and gently work up the lather. Rinse thoroughly	109.00	Drowsiness, dizziness, hypotension or a headache
		GLADDEW-75gm SOAP	Soap		105.00	
62	POVIDONE IODINE (10% W/W)	WOKADINE-15gm OINTMENT	Ointment	Clean affected area. Apply small amount topically to affected area 1—3 times daily. May cover with sterile bandage. Do not use for longer than 1 week.	98.50	No common side effects seen.
63	POVIDONE IODINE (7.5% W/V)	WOKADINE-500ml SOLUTION	Solution	Clean affected area. Apply small amount topically to affected area 1—3 times daily. May cover with sterile bandage. Do not use for longer than 1 week.	201.60	Skin irritation, Allergic Skin Reaction, Acneiform eruptions, Thyroid imbalances

64	POVIDONE IODINE	PLUS SPRAY-50gm	SPRAY	clean the affected area and spray a small amount of the drug 1 to 3 times daily	79.00	Swelling, nausea, headache, dizziness, vomiting, constipation, pruritus, pain in extremity.
65	NEOMYCIN BECLOMETASONE AND LIGNOCAINE	OTIFLOX NEW-5ml EAR DROPS	Drops	Apply on the affected area of the skin after cleaning and drying it well. It is usually recommended to use about 2-3 times a day.	58.91	Ear discomfort, Application site irritation.
66	CARICA PAPAYA	POP-E-TABLET	Tablet	one tablet containing 1.1g of carica papaya should be taken three times a day, for five days.	300.00	Severe stomach/abdominal pain, nausea/vomiting, slow heartbeat, severe drowsiness, inability to move.
67	SILVER NITRATE	SILVEREX-20gm GEL	Gel	Apply the gel on the wound/ affected area at least once daily	99.00	Nausea, vomiting, diarrhoea; hypersensitivity, skin reactions; haematuria; crystalluria; thrombocytopenia, leucopenia, eosinophilia.
68	DORZOLAMIDE HYDROCHLORIDE (DORZOLAMIDE HCL), A CARBONIC ANHYDRASE	DORZOX-5ml EYE DROPS(DORZOLOMIDE)	Drops	1 droplet into each affected eye, 1 - 3 times per day.	360.50	Blurred vision, cloudy vision, double vision, drooping eyelid
69	BENZALKONIUM CHLORIDE+ZINC OXIDE	HAPPY NAP-20gm CREAM	Cream	Put a thin layer on the affected skin(diaper rash)1-3 times a day. Do not wash the treated area after immediately applying Happynap Cream.	92.00	Hives; difficulty breathing; swelling of your face, lips, tongue, or throat.
70	SULPHONATED SURFACTANT	ACNE AID-100gm BAR	Soap	Make a lather with warm water and massage into the affected skin using the finger tips or a facecloth. Rinse and repeat as necessary.	125.80	Peeling, itching, irritation, and reddened
71	TEA TREE OIL, ALOE VERA, A	ACNESTAL-75gm SOAP	Soap	Apply soap on face and shoulder areas Leave it for a minute Wash with water and gently dry the area Can be used for 2- 3 times a day.	88.00	Peeling, itching, irritation, and reddened
72	SALICYLIC ACID TOPICAL,SULFUR TOPICAL.	ACNIL SOAP	Soap	Adults: .Wet the face or body with water Apply the soap and gently work up the lather. Rinse thoroughly. Can be used for 2- 3 times a day.	60.00	Skin irritation, Inflammation of skin, Rash, Contact dermatitis, Irritation to the gastrointestinal tract, Nausea, Vomiting, Diarrhea.

73	CHLORHEXIDINE GLUCONATE	CLOHEX ADS-200ml MOUTH WASH	Mouthwash	Swish for 30 seconds to 1 minute and spit it. Do not swallow. Avoid eating, drinking, or smoking for at least 30 minutes after rinsing.	149.00	Respiratory tract infectionsevere, Skin peelingsevererare, Hypersensitivitysevererare, Inflammation of gumsevere, Tooth pain, Dry mouth, Staining of tooth, Altered sense of taste.
----	-------------------------	-----------------------------	-----------	---	--------	---

ANTI-CANCER

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	CYLOPHOSPHAMIDE	PHOSMID - 1GM	Injection	Malignancies Low dose regimen: 2-6 mg/kg/wk in divided doses. IV Malignancies Moderate dose regimen: 10-15 mg/kg /wk; high dose regimen: 20-40 mg/kg	79.00	Congestive heart failure; leucopenia poor wound healing; anorexia. Nausea, vomiting; alopecia; oral mucosal ulceration; anemia thrombocytopenia, nonhaemorrhagic cystitis.
2	DOXORUBICIN	METODOX-50 -50MG	Injection	AIDS-related Kaposi's sarcoma As pegylated liposome: 20 mg/m2 once 2-3 wkly. Ovarian carcinoma As pegylated liposome: 50 mg/m2 once 4 wkly. Metastatic breast carcinoma w/ cyclophosphamide: 60-75 mg/m2 once 3 wkly.	304.00	Leucopenia, thrombocytopenia, nausea, vomiting, diarrhoea. Rarely facial flushing, rash, alopecia. Blurred vision, headache, seizures, paraesthesia, confusion, malaise, lethargy, skin pigmentation.
3	FLUOROURACIL	CHEMOFLURA - 500MG-10ML	Injection	12 mg/kg/day (max: 0.8- 1g/day) for 3-4 days	21.60	Leucopenia, thrombocytopenia, stomatitis, GI ulceration, bleeding and diarrhoea, haemorrhage from any site (stop treatment)
		FLURACIL - 250 MG			11.61	
4	METHOTREXATE	FOLITRAX - 5 MG	Injection & Tablet	Tab: 2.5 - 30 mg/day IV 10-60 mg/m2	38.27	Ulceration of the mouth and GI disturbances (e.g. stomatitis and diarrhoea), bone marrow depression, hepatotoxicity, renal failure, skin reactions, alopecia, ocular irritation, arachnoiditis
		FOLITRAX 15MG -1ML			38.50	
		ONCOTREX - 2.5MG			54.64	
		ONCOTREX - 5MG			84.80	
5	MITOMYCIN -C & SODIUM CHLORIDE	MITOMYCIN -C - 2MG	Injection	4-10 mg (0.06-0.15 mg/kg)	148.50	GI toxicity, dermatitis, alopecia, fever, malaise; local tissue necrosis, ulceration, cellulitis, cholecystitis, cholangitis, bile duct necrosis; pain, redness, erythema, blisters, erosion, ulceration, skin/muscle necrosis at inj site. Rarely, cardiotoxicity.

6	MYCOPHENOLATE MOFETIL	BIOMYF-500MG	Tablet	1- 1.5 g twice daily	750.00	Diarrhoea, vomiting, GI haemorrhage and perforation; leucopenia; asthenia pain, headache, anaemia, thrombocytopenia, renal tubular necrosis
7	PALONOSETRON (75MG)	THEMISSET-0.75% INJECTION	Injection	Adult: 250mcg by bolus inj over 30 sec to be given approx. 30min before chemotherapy Children (1 month - 17yrs): 20mcg/kg (max 1,500mcg) by infusion over 15 mins to be given approx 30min before chemotherapy	163.90	Headache, Dizziness, Constipation, Diarrhoea.
8	CISPLATIN	CISMAX-10mg/10ml INJECTION	Injection	Administered by slow iv infusion 50-100mg/m ² BSA every 3-4 weeks	70.00	Acute renal failure, Tinnitus, Hearing loss, Leukopenia, Fever, Infection, Nausea, Vomiting, Hypocalcemia
		CISMAX-50mg/50ml INJECTION	Injection		350.00	
9	CARBOPLATIN	CABOO-150mg/15ml INJECTION	Injection	Adult (Previously untreated patient): 400mg/m ² as single short term IV infusion over 15-60min. If previously treated reduce initial dosage by 20-25% (300-320mg/m ²)	809.90	Pain at the injection site severe, Low WBC and Platelet count severe, Fever and chills severe.
		CABOO-450mg INJECTION*	Injection		2222.00	
10	MESNA	CANCENA-200mg/2ml	Injection	Adult and children >4 month: 240mg/m ² (if receiving 1.2g/m ² ifosfamide dose) or 20% of cyclophosphamide dose administered either by short (15-30 mins) or continuous (24 hrs) infusion	26.85	Nausea, Headache, Lightheadedness, Lethargy, Abdominal pain, Flushing, Diarrhoea, Rash, Fever, Flu-like symptoms.
11	CARBOPLATIN	CARBOMAX-450mg INJECTION	Injection	Adult (Previously untreated patient): 400mg/m ² as single short term IV infusion over 15-60min. If previously treated reduce initial dosage by 20-25% (300-320mg/m ²)	2433.00	Loss of appetite, stomach pain, diarrhea, constipation, nausea and vomiting, mouth blistering.
12	FLUOROURACIL	CHEMOFLURA-250mg INJECTION	Injection	Adult: 12mg/kg/day (max 0.8-1g/day) for 3-4 days, maintenance dose: 5-15mg/kg/wk (max 1g/wk. As an infusion, 15mg/kg/day (max. 1g daily) in 500ml NS or 5% glucose over 4 hrs	10.45	Nausea, vomiting, loss of appetite, stomach ache, diarrhea, or darkening of the skin/nails

13	EPIRUBICIN	EPINEON 10mg-INJECTION	Injection	Dose:60-90mg/m2 iv over 5 min, repeated at 3 weeks, maximum dose < 900mg/m2	244.00	Rash, Vomiting, Weakness, Conjunctivitis, Nausea, Mouth sore, Hair loss, Absence of menstrual periods, Itching, Diarrhoea, Hot flashes.
		EPINEON 50mg-INJECTION	Injection		982.00	
14	ETOPOSIDE	ETOVEL-100mg INJECTION	Injection	Dose:50-100mg/m2/day iv for 5 days	195.00	Vomiting, Weakness, Orthostatic hypotension, Allergic reaction, Nausea, Loss of appetite, Hair loss,
		ETOVEL-50mg CAPSULE	Capsule	100-200mg/day oral	560.00	
15	FILGRASTIM	FILNEON-300mg/0.5ml INJECTION	Injection	Adult:5-10mcg/kg daily as a single daily SC inj, as a continuous IV or SC infusion, or as a daily IV infusion over 15-30 mins	1186.65	Bone pain, Weakness, Headache, Rash, Nausea, Increased lactate dehydrogenase level in blood, Vomiting,
16	GEFITINIB	GEFTIGET-250mg TABLET	Tablet	Adult: 250mg once daily	4355.00	Nausea, Rash, Vomiting, Weakness, Loss of appetite, Dry skin, Diarrhoea, Stomatitis
		GEFTINAT-250mg TABLET	Tablet		5900.00	
17	GEMCITABINE	GEMCHEM-1000mg INJECTION	Injection	Adult: 1,000 mg/m2 once weekly via infusion over 30 mins for 3-7 week and then followed by 1 week rest. This cycle is again repeated	1300.75	Nausea, Vomiting, Loss of appetite, Hair loss, Low blood platelets, Allergic reaction, Breathlessness, Decreased white blood cell count, Increased liver enzymes, Anemia
		GEMCHEM-200mg INJECTION	Injection		402.00	
		GEMGET-1000mg INJECTION	Injection		5165.00	
		GEMGET-200mg INJECTION	Injection		1170.00	
18	IFOSFAMIDE&MESNA	HOLOXAN 1gm+UROMITEXAN	Injection	1.2 g/m2 per day for 5 consecutive days. Treatment is repeated every 3 weeks.	463.22	Hair loss, Nausea, Vomiting, Anemia, Blood in urine, Decreased white blood cell count, Increased risk of infection, Neurotoxicity.
		HOLOXAN 2gm+UROMITEXAN	Injection		1157.96	
19	LEUPROLIDE / LEUPRORELIN	LUPRODEX-3m(11.25)INJECTION	Injection	Adult: 1mg as single daily dose by SC inj.	#####	Decreased libido, Testicular atrophy, Increased sweating, Fatigue, Muscle weakness, Erectile dysfunction, Bone pain, Hot flashes
20	METHORTREXATE	METHOFAST-50mg/2ml INJECTION	Injection	Adult: 15-30mg daily for 5 days	38.00	Nausea, vomiting, stomach pain, drowsiness, or dizziness
21	DOXORUBICIN	METODOX-10mg INJECTION	Injection	Adult: 60-75 mg/m2 BSA once every 3 week	90.20	Vomiting, Allergic reaction, Nausea, Mouth sore, Hair loss, Stomach pain, Diarrhoea, Decreased appetite, Eye disorder, Increased thirst, Decreased white blood cell count (neutrophils), Heart beat irregular, Nail disorder.
22	PEGFILGRASTIM	PEGNEON-6mg/0.6ml-INJECTION	Injection	Adult: 6 mg subcutaneously once a week for 2 doses , Child: Less than 10 kg: 0.1 mg/kg 10 to 20 kg: 1.5 mg 21 to 30 kg: 2.5 mg 31 to 44 kg: 4 mg	7200.00	sudden or severe pain in your left upper stomach spreading up to your shoulder;

23	OXALIPLATIN	OXYREA-100mg INJECTION	Injection	Adult:85mg/m ² every 2 wk,given via infusion over 2-6 wk for 12 cycles	1376.00	Decrease in platelet counts, nausea, vomiting and diarrhea,constipation and abdominal pain.
		OXYREA-50mg INJECTION	Injection		751.00	
24	PACLITAXEL	PACLIGET-260mg INJECTION	Injection	Adult:135mg/m ² infused over 24 hr and repeat at 3 weeks intervals.	9000.00	Difficult or painful urination, Dizziness when getting up suddenly from a sitting position, Fever and chills.
25	PALANOSETRON	PALOCHEM 0.25mg/5ml INJECTION	Injection	Adult: 250mcg by bolus inj over 30 sec to be given approx.30min before chemotherapy Children(1 month - 17yrs):20mcg/kg(max 1,500mcg) by infusion over 15 mins to be given approx 30min before chemotherapy	72.00	headache, problems with bowel movement, chills, decreased urge to urinate, acidity, drowsiness, drying of the mouth, cough, bloating.
26	LOMUSTINE	BELUSTINE-40mg CAPSULES	Capsule	single dose(40mg) every 6 to 8 weeks	1125.00	headache, stomach pain, nausea, diarrhea, vomiting, gas.
27	DOCETAXEL	DOCAX -120mg/3ml INJECTION	Injection	Adult: 60-100mg/m ² by infusion over 1 hr every 3 wk	2756.00	Nerve damage, Fluid retention, Muscle pain, Vomiting, Weakness, Nausea, Infection, Difficulty in swallowing, Loss of appetite, Hair loss, Diarrhoea, Skin reaction, Breathing difficulty.
28	IRINOTECAN	INTENSIC-100mg/5ml INJECTION	Injection	Adult:125 mg/m ² infused IV over 90 mins once weekly for 4-6 weeks	781.00	Fatigue, Nausea, Vomiting, Weakness, Hair loss, Fever, Anemia, Diarrhoea, Decreased white blood cell count, Decreased appetite, Mucosal
		INTENSIC-40mg/2ml INJECTION	Injection		442.00	
29	IFOSFAMIDE	IPOGET-1gm INJECTION	Injection	Adult: 1.2g/m ² /day for 5 days (slow iv infusion for 30 mins)repeat every 3 week	493.00	Nausea, Vomiting, Anemia (low number of red blood cells), Hair loss, Decreased white blood cell count, Infection, Blood in urine, Cns toxicity.
30	PACLITAXEL	TAXELEON-100mg-16.5ml/7ml INJE	Injection	Adult:135mg/m ² infused over 24 hr,175mg/m ³ infused over 3 hr	618.00	Nausea, Vomiting, Weakness, Rash, Low blood platelets, Upper respiratory tract infection, Urinary tract infection, Bleeding, Anemia.
		TAXELEON-260mg/43.34ml INJECTI	Injection		1536.00	
31	ZOLEDRONIC	ZOLEPHOS-5mg/100ml INJECTION	Injection	Adult: 4-5mg as a single infusion over at least 15 min.Retreatment after atleast 7 days if necessary.	3000.00	Headache, Back pain, Musculoskeletal (bone, muscle or joint) pain, Indigestion, Heartburn, Diarrhoea.
32	TAMOXIFEN	ENTAX-20mg TABLET	Tablet	20 mg given in 1-2 divided doses. Max: 80 mg/day.	28.00	Vaginal discharge, Hot flashes.
33	DONEPEZIL + MEMANTINE	DONAMEM-10mg TABLET	Tablet	10 mg tablet taken once a day with or without food, in the evening just before bedtime.	183.00	Nausea, Vomiting, Loss of appetite, Diarrhoea, Headache, Dizziness, Bleeding under the skin.

		DONAMEM-5mg TABLET	Tablet		125.00	
34	CITICOLINE (250MG)	CITIZAC-2ml INJECTION	Injection	Adult: up to 1 g IM/IV daily	99.00	Decreased blood pressure, Stomach pain, Diarrhoea, Injection site reactions (pain, swelling, redness), Irregular heart rate.

ANTI-COAGULANTS AND THROMBOLYTIC

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	APIXABAN	ELIQUIS-2.5MG	Tablet	Adult: 2.5 mg - 5 mg BID	725.00	Anemia, hemorrhage, hypotension, GI upset, CNS toxicity
2	CILOSTAZOL	PLETOZ- 50MG	Tablet	100 mg bid	77.00	dizziness, palpitations, diarrhoea, abnormal stools; pain, infection; peripheral oedema,
		STILOZ-50			97.50	
3	CLOPIDOGREL	CLOPILET - 75 MG	Tablet, Capsule	75 mg OD	71.01	Dyspepsia, abdominal pain, nausea, gastritis, vomiting, flatulence, constipation, gastric and duodenal ulcers.
		CLOPILET - A - 150MG			38.11	
		CLOPILET - A - 75MG			34.25	
		PLAVIX-75MG			147.44	
4	ENOXAPARIN SODIUM	ENCLEX - 60 MG - 0.6 ML	Injection	20-40 mg OD	552.50	Haemorrhage (including at the inj site), peripheral or unspecified oedema, anaemia, haematuria, ecchymosis, fever, confusion, nausea, diarrhoea, dyspnoea, inj site pain.
		ENOXION 40 MG			504.00	
		ENOXION-60mg-INJECTION			580.70	
		ENCLEX-40mg INJCETION			400.72	
		CLEXANE-80mg INJECTION			1069.65	
5	FONDAPARINUX SODIUM	FONDAZEST-2.5mg/0.5ml INJECTION	Injection	The usual dose is 2.5 mg once a day, injected at about the same time each day. If there is kidney disease, the dose may be reduced to 1.5 mg once a day.	820.00	Bleeding, Anemia (low number of red blood cells).
6	ERYTHROPOITINE	ZYROP-4K	Injection	600 u/kg twice wky for 3 wk before surgery	1396.00	Hypertension, myalgia, arthralgia, flu-like syndrome, rashes and urticaria.
		EPOSIS 4000IU			850.00	
7	ETHAMSYLATE	ETHAMSYLATE - 2 ML	Injection	Adult: 250-500 mg Low birth-weight neonate: 12.5 mg/kg IM/IV Inj every 4-6 hr.	23.00	Transient hypotension.
8	HAEMOCOAGULASE INJ	BOTROPASE - 1 ML	Injection		92.25	No common side effects seen.
9	HEPARIN SODIUM	THROMBOPHOB - 20 GM	Injection, Ointment	75-80 U/kg or 5,000 U then 18 U/kg/hr or 1,000-2,000 U/hr	75.00	Hypersensitivity reactions ainful, ischaemic and cyanosed limbs; osteoporosis delayed and cyanosed limbs; osteoporosis delayed transient alopecia. prianism. rebound
		LOFH-25,000 IU			237.30	
					6950.00	

						hyperlipaemia.
10	NICOUMALONE	ACITROM - 1	Tablet	4-12 mg on the 1st day and 4-8 mg on the 2nd day	150.00	Alopecia; fever, nausea, vomiting, diarrhoea;skin rash; cholestatic liver damage Haemorrhage.
		ACITROM - 2 MG			235.00	
		ACITROM - 3MG			300.00	
11	PROTAMINE SULPHATE	PROTA - 5ML	Injection	1-1.5 mg per 100 u of heparin	48.88	Hypotension, bradycardia, pulmonary and systemic hypertension, dyspnoea, warm sensation, transitory flushing, nausea and vomiting, lassitude.
12	RETEPLASE- RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR	RETELEX - 18MG KIT	Injection	10 u via slow IV.Repeat dose once, 30 mins after the start of 1st inj.	#####	Localised inj site bleeding, GI bleeding, anaemia.
13	STREPTOKINASE	ICIKINASE - 1500000 I.U	Injection	1.5 million u	2270.00	Fever, chills, back pain, abdominal pain, nausea, vomiting, arrhythmia, bruising, rash, pruritus, acute renal failure,Haemorrhage; anaphylactic shock.
14	TERLIPRESSIN	TERLIEON - 1MG/10ML	Injection	1-2 mg	1467.00	Abdominal cramps, cardiac arrhythmias, headache, transient blanching, increased arterial pressure. Potentially Fatal: MI, cardiac failure.
15	TIROFIBAN	AGGRIBLOC - 5MG -100ML	Injection	0.4 mcg/kg/min for 30 mins, then 0.1 mcg/kg/min for≥24-48 hr	5230.37	bleeding, thrombocytopenia, bradycardia, dissection of the coronary artery, dizziness, oedema, pelvic pain
16	TRANEXAMIC ACID	KLOTIN - 5ML	Injection,Tablet	1-1.5 g 2-3 times/day.	69.20	Diarrhoea, nausea, vomiting, disturbances in colour vision, giddiness, hypotension (after rapid IV inj), thromboembolic events
		TRAPIC			187.00	
	TRANEXAMIC ACID + MEFANAMIC ACID	TRAPIC - MF			211.00	
17	TRANEXAMIC ACID (100MG)	TRAPIC-5ml INJECTION	Injection	Adult: 0.5-1g or 10mg/kg tid or 25-50 mg/kg daily by continous infusion Child: 10mg/kg bid or tid	69.85	Tiredness, Injection site reactions (pain, swelling, redness), Nasal congestion, Musculoskeletal (bone, muscle or joint) pain.
18	TRANEXAMIC ACID (250MG) + ETHAMSYLATE (250MG)	TRAPIC-E-TABLET	Tablet	1 – 1.5 g three times per day.	188.00	Nausea, Stomach pain, Diarrhoea, Headache, Nasal congestion, Sinus pain, Sinus inflammation, Rash, Tiredness, Musculoskeletal (bone, muscle or joint) pain, Anemia (low number of red blood cells).

ANTI-DEPRESSANTS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMITRIPTYLINE HCL	TRIPTAZ - 10 MG	Tablets	Adult:50-200mg	21.50	GI upset, cns toxicity, heart block, arrythmias, orthostatic hypotension, palpitation, dry mouth, urinary retension, blurred vision.
		TRYPTOMER - 10 MG			43.00	
		TRYPTOMER - 25MG			36.86	
2	CLONAZEPAM AND ESCITALOPRAM	CITALIN FORTE-TABLET	Tablets	CLONAZEPAM 0.5mg+ ESCITALOPRAM 10mg combination tablet taken once daily	103.00	Nausea, Vomiting, Diarrhoea, Confusion, Memory impairment, Uncoordinated body movements, Drowsiness, Tiredness, Delayed ejaculation, Anorgasmia (decreased orgasm), Low sexual desire.
3	DULOXETINE HCL	SYMPTA - 20 MG	Tablets	30-80mg	57.50	GI upset, cns toxicity, agitation, insomnia, raise in BP
4	FLUOXETINE	FLUDAC - 20 MG	Capsules	20-50 mg	58.10	GI upset, cns toxicity, epistaxis, serotonin syndrome .
		PRODEP - 10MG			36.00	
		PRODEP - 20MG			40.17	
5	MIRTAZAPINE	MIRTAZ - 15 MG	Tablets	15-45mg	79.00	GI upset, cns toxicity, agitation, insomnia, raise in BP
6	SERTRALINE	SERTA - 25 MG	Tablets	50-200mg	38.00	GI upset, cns toxicity, epistaxis, serotonin syndrome .
		SERTA - 50 MG			70.00	
7	DOSULEPIN	PROTHIADEN 25MG	Tablets	25mg tid, gradually increased to 50 mg tid if necessary	46.00	cholestatic jaundice, sexual dysfunction, exacerbation of psychotic manifestation, cardiac arrhythmias, hypotension, dry mouth constipation
		PROTHIADEN 75MG			95.50	
8	ESCITALOPRAM OXALATE	NEXITO - 10MG	Tablets	5 - 20 mg/day	76.00	Nausea, diarrhoea, increased sweating, insomnia, impotence, ejaculation disorder, fatigue, somnolence; postural hypotension, sinusitis, taste disturbances. Increased appetite and wt gain.
		NEXITO - 5MG			45.00	
9	IMIPRAMINE HYTROCHLORIDE	ANTIDEP	Tablets	Initial: 75 mg/day, up to 150-200 mg/day if needed	8.61	Sinus tachycardia, AV/bundle-branch block, postural hypotension, dry mouth, wt loss/gain constipation, urinary hesitancy/retention, impotence
10	VENLAFAXINE ER	VENIZ XR - 75 MG	Capsules	75 mg/day in 2 or 3 divided doses	68.00	Changes in behaviour, suicidal ideation, agitation, tremor, nervousness, anxiety, insomnia, confusions, abnormal dreams, HTN, nausea, headache, asthenia,

ANTI-DIABETIC

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	GLIBENCLAMIDE	DAONIL - 5 MG	Tablet	Adult dose: start at 1.25 - 2.5 mg gradually increase upto 20 mg/day	11.11	Abdominal cramps, diarrhea, joint and muscle pain, leukopenia
	GLIBENCLAMIDE + METFORMIN	GLUCORED			24.50	
		GLUCORED FORTE			32.50	
		GLUCORED FORTE - 850 MG			45.00	
		DAONIL - M XR			38.30	
		DUOTROL			22.70	
2	GLICLAZIDE	DIAMICRON XR - 60MG	Tablet	Adult dose: 160 mg in two divided doses	152.28	Abdominal cramps, diarrhea, joint and muscle pain, leukopenia
		DIAMICRON-MR			63.39	
	GLICLAZIDE +METFORMIN	RECLIMET			120.00	
3	GLIMEPIRIDE	GP-1MG	Tablet	Adult dose: Initially 1mg once daily followed by maximum upto 6 mg	38.49	Abdominal cramps, diarrhea, joint and muscle pain, leukopenia
		GP-2MG			62.77	
		GLIMULIN - 1 MG			60.15	
		GLIMULIN - 2 MG			107.40	
		GEMER - 2 MG			81.00	
	GLIMEPIRIDE + METFORMIN	GEMER -1MG			54.00	
		GLIMESTAR-M1			39.00	
		GLIMESTAR-M2			59.40	
		GLIMULIN-MF			111.00	
		GLIMY M1			91.50	
		GLYCOMET GP - 1 MG			80.50	
		GLYCOMET-GP - 2MG			118.25	
		ISRYL-M1			22.50	
		ISRYL-M2			36.00	
		BETAGLIM - 1 MG			42.95	
GLYCOMET GP - 0.5 MG	44.85					
4	GLIPIZIDE	GLYNASE - 5 MG	Tablet	Adult dose: initially 5mg increased to the maximum of 15mg	5.07	Abdominal cramps, diarrhea, joint and muscle pain, leukopenia
		SEMI GLYNASE - 2.5MG			2.75	
	GLIPIZIDE + METFORMIN	GLYNASE - MF			11.53	
5	INSULIN ASPART	NOVORAPID FLEXPEN	Injection	0.2-0.4U/kg/day	612.65	Hypoglycemia, local reactions, allergy, edema
6	INSULIN GLARGINE	LANTUS CARTRIDGE - 3 ML	Injection	0.2U/kg or upto 10U once daily	493.60	Hypoglycemia, local reactions, allergy, edema
7	METFORMIN	GLYCIPHAGE - 500 MG	Tablet	Adult dose: 0.5 - 2.5 g in 2 divided doses	30.26	lactic acidosis, vitamin B12 deficiency

		GLYCOMET - 1 GM			59.17	
		GLYCOMET - 850MG			25.25	
		GLYCOMET SR- 500MG			36.12	
		CARBOPHAGE-XR - 500MG			18.00	
		CETAPIN - XR -500MG			33.00	
8	SITAGLIPTIN PHOSPHATE	ISTAVEL - 50MG	Tablet	25 - 100 mg OD	269.00	Allergic reaction, myalgia, GI discomfort
9	VOGLIBOSE	VOLIBO - 0.2 MG	Tablet	Adult dose: 0.2 - 0.3 mg OD	78.00	Flatulence, abdominal discomfort, loose stool
		VOLIBO - 0.3 MG			109.50	
		VOLIBO - M - 0.2 MG		84.25		
		VOLIBO - M - 0.3 MG		114.00		
10	VILDAGLIPTIN	GALVUS-50MG	Tablet	50mg tablet taken twice daily	328.00	Allergic reaction, myalgia, GI discomfort
11	BIPHASIC ISOPHANE INSULIN INJECTION	HUMAN MIXTARD 30/70	Injection	Type 1 diabetes: Ketones moderate or less: 0.5 U/kg/day SC, Ketones large: 0.7 U/kg/day	148.37	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision
12	MIXTURE OF INSULIN & PROTAMINE CRYSTALLISED	NOVOMIX FLEXPEN	Injection	0.5-1.0 IU/kg/body weight/day	611.31	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision
13	SOLUBLE INSULIN	HUMAN ACTRAPID	Injection	Adult(IM):Initial loading dose of 20 units ,followed by 6 units/hr until blood glucose	149.40	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision
14	ISOPHANE INSULIN	HUMAN INSULATARD	Injection	0.3-1.0 IU/kg/body weight/day.Given once or twice a day	149.40	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision

ANTI-EMETIC

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	DOMPERIDONE	DOMSTAL - 30 ML	Tablets, Syrup	Adult: 10-20 mg every 4-6 hours	35.60	Galactorrhea, Gynaecomastia, weakness, headache
		DOMSTAL			26.04	
	DOMPERIDONE + OMEPRAZOLE	DOMSTAL - O		Paed: 0.2-0.3mg/kg 3-4 times a day	93.00	
2	ONDANSETRON	EMESET - 30ML	Syrup, Tablets,	Adult; IV 0.15 mg/Kg, 8 mg Orally OD Paed: 4-12 yrs 4 mg OD	36.14	Headache, GI upset, constipation
		EMESET - 4MG	Injections		55.14	
		NEOMIT - 4 MG - 2 ML			16.60	
3	PROCHLOPERAZINE MALEATE	STEMETIL	Injections, Tablet	Adult: 5-10 mg TID/QID Injection: 12.5mg IM	49.40	Drowsiness, dizziness, amenorrhea blurred vision, skin reactions
		STEMETIL-MD-TABLET			65.00	

4	LEVOSULPIRIDE	NEOPRIDE	Tablet	25 mg 3 times/day	53.00	Amenorrhoea, gynaecomastia, . galactorrhoea, changes in libido
5	METOCLOPRAMIDE	PERINORM	Tablet, Injection	10-15 mg 4 times/day	4.67	Extrapyramidal symptoms (usually acute dystonic reactions); parkinsonism; tardive dyskinesia; restlessness, drowsiness, dizziness
		PERINORM			12.08	
6	PROMETHAZINE (5MG/5ML)	PHENERGAN-100ml ELIXIR	Elixir	Children(2-5yrs):5mg Children(5-10yrs):10mg Children >10 yrs and adults: 25mg To be taken in the night before the journey and then repeat after 6-8 hours as required.	28.5	Liver problems that may cause the eyes or skin to go yellow (jaundice),Muscle stiffness or shaking.
7	PROMETHAZINE (25MG)	PHENERGAN-2ml INJECTION	Injection	Adult: 125-25mg, repeated at intervals of not<4hr.(Max:100mg daily)	8.35	Drowsiness, dizziness, constipation, blurred vision, or dry mouth.
8	APREPITANT	APRINEON-125/80mg CAPSULE (APREPITANT)	Capsule	Day 1: 125 mg orally once 1 hour before chemotherapy, Days 2 and 3: 80 mg orally once a day, 1 hour before chemotherapy OR in the morning (if chemotherapy is not given on Days 2 and 3), Duration of therapy: 3 days/cycle	1260	nausea, vomiting, heartburn, stomach pain, diarrhea or constipation, loss of appetite, hiccups, hair loss, headache, dizziness, tired feeling.

ANTI-EPILEPTICS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	CARBAMAZEPINE	TEGRITAL - 200MG	Tablet, syrup	Adult: 200- 400 mg TDS Paed: 15 -30 mg/kg/day	15.90	sedation, dizziness, vertigo, diplopia, ataxia
		TEGRITAL -100ML			15.24	
		TEGRITAL -400 MG			37.60	
		TEGRITAL-CR - 200MG			23.47	

2	OXCARBAZEPINE	OXETOL-300mg TABLET(OXCARBAZEPINE)	Tablet	Adult: As monotherapy or adjunctive therapy:Initially,600mg daily in 2 divided doses,may increase if necessary in max increments of 600mg daily at wkly interval. Maintenance dose:600-1200mg daily or up to 2,400 mg daily. Child:≥6 yrs:8-10mg/kg daily in 2 divided doses,may increase if necessary in increments of 10mg/kg daily at wkly interval. Maintenance dose:30mg/kg daily	88.00	Dizziness, drowsiness, tiredness, nausea/vomiting, stomach/abdominal pain, headache, trouble sleeping, or constipation
3	DIVALPROEX	DICORATE ER - 250MG	Tablet	Adult: 10 mg/kg/day	56.00	Somnolence, heart block, fatigue, restlessness, excitation
4	GABAPENTIN	GABAGESIC - 300MG	Tablet, Capsule	Adult: 300 - 60mg OD- TDS	98.00	Mild sedation, dizziness and unsteadiness
		GABANTIN - 100MG			50.00	
		GABANTIN - 300MG			135.00	
		GABANEURON			112.85	
		NEUCOBAL G			141.00	
		RENERVE G			173.50	
		GABAPIN-300mg CAPSULE			240.00	
5	GABAPENTIN(300 MG), METHYLCOBALAMIN	GABAGESIC-M-TABLET	Tablet	Dose: 300mg to 600mg about three times a day.	134.00	Dizziness, Sleepiness, Tiredness, Uncoordinated body movements.
6	LEVETIRACETAM	LEVIPIL - 500MG	Tablet, Injection	Adult: 500 mg BD - 3g OD	118.00	Somnolence, asthenia, dizziness, vertigo, depression, emotional instability, hostility,
		LEVIPIL - 5ML			147.00	
		EPITRA-250mg TABLET			59.47	
		EPITRA-500mg TABLET			124.32	
		EPITRA-5ml INJECTION			110.88	
		LEVIPIL-100ml SYRUP			382.92	diplopia.
		LEVIPIL-250mg TABLET			59.45	
7	PHENYTOIN SODIUM	DILANTIN - 2 ML	Tablet, Injection	Adult: 100 mg OD-TID 15 - 20 mg/kg IV Paed: 5-10 mg/kg/day	11.50	Gum hypertrophy, hirsutism, megaloblastic anemia, HS reaction
		EPTOIN - 100 MG			164.58	
		EPTOIN - 50 MG			11.50	
8	PREGABALIN	GABAWIN-25MG	Tablet, Capsule	Adult: PO Adjunct in partial seizures Initial: 150 mg/day, may increase to 300 mg/day after a wk. Anxiety Initial:150	45.00	Somnolence, dizziness, headache,
		GABAWIN-50MG			72.00	
		LYRICA - 75 MG			768.56	
		PREGABA - 75MG			110.00	

		PREGABA - M		mg/day, may increase in	104.05	
		PRELOGIC		increments of 150 mg wkly.	125.00	
		PREGATOR - 75 MG		Max:600mg/day.Neuropathic pain Initial: 150 mg/day, may increase to 300 mg/day after 3-7 days. Max: 600 mg/day after a 7-day interval.	159.81	
9	SODIUM VALPROATE	ENCORATE 100MG -5ML	Tablet, Syrup	Adult: 200-800 mg TDS	30.08	GI upset, drowsiness, ataxia & tremor
		VALPARIN - 100 ML	Injection	Paed: 15 - 30 mg/kg/day	67.09	
		VALPARIN -200 MG			35.53	
		ENCORATE CHRONO - 200MG (C.R			46.00	
10	SODIUM VALPROATE + VALPROIC ACID	VALEPTOL-CR-200mg TABLET(SODIUM VALPROATE)	Tablet	10-15mg/kg/day ,increased by 5-10mg/kg/day at weekly interval.Max dose:60mg/kg/day	45.00	Nausea, Vomiting, Weight gain, Loss of appetite, Swelling of gums, Headache, Sleepiness, Tremor, Hair loss, Liver injury.
		VALEPTOL-CR-300mg TABLET(SODIUM VALPROATE)	Tablet		67.00	
11	SODIUM VALPROATE (100MG)	VALPROL-100mg/5ml-INJECTIN	Injection	Initial dose: 10 to 15 mg/kg orally or IV daily; increase by 5 to 10 mg/kg per week if necessary according to clinical response Maintenance dose: 10 to 60 mg/kg daily Maximum dose: 60 mg/kg daily	29.29	Injection site reactions (pain, swelling, redness), Loss of appetite, Nausea, Vomiting, Sleepiness, Hair loss, Weight gain, Liver function abnormal, Tremor.

ANTI-FUNGALS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	FLUCONAZOLE	AF 400 MG	Tablet	Adult: 100 - 400 mg OD Paed: 6 - 12 mg/kg/day OD	23.00	GI upset, Headache
		AF- 150 MG			9.90	Taste perversion, HS reaction
		FORCAN I.V- 100ML			118.50	Headache, Cutaneous rash, Anaphylaxis, Dry mouth, Alopecia, Seizure, Dizziness, Tremors.
		NUFORCE 3 KIT			99.00	
2	KETOCONAZOLE	HYPHORAL-200MG	Tablet, Lotion, Soap	Adult: 200 -400 mg OD Child> 2yrs: 3.3 - 6.6 mg/kg OD	199.00	Urticaria, Anaphylatic
		KZ - 75GM SOAP			155.00	Reactions, GI upset, Constipation,
		NOVALE - 75 GM BAR			109.00	
		AFK - LOTION			108.00	
		SCALPE - 75 ML			250.00	

		ARCOLANE - 60ML			238.29	
		KZ-CREAM			185.00	
		KETO-LOTION			84.00	
3	TERBINAFINE	SEBIFEN-250MG	Tablet, Cream	Adult: 250 mg OD	288.00	Headache, GI upset, Taste disturbance, insomnia, Abdominal pain, Diarrhoea, Cutaneous rash, URTI, Visual disturbances, cutaneous angioedema.
		TERBEST			84.00	
		TERBEST CREAM 15GM			53.00	
		TYZA-15GM CREAM			69.00	
		TYZA-250MG			108.50	
4	CLOTRIMAZOLE	CANDID - V - 30 GM GEL	Tablet, Cream, Lotion, Capsule, Powder, Ear drops, Mouth paint, Vaginal suppositories.	Vaginal: Adult:As pessary:100mg daily for 6 days,200mg daily for 3 days or 500mg as a single dose,alternatively apply 1,2 or 10% cream Ear: Adult: Instill 10 drops into the canal of the affected ear(s) once daily Child: 6 month -13 yrs instill 5 drops ,duration depending upon the disease Topical(creams,gel,suspension,s olution,powder): Apply onto affected area once daily(nails),bd(skin)	69.25	Erythema, Stinging, Irritation, redness, blistering, peeling, swelling, itching, burning sensation, stomach pain, nausea, vomiting and fever.
		CANDID - V3 TABLET			55.00	
		CANDID - B CREAM -- 20GM			89.50	
		CANSOFT-CL - CAPSULE (3'S)			73.50	
		CANDID TV - LOTION			179.50	
		CANDID - 20 GM CREAM			83.50	
		CANDID -100GM POWDER			104.00	
		CANDID - 10 ML EAR DROPS			53.00	
		CANDID - 20 ML MOUTH PAINT			89.00	
		CANDID V6 TABLET			58.15	
		CANDID - V3 TABLET			44.50	
		CANDID - CL - CAPSULES(3 NOS)			93.50	
5	SERTACONAZOLE NITRATE	ONABET - 15 GM CREAM	Cream	Adult: Apply to affected area bid for 4 wk	130.00	Swelling, itching, burning sensation, stomach pain,redness, oozing, dry skin
		ONABET - 30 GM CREAM			230.00	
		ONABET-B - 10 GM CREAM			118.00	
6	ITRACONAZOLE	I-TYZA-100MG	Capsule	100-200 mg OD/BID	87.50	Urticaria, anaphylactic
		ITAZOL - 100MG			357.50	Reactions, GI upset,cutaneous angioedema, hepatotoxicity,thrombocytopenia, hearing loss, tinnitus
7	KIT OF TWO ONE ORNIDAZOLE, FLUCONAZOLE AND ONE AZITHROMYCIN	AF-KIT TABLET	Tablet	Ornidazole: Adult:0.5-1.5gm daily for 1-7 days Azithromycin:1 gm as a single dose 1 hr before or 2 hrs after food with plenty of water .	21.00	GI Irritation, Nausea, Vomiting
8	MICONAZOLE	DK - GEL	Gel	Adult & children: Apply thinly over affected are bd for 2-6 week. Continue treatment for atleast 1 wk after symptoms have disappeared.	116.00	Redness, severe blistering, irritation of skin, itching, peeling, dry skin, dry mouth, sore tongue, tooth pain, headache.

9	NATAMYCIN (5%)	NATAFORCE EYE DROPS	Drops	Adult: As 5% ophth susp: Instill 1 drop in the conjunctival sac every 1-2 hr,reduce to 1 drop 6-8 times daily after 3-4 days.	111.09	mild eye irritation or discomfort (redness stinging or burning)
10	KETOCONAZOLE	TARRY SHINE-100ml SHAMPOO	Lotion	Adult: As 2% shampoo: Apply on scalp once daily for up to 5 days. For prophylaxis: use once daily for a max of 3 days before exposure to sunlight	230.00	Application site reactions (burning, irritation, itching and redness)
11	TERBINAFINE (1% W/W)	TYZA-30gm CREAM	Cream	Adult: Apply once or bd to affected area(s) for 1-2 weeks	135.00	Skin peeling, Itching.

ANTI-HISTAMINE

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	CETRIZINE HCL	CETRIZET - 10MG	Tablets	Adult: 10 mg OD Paed: children >6 yrs 5-10mg	20.00	dizziness, dry mouth, GI upset, disturbed coordination, confusion.
		AVIL NU - 10 MG			20.00	
		CETZINE-60ml SYRUP	Syrup		36.14	
		ALERID-10mg TABLET	Tablet		17.33	
2	LEVOCETIRIZINE	ALLERONE-M-TABLET	Tablet	Adult: 2.5-5mg once daily in the evening Child(2-5 yr):1.25mg once daily;6-11yr :2.5mg once daily;≥12yr:2.5-5mg once daily.All doses to be taken in the evening.	108.00	Drowsiness, tiredness, and dry mouth
		LECETIRICAS-5mg TABLET	Tablet			
3	CETRIZINE HCL + AMBROXYL	RELENT		OD, 2-5 yrs: 2.5-5 mg/day	44.80	Stomach upset, Dryness in mouth, Fatigue, Dizziness, Sedation, Allergic reaction.
4	CINNARIZINE	STUGERON	Tablets	Adult: 25-50 mg TID	130.50	dizziness, dry mouth, GI upset, disturbed coordination, confusion.
5	FEXOFENADINE	ALLEGRA - 180MG	Tablets	Adult: 120-180 mg OD-BD Paed:6-11 yrs: 30 mg BD	129.40	dizziness, dry mouth, GI upset, disturbed coordination, confusion.
		ALLEGRA 120MG			141.50	

6	LEVOCETIRIZINE DIHYDROCHLORIDE	LECOPE-AD	Tablets, Syrup	Adult: 5 mg OD	37.50	dizziness, dry mouth, GI upset, disturbed coordination, confusion.
		LEVORID			45.50	
		LE ZYNCET -10MG			90.35	
		LE ZYNCET			46.25	
		HH LEVO-5MG			49.00	
		TELEKAST-L KID - 60ML			75.00	
		XYZAL-M - 5MG			95.90	
		LEZYNCET - D			46.70	
7	PHENIRAMINE MALEATE	PHENIRAMINE MALEAE 2ML	Tablets, Injections	Adult: 25-50 mg BID Paed: 6-14 yrs: 15-22.5 mg; 1-5 yrs: 15 mg; 4 months to 1 year:7.5mg	3.04	dizziness, dry mouth, GI upset, disturbed coordination, confusion.
		AVIL - 25 MG			5.70	
8	CHLORPHENIRAMINE, PARACETAMOL, AND PHENYLEPHRINE	CORIMINIC-15ml DROPS	Drops, Syrup	Drops(1mg/ml):6 months to 11 months: 0.75 mL orally 4 times a day. 1 year to 2 years: 1 mL orally 4 times a day. Oral syrup(4mg/5ml):2 to 5 years: 1.25 ml orally every 4 hours to 6 hours, not to exceed 7.5 mL in 24 hours. 6 to 11 years: 2.5 ml orally every 4 hours to 6 hours, not to exceed 15 mL in 24 hours. 12 years or older: 5 ml orally every 4 hours to 6 hours, not to exceed 30 mL in 24 hours.	54.90	Drowsiness, dizziness, dry mouth/nose/throat, headache, upset stomach, constipation, or trouble sleeping
		CORIMINIC-60ml SYRUP			66.90	
9	PROMETHAZINE HYDROCHLORIDE	PHENERGAN - 100 ML	Tablets, Syrup, Injection	Adult: 25mg every 4-6 hours Paed: 0.25-0.5 mg/kg every 4-6 hours	23.98	dizziness, dry mouth, GI upset, disturbed coordination, confusion.
		PHENERGAN - 2 ML			6.92	
		TIXYLIX - 60ML			64.90	
		AVOMINE - 25 MG			28.90	
10	PROMETHAZINE TEOCLATE	AVOMINE-25mg TABLET	Tablet	Adult: 25mg at night,increase to 25mg bd if necessary. Alternatively,10-25mg 2-3 times daily. Child(2- 5yr):5-15mg daily in 1 or 2 divided doses; (>5-10 yr):10- 25mg daily in 1 or 2 divided doses.	3.48	Tiredness,Sleepiness
11	CYPROHEPTADINE TABS 5MG	CIPLACTIN	Tablets	4 mg tid. Usual: 12-16 mg/ day in 3-4 divided doses. Max: 32 mg/day.	23.50	Confusion, disturbed coordination,dizziness, excitation, euphoria, hallucinations, headache, hysteria, insomnia, irritability,

12	DESLORATADINE	DAZIT-5MG	Tablets	5 mg once daily.	64.00	Pharyngitis, dry mouth, myalgia, fatigue, somnolence, headache, dizziness, insomnia, dysmenorrhoea,
	DESLORATADINE + MONTELUKAST	MONDESLOR			139.00	
13	HYDROXYZINE HCL	HYRAX-10	Tablets, Syrup	25 mg at night, up to 25 mg 3-4 times/day.	17.00	Dizziness, ataxia, fatigue, drowsiness, hallucination, seizure, dry mouth, nervousness, xerostomia, involuntary movements, slurred speech,
		HYRAX - 100 ML			44.00	
		ATARAX -10 MG			25.17	
		ATARAX - 25MG			45.00	
14	LORATADINE	ALASPAN - 10 MG	Tablets	10 mg once daily or 5 mg 12 hrly.	83.00	Dizziness, ataxia, fatigue, drowsiness, hallucination, seizure, dry mouth, nervousness, xerostomia, involuntary movements, slurred speech,
15	MIZOLASTINE	ELINA	Tablets	10 mg/day.	97.00	dry mouth, diarrhoea, abdominal pain, nausea; transient drowsiness, headache, dizziness; anxiety and depression; raised liver enzymes
16	OLAPATADINE HCL	OLOPAT OD DROPS-3ML	Eye drops	Adult & Child: ≥3yrs: As 0.2% solution : 1 drop in the affected eye(s) once daily.	119.00	Eye irritation, burning sensation
17	CINNARIZINE-20MG+DIMENHYDRINATE-40MG	DIZITAC-TABLET	Tablets	Tablet taken for 3 times daily for balance problems and 8th hrly for motion sickness(before food)	80.00	Nausea, Dyspepsia, Weight gain, Drowsiness, Sleepiness, Dizziness, Weakness, Decreased blood pressure, Coordination impaired, Impaired concentration.
18	HYDROXYZINE HYDROCHLORIDE	HYRAX-100ml SYRUP	Syrup	Adult: Initially, 25mg at night increased if required up to 25 mg 3-4 times daily. Child: 6months -6yrs initially, 5-15mg/day, increased to 50mg/day in divided doses; >6 yrs initially, 15-25mg/day, increased up to 50-100mg/day in divided doses.	53.00	Dizziness, Headache, Drowsiness, Fatigue, Nervousness, Blurred Vision, Tremors, Seizures And Convulsions.
19	HYDROXYZINE HYDROCHLORIDE	HYRAX-10mg TABLET	Tablets	Adult: Initially, 25mg at night increased if required up to 25 mg 3-4 times daily. Child: 6months -6yrs initially, 5-15mg/day, increased to 50mg/day in divided doses; >6 yrs initially, 15-25mg/day, increased up to 50-100mg/day in divided doses.	2.04	Sedation, Nausea, Vomiting, Stomach upset, Constipation, Diarrhoea, Loss of appetite.

20	OLAPATADINE (TOPICAL) (0.1% W/V)	OLOPAT-5ml DROPS	Drops	Adult & Child: ≥3yrs: As 0.2% solution : 1 drop in the affected eye(s) once daily.	126.95	Headache, blurred vision, burning/stinging/redness/dryness of the eye, eyelid swelling, or a feeling as if something is in your eye
		OLOPAT-OD-3ml DROPS	Drops		142.00	
21	NAPHAZOLINE HCL CHLORPHENIRAMINE MALEATE	REDIF-10ml EYE DROPS	Drops	Instill one or two drops in the affected eye(s) 3 times a day	51.30	Eye irritation, Burning eyes, Stinging in the eyes.
22	CHLORPHENIRAMINE (2MG/5ML) + PHENYLEPHRINE (5MG/5ML)	T-MINIC-15ml DROPS	Drops	Drops (1mg/ml): 6 months to 11 months: 0.75 mL orally 4 times a day. 1 year to 2 years: 1 mL orally 4 times a day. Oral syrup (4mg/5ml): 2 to 5 years: 1.25 ml orally every 4 hours to 6 hours, not to exceed 7.5 mL in 24 hours. 6 to 11 years: 2.5 ml orally every 4 hours to 6 hours, not to exceed 15 mL in 24 hours. 12 years or older: 5 ml orally every 4 hours to 6 hours, not to exceed 30 mL in 24 hours.	63.80	Drowsiness, dizziness, headache, blurred vision, upset stomach, nausea, constipation, or dry mouth/nose/throat
		T-MINIC-60ml SYRUP	Syrup		68.20	

ANTI-HYPERTENSIVE

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMLODIPINE + HYDROCHLOROTHIOZIDE	AMLONG-H	Tablet	Adult : 5-20 mg OD	54.50	Headache, pedal edema, palpitation, abdominal discomfort
		AMLODIPINE			AMLONG - 10 MG	
		AMLONG - 2.5 MG			29.85	
		AMLONG - 5 MG			49.35	
		STAMLO -2.5 MG			31.00	
		STAMLO -5 MG			49.05	
	AMLODIPINE + ATENOLOL	STAMLO BETA			104.00	
		AMLONG - A			88.00	
	AMLODIPINE + METOPROLOL	AMLONG MT - 50 MG	65.00			
2	ATENOLOL	ATPARK - 50 MG	Tablet	Adult: 25 - 100 mg OD Paed: 1-2 mg/kg/day in single or 2 divided doses	33.52	Fatigue, bradycardia, cold extremities, epigastric discomfort

		ATPARK - 25 MG			30.83	
3	CARVEDILOL 3.175 MG	CARDIVAS 3.125 MG	Tablet	Adult: 12.5 - 25 mg OD	39.00	Fatigue, bradycardia, cold extremities, epigastric discomfort
		CARDIVAS-12.5mg TABLET			82.00	
		CARDIVAS-6.25mg TABLET			52.00	
4	CILNIDIPINE	NULONG - 10MG	Tablet	Adult: 5 -20 mg OD	68.50	Headache, pedal edema, palpitation, abdominal discomfort
		NULONG - 5MG			36.00	
5	CLONIDINE	ARKAMIN	Tablet, Injection	Adult: start with 0.1 mg BID max upto2.4mg/day Paed: 5-7 mcg/kg/day	1.38	Constriction of pupil, drowsiness, bradycardia
		CLONEON			47.00	
6	DILTIAZEM HYDROCHLORIDE	DILZEM - 30 MG	Tablet	Adult:30-120 mg TDS Paed: 1.5 - 2 mg/kg/day	26.15	GI upset, AB block, bradycardia, CHF, edema
		DILZEM-CD - 90MG			105.00	
		DILZEM SR - 90 MG			94.00	
7	ENALPRIL MALEATE	ENVAS - 5 MG	Tablet	Adult: 5 -20 mg	51.44	Dry cough, orthostatic hypotension, GI upset
		ENVAS - 2.5 MG			30.83	
8	LABETALOL	LABLOL - 4ML	Injection	Adult: 20-80 mg i.v upto max 300 mg	326.50	Fatigue, bradycardia, cold extremities, epigastric discomfort
9	LOSARTAN	LOSAR - 25 MG	Tablet	Adult: 25 - 100 mg/day in 1-2 divided doses Children > 6yrs: 0.7 -1.4 mg/kg OD	43.37	Hypotension, tachycardia, bradycardia, muscle cramps
		LOSAR - 50 MG			74.65	
		REPACE - 50 MG			47.99	
		COVANCE - 50 MG			47.97	
	LOSARTAN HYDROCHLOROTHIOZIDE	LOSAR-H		120.40		
10	METOPROLOL SUCCINATE	BETAONE XL - 25 MG	Tablet, Injection	Adult: TAB 100 mg OD - 200 mg BD Inj 5-15 mg	37.74	Fatigue, bradycardia, cold extremities, epigastric discomfort
		BETAONE XL - 50 MG			55.92	
		MET XL - 50 MG			87.16	
		PROLOMET - XL - 12.5 MG			29.00	
		PROLOMET - XL - 25 MG			39.12	
		METOLAR - XR - 50 MG			105.50	
		METOLAR 12.5 MG			27.50	
		METOLAR-5ML			12.31	
		BETALOC-25mg TABLET	Tablet		91.50	
BETALOC-INJECTION	Injection		15.10			
11	NIFEDEPINE	DEPIN - 5 MG	Tablet, Capsule	Adult: 30 - 120mg OD	48.00	Headache, pedal edema, palpitation, abdominal discomfort
		NICARDIA RETARD - 10 MG			24.70	
		NIFEDINE -10MG			13.08	
12	OLMESARTAN MEDOXOMIL AND COMBINATION	OLMETRACK - 20 MG	Tablet	Adult: 20-40 mg OD	76.69	Hypotension, tachycardia, bradycardia, muscle cramps
		OLMEZEST-40 MG			150.00	

		OLMEZEST-AM			100.00	
		OLMEZEST H - 40 MG			155.00	
13	RAMIPRIL	HOPACE - 5MG	Tablet	Adult: 2.5 - 20mg OD	180.00	Hyperkalemia, angioedema, cough, renal dysfunction
14	RAMIPRIL+HYDROCHLORO THIAZIDE	CARDACE H-2.5mg/12.5mg TABLET	Tablet	RAMIPRIL(2.5/5mg)+HYDRO CHLOROTHIAZIDE(12.5mg): Tablet taken once daily	78.46	Decreased blood pressure, Cough, Increased potassium level in blood, Fatigue, Weakness, Dizziness, Renal impairment.
		CARDACE H-5mg/12.5mg TABLET	Tablet		123.65	
15	SODIUM NITROPRUSSIDE	NITOSIDE-50MG	Injection	Adult: 0.5 - 10 mcg	144.00	Palpitation, nervousness, lactic acidosis, GI discomfort
				/kg/min		
16	TELMISARTAN	CRESAR - 20MG	Tablet	Adult: 20-80 mg/day	36.00	Hypotension, tachycardia, bradycardia, muscle cramps
		CRESAR - 40MG			75.50	
		TAZLOC-40MG			78.71	
		TELMA - 20 MG			65.60	
		TELSARTAN - 80 MG			142.00	
	TELMISARTAN AND COMBINATION	TELMA - 40 MG		8.19		
		ARBITEL - 20MG		45.00		
		ARBITEL - 40 MG		81.50		
		TELMA AM		143.40		
		TELMA-CT -40MG		144.75		
		TELMA - H -80MG		269.18		
		TELMA - H		161.81		
		CRESAR-H		76.00		
17	VERAPAMIL HCL 40MG	CALAPTIN - 40 MG	Tablet	Adult: 80 - 160 mg bd	8.03	Headache, pedal edema, palpitation, abdominal discomfort
18	ALPHAMETHYLDOPA	ALPHADOPA	Tablet	Adult: 0.5 - 2g in divided doses Paed: 10mg/kg/day in 2-4 divided doses	160.00	Sedation, headache, orthostatic hypotension, edema, GI upset, bone marrow depression
19	BISOPROLOL & HYDROCHLOROTHIAZIDE	LODOZ - 2.5 MG	Tablet	5-10 mg once daily. Max: 20 mg/day.	58.00	Bradycardia, worsening of pre-existing heart failure, hypotension, dizziness, headache, GI disturbances
20	BISOPROLOL FUMARATE	CONCOR COR-2.5mg TABLET(BISOPROLOL FUMARATE)	Tablet	Adult:5-10mg once daily(max.20mg/day)	53.35	Nausea, Headache, Fatigue, Constipation, Diarrhoea, Dizziness, Cold extremities.
		CONCOR-5mg TABLET(BISOPROLOL FUMARATE)	Tablet		72.38	

21	BOSENTAN	LUPIBOSE - 62.5 MG	Tablet	62.5 mg bid upto 125 mg BID	762.50	Headache, nasopharyngitis, flushing, fluid retention (e.g. peripheral oedema), hypotension, palpitations, dyspepsia
22	LISINOPRIL	LISTRIL 2.5 MG	Tablet	Initial: 10 mg/day given at bedtime. Maintenance: 20 mg once daily, up to 80 mg/day if needed.	52.05	Headache, fatigue, persistent and non-productive cough, chest and abdominal pain, dizziness, upper resp tract infection, asthenia, rash, orthostatic effects, hypotension,
23	PRAZOSIN	PRAZOPRESS - 1 MG	Tablet	0.5 mg bid or tid	50.00	During cataract surgery, Intraoperative Floppy Iris Syndrome(IFIS)may occur. Postural hypotension, syncope, tachycardia, palpitations, lackofenergyDzziness,
		MINIPRESS XL - 2.5 MG			300.30	
		MINIPRESS XL - 5MG			353.76	
24	PROPRANOLOL	INDERAL- 10 MG	Tablet	40 - 80 mg BID	12.60	Bradycardia, hypotension, syncope, shock, angina pectoris.Lightheadedness, giddiness, ataxia, dizziness, irritability, sleepiness, hearing loss, and visual disturbances to vivid dreams, hallucinations, and confusion.
		INDERAL-40MG			29.61	
		CIPLAR - LA - 40MG				
25	NIMODIPINE (30MG)	NIMODIP-30mg TABLET	Tablet	Adult: 60mg 4 hrly beginning within 4 days of onset of haemorrhage & continued for 21 consecutive days	160.00	Dizziness, lightheadedness, flushing, or swelling ankles/feet.
26	PAPAIN+UREA	AUTOLYSE-15gm OINTMENT	Ointment	Clean the affected area with saline or with mild wound cleansing solution.Do not use hydrogen peroxide.Apply ointment directly to the lesion cover with appropriate dressing and secure into place.Wash the lesion each time when dressing is changed	159.00	Mild stinging or burning of the skin where the medicine is applied.

ANTI-MALARIAL

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ARTEMETHER & LUMEFANTRINE	LUMERAX - 80 MG	Tablet	80 mg/day.	175.00	Mild GI disturbance, dizziness, tinnitus, reduction in reticulocyte and leucocyte counts, nausea, vomiting, abdominal pain, bradycardia, 1st-degree heart block
2	ARTESUNATE	FALCIGO	Injection	Adult: IV/IM Initial: 2.4 mg/kg.	224.82	Transient and reversible , rash,

				Repeat 12 hr and 24 hr later, then once daily thereafter.		reticulocytopenia, drug fever bradycardia, transient 1st-degree heart block and reversible elevation of serum transaminases.
3	CHLOROQUINE PHOSPHATE	LARIAGO - 250 MG	Tablet, Suspension	Initial: 600 mg, then 300 mg 6-8 hr later on day 1. Continue w/ 300 mg/day on days 2 and 3	8.21	Retinopathy, hair loss, photosensitivity, tinnitus, myopathy (long-term therapy). Psychosis, seizures, leucopenia and rarely aplastic anaemia, hepatitis, GI upset, dizziness, hypokalaemia
		LARIAGO - 60 ML			18.62	
		LARIAGO - DS			8.04	
4	HYDROXY CHLOROQUINE	HCQ	Tablet	200-400 mg/day. Max: 6.5 mg/kg/day	65.49	Retinopathy, hair loss, tinnitus, photosensitivity, myopathy (long-term therapy). Psychosis, seizures, leucopenia and rarely aplastic anaemia, hepatitis
5	PRIMAQUINE PHOSPHATE	MALIRID - DS	Tablet	15 - 30 mg/day	40.00	Nausea, vomiting, epigastric distress abdominal cramps, leukopenia, leucocytosis, agranulocytosis, methaemoglobinemia in NADH methaemoglobin reductase-deficient individuals.

ANTI - PARASITIC

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	METRONIDAZOLE	FLAGYL - 100 ML	Tablet, Suspension	Adult: TAB 250 mg - 750mg INJ: 15mg/kg IV	12.83	GI upset, metallic taste, seizures, Peripheral neuropathy
		FLAGYL - 400 MG			12.29	
		METRONIDAZOLE - 400			12.29	
		FLAGYL-60ml SUSPENSION			17.01	
		METROGYL GEL-30gm				
	METRONIDAZOLE + NORFLOXACIN	NOR- METROGYL (SUSP) -30 ML		Paed : 100mg +100 mg	12.54	Dizziness, headache, stomach upset, nausea, vomiting, loss of appetite, diarrhea, constipation, or metallic taste in your mouth
2	ORINDAZOLE	ORDENT (ORINDAZOLE+ OFLOXACIN)	Tablet	Adult: 500 mg - 1.5g Child 10-40mg/kg OD	97.00	GI upset, vertigo, headache
		DAZOLIC - 500 MG (ORINDAZOLE)			90.00	
3	SATRANIDAZOLE	SATROGYL - 300 MG	Tablet	Adult: 300 - 600 mg OD	42.25	Headache, dry mouth, weakness and dizziness.
	SATRANIDAZOLE + OFLOXACIN	SATROGYL - O			106.00	
4	SECNIDAZOLE	SECNIL FORTE - 1 GM	Tablet	Adult: 1.5-2g OD	100.00	Nausea, gastralgia, change of taste,

						metallic taste, stomatitis, urticaria, rashes, leucopenia. Rarely vertigo, moderate neurological, digestive disturbances.
5	TINIDAZOLE	FASIGYN - 500MG	Tablet	Adult: 2g OD	28.00	Transient leucopenia, ataxia, convulsions, peripheral neuropathy, sensory disturbances, vertigo, metallic taste, flushing. Potentially Fatal: Hypersensitivity.
6	DIETHYL CARBAMAZINE CITRATE	HETRAZAN - 100 MG	Tablet	Filariasis; Loiasis; Toxocariasis Inital: 1 mg/kg/day, may increase to 6mg/kg/day	52.50	Fever, headache, vomiting, dizziness drowsiness, nausea, chills.
7	IVERMECTIN	IVERA - 30ML	Tablet, Shampoo	150-200 mcg/kg OD	74.00	Diarrhoea, nausea, vomiting, dizziness, pruritus, urticaria, rash, arthralgia, fever, myalgia, asthenia, postural hypotension, tachycardia, oedema, lymphadenopathy, sore throat, cough, headache, somnolence, transient eosinophilia, raised liver enzyme values.
		IVERMECTOL - 12 MG			59.50	
		IVERMECTOL - 6 MG			35.50	
	IVERMECTIN + ALBENDAZOLE	BANDY PLUS		Adult : 6mg - 400mg	19.00	
8	ALBENDAZOLE	BANDY SUSPENSION	suspension	Child: 15mg/kg daily in 2 divided doses.Max.800mg/day.	16.20	skin rashes, swelling, nausea, vomiting headache, drowsiness, dizziness, and diarrhea.
		ZENTEL-SYRUP	syrup		16.21	vision changes, yellowing eyes or skin, severe stomach or abdominal pain,

ANTI-PARKINSONISM

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	BROMOCRIPTINE TABS 2.5MG	B-CRIP--2.5MG	Tablet	Adult: 1.25 -20 mg/day	142.80	GI upset, hallucination,hypotension
2	LEVODOPA&CARBIDOPA	SYNDOPA - 110 MG	Tablet	Adult: start with 1 tablet (carb: 10mg, leva: 100mg) TID increase gradually	17.51	GI upset, postural hypotension, cardiac arrythmia
		SYNDOPA PLUS			23.73	
3	SELEGILINE HCL	SELGIN - 5MG	Tablet	Adult: 10mg /day at breakfast & lunch	46.50	GI upset, postural hypotension, cardiac arrythmia
4	TRIHXYPHENIDYL HCL	PACITANE - 2 MG	Tablet	Adult: 2-10 mg/day	47.26	Dryness of mouth, impairment of memory, confusion
5	CITICHOLINE	CITINERVE - 500 MG	Tablet, injection	PO Parkinson's disease;	400.00	Head ache, Diarrhoea

		CITILIN-250MG - 2ML		Cerebrovascular disorders; Head injury 200-600 mg/day in divided doses. IV/IM Parkinson's disease;	106.75	
6	DONEPEZIL	DONECEPT - 5MG	Tablet	5 mg/day in the evening, up to 10 mg/day after 4-6 wk.	94.50	Nausea, vomiting, anorexia, wt loss diarrhoea, insomnia, fatigue, muscle cramps; headache and dizziness; syncope, bradycardia; convulsions;

ANTI-PSYCHOTICS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMISULPRIDE	AMISANT - 200MG	Tablet	Adult: PO 400-800 mg bid. Max: 1.2 g/day.	118.00	insomnia, wt gain, QT prolongation, hypotension, bradycardia, GI disorders hyperglycaemia; breast pain, gynaecomastia, galactorrhoea.
		SULPITAC - 100 MG			87.00	
2	CLOZAPINE	SIZOPIN -50MG	Tablet	Adult:50-300mg /Day	53.00	Agranulocytosis, blood dyscrasis, metabolic syndrome
		SIZOPIN - 25 MG			32.00	
3	HALOPERIDOL	SERENACE 0.25MG	Tablet, Injection	Adult:2-20 mg/day	11.95	Parkinsonism, Autonomic effects
		SERENACE - 1 ML			6.52	
		SENORM-LA			190.00	
4	OLANZAPINE	OLEANZ - 10MG	Tablet	Adult:2.5-10 mg/day	61.67	Agranulocytosis, blood dyscrasis, metabolic syndrome
		OLEANZ - 5MG			32.40	
5	CHLOPROMAZINE	CHLOPROMAZINE-50mg TABLET	Tablet	Adult:25mg tid;may be given as a single 75mg dose at night.Maintenance: 25-100mg tid increased to ≥1g daily as required in psychotic patients. Child:1-12 yr:500mcg/kg every 4-6hr.Max:>5yr:75mg daily;1-5yr:40mg daily. Elderly:Initially,1/3-1/2 the normal adult dose.	4.85	Drowsiness, dizziness, lightheadedness, dry mouth, blurred vision, tiredness, nausea, constipation, weight gain, and trouble sleeping
6	QUETIAPINE	QUTIPIN - 50MG	Tablet	Adult:50-400 mg/day		Sedation, postural hypotension, urinary incontinence
7	RISPERIDONE	RESPIDON-2MG	Tablet	Adult:2.-12 mg/day	30.00	Increase in prolactin levels , postural hypotension, parkinsonism
8	LITHIUM CARBONATE	LITHIC SR - 400MG	Tablet	Adult:600-1200mg/day	39.00	Thirst, polyurea, tremors, seizures CNS toxicity, GI upset, renal
		LITHIC SR - 450MG				

						diabetes insipidus, goitre
9	ZUCLUPENTHIXOL DECANOATE INJ	CLOPIXOL DEPOT-200 - 1ML	Injection	IM Acute psychosis As acetate ester: 150 mg, repeat if needed 2-3 days	370.40	Drowsiness, blurred vision, tachycardia, nausea, dizziness, headache, excitement, postural
				later. Max cumulative dose: 400 mg/course. Max number of inj: 4/ course Max duration:2wk		hypotension, hyperprolactinaemia, early phase of treatment; urinary frequency or incontinence; tardive dyskinesia.
10	ARIPIPRAZOLE	ARPIZOL - 15 MG	Tablet	Adult: 10 mg or 15 mg once daily	116.00	GI disorders, headache, anxiety, wt gain, insomnia, lightheadedness, drowsiness,
		ARPIZOL-10mg TABLET	Tablet		103.00	

ANTI-TUSSIVES & EXPECTORANTS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMBROXOL+	MUCOLITE	Tablet, Syrup, Drops	Ambroxol: 15 ,mg Guaiphenesin 50mg Terbutaline: 1.25 mg	25.00	Mild GI - effects, Allergic reaction
	TERBUTALINE + GUAIPHERNESIN	MUCOLITE - 100 ML			59.00	
		MUCOLITE SR - 75 MG			62.50	
		MUCOLITE - 15ML			19.60	
		AERODIL - 100ML			68.00	
		CHESTON PLUS - 100ML			48.50	
2	AMBROXOL+GUAIFENESIN+ LEVOSALBUTAMOL	CHERICOF-LS-100ml SYRUP	syrup	Adult: 30-120mg daily in 2 -3 divided doses	71.00	Nausea, Vomiting, Diarrhoea, Stomach upset, Stomach pain, Allergic reaction, Dizziness,
		ASCORIL-LS-SYRUP	syrup		90.00	
3	BROMHEXINE HYDROCHLORIDE	BROMHEXINE - 100 ML	Tablet, Syrup	Adult 4-8 mg tablet	74.00	Headache, Sweating, Skin rashes
		BROMHEXINE - 8MG			3.80	
4	BROMHEXINE 4 MG+TERBUTALINE	BRONCOREX-50ml SYRUP	syrup	Bromhexine-4 mg,two-four times a day Terbutaline Sulphate-1.25 mg,three times per day. Guaiphenesin-50 mg,200 to 400 mg 4 hourly	57.15	Nausea, Diarrhoea, Bloating, Indigestion, Vomiting, Stomach pain, Sweating, Dizziness, Headache, Skin rash, Urticaria, Tremor, Increased heart rate.
5	BROMHEXINE HYDROCHLORIDE +	ASCORIL EXPECTORANT - 100 ML	Expectorant	Adults and Children >12 years: 10 mL (2 tsp) 3 times a day. Children 6- 12 years: 5 mL (1 tsp) 3 times a day; 2-5 years: 2.5 mL (½ tsp) 3 times a day; <2 years: Not recommended.	76.00	drowsiness and dizziness

6	TERBUTALINE + GUAIPHENESIN	BRO - ZEDEX - 100 ML	syrup	Adults -10ml 2-3 times/day, 3 – 6 years 2.5 ml TID, 6-12 years 5 ml TID	75.00	Nausea, diarrhoea rash, angioedema, pruritus, urticaria, sweating, dizziness, vomiting, headache and pain in the upper part of abdomen.
		ASTHAKIND-100ml EXPECTORANT	Expectorant		58.56	Nausea or vomiting
7	BROMHEXINE 4 MG+GUAIFENESIN 50 MG+MENTHOL 2.5 MG+TERBUTALINE	BRO-ZEDEX-100ml SYRUP	syrup	Adults -10ml 2-3 times/day, 3 – 6 years 2.5 ml TID, 6-12 years 5 ml TID	98.00	Nausea, diarrhoea rash, angioedema, pruritus, urticaria, sweating, dizziness, vomiting, headache and pain in the upper part of abdomen.
8	CHLORPHENIRAMINE MALEATE	PIRITON EXPECTORANT - 100ML	Syrup	Chlorpheniramine: 4mg Dextromethorphan : 10mg/5 ml	58.60	Dizziness, GI disturbance
	CHLORPHENIRAMINE MALEATE + DEXTROMETHORPHAN	PIRITON-CS - 100ML			70.65	
		AERODIL-DX - 100ML			70.00	
		ASTHAKIND-DX - 60 ML			42.35	
		ASCORIL-D EXPECTORANT-100ML				
9	CHLORPHENIRAMINE MALEATE 2 MG,DEXTROME THORPHAN HYDROBROMIDE 10 MG, PH ENYLEPHRINE HYDROCHLORIDE	CHERICOF-SYRUP	syrup	Child:2-6yrs-1/2tsp;6-12- 1tsp;Adult-2tsp three to four times/day	76.50	Nausea, Vomiting, Loss of appetite, Headache.
10	CHLORPHENIRAMINE MALEATE 4 MG+CODEINE PHOSPHATE	CODISTAR-100ml SYRUP***	syrup	CHLORPHENIRAMINE MALEATE: 4 mg+CODEINE PHOSPHATE: 15-30 mg 4 times/day	98.49	Nausea, Vomiting, Constipation, Sleepiness, Lightheadedness, Dizziness, Shortness of breath, Sweating.
11	CHLORPHENIRAMINE MALEATE + LEVODROPROPIZINE	RESWAS	Syrup	CHLORPHENIRAMINE(2mg/5 ml): 2 times / day; LEVODROPROPIZINE(30mg/5 ml): Adult and children>12 yrs:60 mg 3 times/day;Child: >2 yr: 1 mg/kg tid for ≤7 days.	63.00	Dizziness, GI disturbance
12	PHENYLEPHRINE HCL	PHENPRESS 10MG - 1ML	Injection		227.00	Anxiety, reflex bradycardia, tachycardia, arrhythmias, headache, cold extremities/ gangrene, hypertension, nausea, vomiting, sweating, weakness, fear, restlessness

13	MENTHOL,CAMPHOR,TERPINEOL,EUCALYPTOL,CHLOROTHYMOLOL	KARVOL PLUS-CAPSULE	Capsule	Snip tip of cap and squeeze contents into half ltr hot or boiling water or on a tissue/cloth handkerchief and inhale vapours deeply and freely. At night, the inhalant can be applied on the pillow or night dress.	57.30	Nasal irritation, Dryness in the nasal cavity, Sore throat, Burning, Drowsiness, Vomiting, Abdominal pain, Coma.
14	LEVOCLOPERASTINE	LUPITUSS-100ml SYRUP	syrup	10-20 mg of levocloperastine is to be taken orally, thrice daily	95.50	Dizziness, drowsiness, nausea, and vomiting
15	DEXTROMETHORPHAN	ALEX COUGH-LOZENGES	Lozenges	Adult & child >12yrs: As loz: suck not more than 10 loz daily Child: As loz(6-12 yr) suck not more than 2 loz within any 4 hr, Max:7 loz daily.	78.00	Sleepiness, Dizziness, Excitation, Confusion, Convulsion, Respiratory depression, Nausea, Vomiting, Diarrhoea, Rash.
16	BENZONATATE	BENZ PEARLS-CAPSULE	Capsule	100mg capsule taken with or without food 3 times daily	7.33	Drowsiness, dizziness, headache, nausea, stomach upset,constipation, and stuffy nose
17	BROMHEXINE	BROMHEXINE-8mg TABLET	Tablet	Adult & child ≥12 yrs: 8-16mg tid. Child:(2-5yr) 8 mg daily in 2-3 divided doses;(6-11 yr): 4-8 mg tid	45.90	Feeling of fullness in the stomach (bloating), diarrhea, dizziness, headache, indigestion, nausea, sweating and skin rashes.
18	AMYL METACRESOL + DEXTROMETHORPHAN	TUSQ-D-LOZENGES	Lozenges	Adult: 1-2 lozenges every 4-6 hrs. Child: 6-12 yrs: 1 lozenge every 4-6 hrs.	39.00	Stomach upset, Sore tongue.
19	PHENYLEPHRINE + CHLORPHENIRAMINE + DEXTROMETHORPHAN	TUS-Q-DX-SYRUP	syrup	Oral: Adults: 1-2 teaspoonful 3-4 times daily, Children: 1/2-1 teaspoonful thrice daily	74.00	Dizziness, drowsiness, nausea, and vomiting.
20	GUAIFENESIN + TERBUTALINE + BROMHEXINE	TUSQ-X-100ml EXPECTORANT	Expectorant	Oral: Adults and children above 12 years : 10 ml 3 times daily, Children 2-6 years : 2.5 ml 3 times daily, Children 6-12 years : 5 ml 3 times daily	74.00	Nausea, Vomiting, Diarrhoea, Stomach pain, Stomach upset, Dizziness, Headache, Rash, Allergic reaction, Urticaria, Palpitations, Tremor, Increased heart rate.
21	CHLORPHENIRAMINE + DEXTROMETHORPHAN	ZEDEX-100ml SYRUP	syrup	Adult: 10 ml. Child 2-6 yrs: 2.5 ml; 6-12 yrs: 5 ml. All tid-qid	99.00	Dizziness, drowsiness, nausea, and vomiting.
22	HONEY AND CLOVE	KOFLET-H-LOZENGES	Lozenges	1 loz-three to four times a day	24.00	clinically safe.

ANTI-ULCER

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ANTACID GEL	DIGENE GEL - 200 ML	Tablets, Syrup	" Chew 1 -2 tablets four times a day Syrup:Shake bottle before use. Should be taken undiluted, but the suspension may be followed by a sip of water if	81.90	GI upset, headache, dry mouth confusion, drowsiness
		GELUSIL XTRA COOL - 200ML			85.00	
		DIGENE		desired. – 10 to 20ml four times daily – should be taken 15 minutes before meals and at bedtime – do not exceed recommended dosage."	12.00	
		MUCAINE - 200 ML GEL				
2	ESOMEPRAZOLE	RACIPER- 40 MG	Tablets, Capsules	Adult:20-40 mg OD	63.50	GI upset, headache, dry mouth confusion, drowsiness
		RACIPER-20 MG			37.50	
		ESOZ - 20MG			64.50	
		ESOZ - 40MG			102.50	
	ESOMEPRAZOLE + LEVOSULPRIDE	SOMPRAZ-L	165.00			
3	FAMOTIDINE	FAMOTIDINE - 20 MG	Tablet	Adult:20-40 mg BD	2.30	GI upset, hypotension, tachycardia, rash
		TOPCID - 40 MG			3.60	
4	OMEPRAZOLE	OMEZ - 20 MG	Capsules, Injection	Adult:20-40 mg OD Paed: 0.4-0.8 mg/kg/day	52.35	GI upset, headache, dry mouth confusion, drowsiness
		OMEZ			29.80	
5	PANTOPRAZOLE SODIUM + DOMPERIDONE	PANTOCID DSR	Tablet, Capsules, Injection	Adult:40 mg OD	119.00	GI upset, headache, dry mouth confusion, drowsiness
		PANTOCID - D			77.00	
	PANTOPRAZOLE SODIUM	PAN - 40MG	80.00			
		PAN 20MG	52.00			
		PANTOCID - 40 MG	79.00			
		PANTODAC I.V - 40 MG	54.50			
	PANTOPRAZOLE SODIUM LEVOSULPRIDE	PANTOCID-L	181.00			
6	RABEPRAZOLE	ROLES - 20MG	Tablet, Capsules, Injection	Adult:20 mg OD	59.00	GI upset, headache, dry mouth confusion, drowsiness
		RABLET - 20MG			66.00	
		RABIUM 20 MG			57.00	
		RAZO - 20MG			83.00	
	RABEPRAZOLE + DOMPERIDONE	RABIUM D SR	69.50			
		ROLES-D	77.50			
		RABIUM PLUS	163.00			
	RABEPRAZOLE +	ROLES-L	179.50			

	LEVOSULPRIDE					
7	RANITIDINE HCL	RANTAC - 150MG RANITIN - 2 ML RANITIN 150MG	Tablet, Injection	Adult: 150-300mg OD-BD; Injection:50-100mg every 6-8 hours Paed: 1-5 mg?kg?day in 2-4 divided dose njection:1-5mg every 6-8 hours	0.72 3.38 17.52	GI upset, hypotension, tachycardia, rash
8	SUCRALFATE	SUCRAFIL - 200 ML SUCRAFIL - 450 ML SUCRAFIL - O 200 ML	Syrup	Adult 1g TID	158.00 370.00 195.00	Flatulence, Abdominal fullness, GI upset
9	ENTRIC COATED ESOMEPRAZOLE MAGNESIUM AND DOMPERIDONE SR CAPSULES	SOMPRAZ-D - 40MG RACIPER - D	Capsule	20 - 40 mg OD	9.90 98.00	Increased risk of Clostridium difficile-associated diarrhoea (CDAD) and osteoporosis-related fracture. Headache, rash, pruritus, dizziness, fatigue, cough, back or abdominal pain, arthralgia
10	ITOPRIDE HCL	GANATON	Tablet	Adult: PO Antiemetic; As a prokinetic agent 50 mg 3 times/day.	86.50	Rash, diarrhoea, constipation, abdominal pain, headache, sleeping disorders, dizziness, galactorrhea, gynecomastia.
11	LANSOPRAZOLE	JUNIOR LANZOL - 15 MG	Tablet	1-11 yr ≤30 kg: 15 mg once daily in the morning for up 12 wk; >30 kg: 30 mg once daily in the morning for up t 12 wk. May increase doses up to 30 mg bid if patient is still symptomatic after 2 or more wk of treatment. 12-17 yr 15 mg once daily for up t 8 wk.	56.00	Diarrhoea, abdominal pain, nausea, vomiting, flatulence, constipation, headache, drymouth, peripheral oedema, dizziness, sleep disturbances, fatigue, paraesthesia, arthralgia, myalgia, rash, pruritus
12	MESALAMINE	MESACOL - 400 MG	Tablet	1-4 g/day BD/TDS	73.45	Abdominal pain (if new abdominal pain - consider pancreatitis); headache, nausea; flu; fatigue; fever, rash; sore throat; diarrhoea, joint pain
13	MOSAPRIDE CITRATE	MOZA - 5MG	Tablet	5 mg 3 times/day.	45.25	Diarrhoea, abdominal pain;dizziness constipation; headache; insomnia
14	OMEPRAZOLE, TINIDAZOLE AND AMOXICILLIN	HP-KIT CAPSULE	Capsule	omeprazole 20 mg, amoxicillin 750 mg, tinidazole 500 mg: 1 kit daily for 7 days		Nausea, Vomiting, Stomach pain, Flatulence, Indigestion, Dizziness, Tiredness, Headache, Skin rash, Dryness in mouth, Metallic taste.

15	LANSOPRAZOLE	JUNIOR LANZOL-15mg TABLET	Tablet	Adult:30mg once daily in the morning given for 4 - 8 week Maintenance:15mg/day.	101.00	Nausea, Headache, Abdominal pain, Constipation, Flatulence, Diarrhoea.
		JUNIOR LANZOL-30mg TABLET	Tablet		119.00	
16	SUCRALFATE (2GM)	SRFIL ENEMA-60ml	Enema	1-2 gm tablet mixed with 4.5-10ml of water in an enema applicator producing a low volume paste.Can be self-administered twice daily for 6 months.	95.00	Rectal bleeding, gas, bloating, diarrhea, stomach cramps, and rare allergic reactions, such as hives and difficulty breathing.
17	SUCRALFATE (1000MG)	SUCRAFIL-200ml SYRUP	Syrup	Adult: 1g 4 times daily or 2g bid for 4-8wk,may extend up to 12 wk if necessary. Max: 8 g daily	143.00	Dry mouth, upset stomach, gas, and nausea
18	SUCRALFATE + OXETACAINE / OXETHAZAINE	SUCRAFIL-O-200ml GEL	Syrup	Adult: 1 g four times per day on an empty stomach.	213.00	Constipation, Allergic reaction.

ANTI-VIRAL

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ACICLOVIR	ACIVIR DT - 200 MG	Tablets,Injection	Adult: 200-800 mg five times daily Paed: 5-10 mg/kg 5 times daily	81.50	GI upset, increase in liver enzymes, burning, itching, haematological changes, rashes.
		ACIVIR DT - 400 MG			67.35	
		ACIVIR DT - 800 MG			160.00	
		NEOCLOVIR - 500MG			450.00	
		ACIVIR EYE OINTMENT			52.90	
2	TENOFOVIR DISOPROXIL FEMARATE	TENVIR - 300 MG	Tablets	Adult>18 years: 300mg OD	1500.00	Flatulence, peripheral neuropathy, insomnia, body fat redistribution, raised liver enzymes
3	VALCYCLOVIR HCL	VALTOVAL-1GM	Tablets	500mg-2G tid	157.00	Acute renal failure, CNS effects Potentially Fatal: Thrombotic thrombocytopenic purpura/haemolytic uraemic syndrome in patients w/ advanced HIV disease. Acute renal failure, CNS effects
4	TENOFOVIR (300MG)	TENOF-300mg TABLET	Tablet	300mg-once daily paed: 8mg/kg daily	1325.60	Nausea, stomach pain, diarrhea, depression, headache, dizziness, weakness, trouble sleeping, itching or rash.

CNS - OTHERS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ERGOTAMINE, CAFFEINE, PARACETAMOL, PROCHLOPRAZINE	VASOGRain	Tablet	1-2 tabs as per the clinical condition needed.	66.50	Muscle cramps, stiffness, tiredness. Numbness and tingling of extremities Nausea, vomiting, diarrhoea. Ergotism consisting of hypotension gangrene of extremities, anginal pain
2	L-CARNITINE L-TARTRATE, MECOBALAMIN & FOLIC ACID	NUROKIND LC	Tablet	1,500 mcg/day in 3 divided doses.	89.89	Anorexia, nausea, vomiting and diarrhoea
3	PIRACETAM	NOOTROPHIL	Tablet, Injection	PO 2.4 to 7.2 g/day in 2-3 divided doses IV/IM 1-2 g 3 times/day.	449.47	Hyperkinesia, nervousness, depression, diarrhoea, rashes. CNS stimulation, sleep disturbances, dizziness, excitement, insomnia, somnolence, wt gain.
		NOOTROPIL - 100 MG			231.73	
		NOOTROPIL 15 ML - 200 MG			118.75	
	PIRACETAM + GINKGO BILOBA VINPOCETIN	COGNIVEL			174.40	
4	METHYLCOBALAMIN	NUROKIND FAST STRIPS	Strips	1tab/once or twice daily	14.00	headache, nausea, vomiting, dizziness, diarrhoea, back pain, arthritis, back pain, anxiety, nervousness, lack of coordination in body movement
5	FOLIC ACID · METHYLCOBALAMIN · VITAMIN B1	NUROKIND GOLD-TABLET	Tablet	1tab once or twice /daily	89.00	Drowsiness, dizziness, hypotension or a headache
6	MECOBALAMIN	NUROKIND-500mg TABLET	Tablet	1tab once or twice/daily	39.20	Nausea, Vomiting, Headache.
7	METHYLCOBALAMIN	NUROKIND-INJECTION	Injection	500 mcg/day 3 times/wk	15.00	Gastrointestinal discomfort.
8	METHYLCOBALAMIN & ALPHA LIPOIC ACID, FOLIC ACID & PYRIDOXINE HCL	NUROKIND PLUS-CAPSULES	Capsule	1-2 capsule /day	84.00	Nausea, Vomiting, Stomach ache, Loss of appetite, Flatulence, Skin rashes, Bloating.
9	METHYLCOBALAMIN+PYRIDOXINE + NICOTINAMIDE	NUROKIND PLUS-INJECTIONS	Injection	500 mcg/day 3 times/wk	22.00	Injection site related side effects, Flushing, Increased blood sugar level, Blurred vision, Rash, Elevations in liver function tests, Dizziness.
		NEURO-D3-CAPSULE	Capsule	1-2 CAPSULE per day		Allergic reactions like rash or itching, Swelling of the face, throat, and tongue, Severe dizziness, Trouble breathing.
10	MECOBALAMIN, GABPENTIN	NUROKIND-G-TABLET	Tablet	150mg per day may increased upto 300mg after a week	111.50	Dizziness, Sleepiness, Tiredness, Uncoordinated body movements.
11	METHYLCOBALAMIN + NIACINAMIDE+ VITAMIN B6	RE-XITE PLUS-2ml INJECTION	Injection	500 mcg daily IM/IV 3 times/wk.	22.00	Unusual or unpleasant taste in mouth
12	METHYLCOBALAMIN+ NIACINAMIDE + VITAMIN B6 (PYRIDOXINE)	RE-XITE PLUS-CAPSULE	Capsule	1500 mcg/day in 3 divided doses.	16.80	Constipation, diarrhea, or upset stomach

13	METHYLCOBALAMIN THIAMINE MONONITRATE PYRIDOXINE HCL ,FOLIC ACID	RE-XITE-BT-TABLET	Tablet	1 tab once or twice daily	65.00	Hypersensitivity, Abdominal and stomach pain, Blurred vision, Bloating, Headache, Irregular heartbeat, Hair loss, Constipation, Dry mouth.
14	MECOBALAMIN, FOLIC ACID, PYRIDOXINE HYDROCHLORIDE, VITAMIN D3, CYANOCOBALAMIN, VITAM IN B12 (CYANOCOBALAMIN)	RE-XITE-D3-TABLET	Tablet	1 tab once or twice daily	13.00	nausea/vomiting, loss of appetite, unusual weight loss, mental/mood changes, signs of kidney problems (such as change in the amount of urine), bone/muscle pain, headache, increased thirst, increased urination, weakness, tiredness,
15	METHYLCOBALAMIN, ALPHA LIPOIC ACID, INOSITOL	RACENERVE FORTE-TABLET	Tablet	1,500 mcg/day in 3 divided doses	189.00	Gastrointestinal discomfort, Rash may be seen after administration, Nausea, Vomiting, Vertigo, Flushing, Postural hypotension, Swelling of part or whole of the body, Sensation of spinning, Urge to vomit, Racenerve Forte Tablet may also cause side-effects not listed here.
16	METHYLCOBALAMIN, AND PREGABALIN.	NEUGABA-M-TABLET	Tablet	Initial: 150 mg/day, may increase to 300 mg/day after a wk. Max: 600 mg/day.	131.00	Dizziness, Sleepiness, Tiredness, Uncoordinated body movements.
17	CALCIUM PANTOTHENATE + MECOBALAMIN + NIACINAMIDE + PYRIDOXINE + THIAMINE MONONITRATE	RACENERVE-200ml SYRUP	Syrup	5ml bd/tds daily	236.00	Headache. Upset stomach or throwing up.
18	NIACINAMIDE(100 MG), THIA	RENERVE PLUS-INJECTION	Injection	500 mcg/day 3 times/wk	117.00	Pain/redness at the injection site, mild diarrhea, itching, or a feeling of swelling all over the body.
19	GABAPENTIN, METHYLCOBALAMIN AND ALPHA LIPOIC ACID.	TRIGABANTIN-100mg TABLET	Tablet	Initial: 300 mg on the 1st day, 300 mg bid on the 2nd day and 300 mg tid on the 3rd day; alternatively, 300 mg tid on the 1st day. Dose may increase in increments of 300 mg every 2-3 days	127.00	Dizziness, Sleepiness, Tiredness, Uncoordinated body movements.
		TRIGABANTIN-300mg TABLET	Tablet	Initial: 150 mg/day, may increase to 300 mg/day after a wk. Max: 600 mg/day.	207.00	
20	PIRACETAM+CITICOLINE	CITINERVE-P-TABLET	Tablet	7.2 g/day in 2-3 divided doses; may increase by 4.8 g every 3-4 days. Max: 24 g/day in 2-3 divided doses	540.00	Weight gain, Hyperactivity, Stomach pain, Nervousness, Diarrhoea, Decreased blood pressure.

21	PIRACETAM + GINKGO BILOBA + VINPOCETIN	COGNIVEL	Tablet	120-240 mg/day in divided doses.	216.00	Weight gain, Stomach upset, Rash, Flushing (sense of warmth in the face, ears, neck and trunk), Nervousness, Abnormality of voluntary movements.
22	PREGABALIN (75MG) + NORTRIPTYLINE (10MG)	PREGADOC-NT TABLET	Tablet	150 mg/day, may increase to 300 mg/day after 3-7 days. Max: 600 mg/day after a 7-day interval.	159.42	Constipation, Difficulty in urination, Weight gain, Sleepiness, Dizziness, Tiredness, Blurred vision, Dryness in mouth, Increased heart rate, Orthostatic hypotension (sudden lowering of blood pressure on standing), Uncoordinated body movements.
23	PREGABALIN (75MG) + NORTRIPTYLINE (10MG)	PREGALIFT-NT-TABLET	Tablet	Initial: 150 mg/day, may increase to 300 mg/day after a wk. Max: 600 mg/day.	143.00	Constipation, Difficulty in urination, Weight gain, Sleepiness, Dizziness, Tiredness, Blurred vision, Dryness in mouth, Increased heart rate, Orthostatic hypotension (sudden lowering of blood pressure on standing), Uncoordinated body movements.
24	CHOLECALCIFEROL	ARACHITOL-6L-I.U-1ml INJECTION	Injection	0.25-1microgram/day	44.15	Arachitol is Injection site reactions (pain, swelling, redness).
25	PYRIDOXINE +CYANOCOBALAMIN+NICOTINAMIDE+THIAMINE +RIBOFLAVIN +PANTHENOL	NEUROBION FORTE-2ml INJECTION	Injection	500 mcg/day 3 times/wk.	11.22	Itching, Diarrhea, Dizziness, Irritable, Sleepiness, Rash, Eczema, Tingling or numbness, Flushing, Increased blood sugar level.
26	MECOBALAMIN,NIACINAMIDE,PYRIDOXINE	NEUROBION PLUS-TABLET**	Tablet	1,500 mcg/day in 3 divided doses	8.62	Loss Of Appetite, Nausea, Vomiting, Diarrhea, Allergic Reactions (skin Rashes), Itching And Other Injection Site Related Side Effects (pain And Blood Clots)

CVS - OTHERS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ADENOSINE	ADENOCOR	Injection	Adult: 6 mg IV bolus	255.00	Flushing, headache, palpitation, hypotension
		ADNEON 6MG - 2ML		Paed: 0.04 - 0.1 mg/kg IV	239.00	
2	AMIODARONE	AMIODON 150 MG - 3 ML	Injection	Adult: 1-2.2g	71.00	Blue-grey discolouration of skin, photosensitivity, peripheral neuropathy, paraesthesia, myopathy, ataxia
		CORDARONE - 3 ML			69.30	

3	ACETAZOLAMIDE	DIAMOX - 250 MG	Tablet	Adult: 250- 1000 mg/day Paed: 10 -15mg/kg/day in 3 divided doses	64.10	Malaise, fatigue, depression, excitement, headache, wt loss, GI disturbances, drowsiness, paraesthesia, diuresis, metabolic acidosis, electrolyte imbalance
4	CHLORTHALIDONE	CTD - 6.25 MG	Tablet	25-50 mg/day up to 100-200 mg/day	40.00	Dry mouth, thirst, weakness, lethargy, drowsiness, restlessness, seizures, oliguria, hypotension, fatigue, muscle cramps and GI disturbances
5	DIGOXIN	LANOXIN - 0.25 MG	Tablet	Rapid digitalisation: Loading dose: 0.75-1.5 mg in the 1st 24 hr Slow digitilisation: 250 mcg 1-2 times/day. Usual maintenance: 125-250 mcg/day.	14.57	Extra beats, anorexia, nausea and vomiting. Diarrhoea in elderly, confusion, dizziness, drowsiness, restlessness, nervousness, agitation and amnesia, visual disturbances
6	DOBUTAMINE	DOBUTAMINE - 250 MG - 5 ML	Injection	Acute heart failure 2.5-10 mcg/kg, up to 0.5-40 mcg/kg according to response. Cardiac stress test As 1 mg/mL soln: 5 mcg/kg/min for 8 mins, increase dose at 5 mcg/kg /min until 20 mcg/kg/min	295.60	increased heart rate and BP, ectopic beats, palpitation. Nausea, headache, chest pain, palpitation, dyspnoea, paraesthesia, legcramps
7	DOPAMINE	DOMIN - 200 MG / 5 ML	Injection	Initial: 1-5 mcg/kg/min, up to 5-10 mcg/kg/min according to patient's response. For seriously ill patients: up to 20-50 mcg/kg/min may be used.	30.80	Nausea, vomiting, tachycardia, ectopic beats, palpitation, anginal pain, hypotension, vasoconstriction, bradycardia, hypertension
8	ESMOLOL	NEOTACH-100MG - 10ML	Injection	Loading dose (optional): 500mcg/kg over 1 min. Maintenance: 50 mcg/kg/min for 4 min.	228.00	Hypotension, bradycardia, heart block, syncope, peripheral ischaemia, pallor, flushing nausea, vomiting, anorexia, dizziness, somnolence, paraesthesia
		ESOCARD-10ml INJECTION	Injection		220.00	Nausea, Headache, Fatigue, Constipation, Diarrhoea, Dizziness, Cold extremities.
9	FRUSEMIDE	LASIX - 40 MG	Tablet, Injection	20-80 mg	7.40	Hyponatraemia, hypochloaemic

		LASIX - 2 ML			3.31	alkalosis, hypokalaemia, headache, drowsiness, muscle cramps, hypotension,
		LASIX-4ml INJECTION			6.72	
	FRUSEMIDE + SPIRONOLACTONE	LASILACTONE			30.30	
10	HYDROCHLOROTHIAZIDE	HYDRIDE - 25MG	Tablet	Usual: 25-100 mg/day in the morning, may reduce based on response. Severe: Up to 200 mg/day	17.43	Electrolyte disturbances, weakness, hypotension, pancreatitis, jaundice, diarrhoea, vomiting, sialadenitis, cramping, constipation
11	ISOSORBIDE DINITRATE	ISORBITRATE - 10 MG	Tablet	Adult: PO Long term management of angina 20-120 mg/day in divided doses. Max: 240 mg/day.Heart failure 30-160 mg/day in divided doses. Max: 240 mg/day. 25-100mg od	39.79	Hypotension, tachycardia, flushing, headache dizziness, palpitation, syncope, confusion. Nausea, vomiting, abdominal pain. Apprehension, restlessness, weakness and vertigo.
		SORBITRATE - 5 MG			45.90	
		ISOLAZINE - 20MG			6.55	
		SORBITRATE - 10 MG			0.78	
12	ISOSORBIDE MONONITRATE	NITROFIX - 10MG	Tablet	Adult: PO 20 mg 2-3 times/day. Usual range: 20-120 mg/day.	21.40	Hypotension, tachycardia, flushing, dizziness, palpitation, syncope, Nausea, vomiting, abdominal pain. Restlessness, weakness and vertigo
		NITROFIX SR - 30MG			54.00	
13	ISOXSUPRINE HYDROCHLORIDE	DUVADILAN -	Tablet	10-20 mg 3-4 times/day	83.00	Hypotension, dizziness, palpitation vomiting, abdominal distress, severe rash, flushing, tachycardia
14	IVABRADINE	IVABRAD - 5 MG	Tablet	2.5 - 7.5 mg BD	141.00	Luminous phenomena (phosphena), blurred vision, bradycardia, 1st degree AV-block, ventricular extrasystole, supraventricular extrasystoles, palpitations, headache, dizziness, nausea, constipation, diarrhoea
15	MANNITOL	MANNITOL - 100 ML	Injection	iv 0.25-2 g/kg	36.75	Fluid and electrolyte imbalance; (with high doses). Nausea, vomiting, headache, dizziness, convulsions, fever; tachycardia, chest pain; blurred vision thirst;
16	METOLAZONE	ZYTANIX-2.5MG	Tablet	2.5-5 mg/day	119.20	Chest pain, palpitation, necrotising angitis, orthostatic hypotension, syncope, venous thrombosis, vertigo, volume depletion; depression

17	NEBIVOLOL HCL	NEBILONG 5MG	Tablet	1.25-5 mg once	95.00	Headache, insomnia, chest pain, dyspnoea, rash, and peripheral oedema, hypercholesterolaemia, hyperuricaemia
18	NICORANDIL	KORANDIL - 5MG	Tablet	10-20 mg twice daily	107.00	Headache (usually transitory), flushing, dizziness, nausea, vomiting and weakness. Hypotension and reflex tachycardia at high doses.
19	NITROGLYCERIN	ANGIPLATT - 2.5MG	Tablet, Injection, capsules	inj: 5-25 mcg/min. Usual: 10-200 mcg/min. Max: 400 mcg/min PO: Extended release Up to 12.8 mg 3 times/day.	108.00	Facial flushing, dizziness, tachycardia, throbbing headache and tolerance. Large doses can cause vomiting, restlessness, hypotension, syncope, rarely cyanosis and methaemoglobinaemia, impaired respiration, bradycardia.
		ANGIPLATT - 6.5MG			135.00	
		NITROCONTIN - 2.6 MG			160.60	
		NITROCONTIN - 6.4 MG			189.50	
		NITROGLYCERIN 5 ML			41.00	
20	NORADRENALINE	NORAD - 2 ML	Injection	8-12 mcg/min, up to 8-30 mcg/min in refractory shock. Adjust according to BP response. Maintenance: 2-4 mcg/min	186.40	Hypertension, headache, bradycardia, peripheral ischaemia, arrhythmias, skin necrosis, dyspnoea, respiratory difficulty.
		ADRENOR - 2 ML			173.00	
21	PENTOXIFYLLINE EX.R	TRENTAL - 400 MG	Tablet	400 mg 3 times/day	38.81	Nausea, vomiting, dizziness, headache, flushing; angina, palpitations; occasional cardiac arrhythmias; hepatitis, jaundice.
22	SILDENAFIL	ASSURANS - 20 MG	Tablet	Adult: PO 50 mg about 1 hr before sexual intercourse.	165.00	Headache, flushing, AV block, migraine, respiratory tract infection, angina pectoris, syncope, tachycardia, postural hypotension, MI, cerebral thrombosis
23	SPIRONOLACTONE	ALDACTONE - 100 MG	Tablet	100 mg/day, up to 400 mg/day	81.00	Fluid or electrolyte imbalance, GI upset, gynaecomastia, drowsiness, headache, hyponatraemia; tachycardia, hypotension, oliguria, hyperkalaemia
		ALDACTONE - 25 MG			30.50	
24	TADALAFIL	PULMOPRESS - 20MG	Tablet	10 mg once daily	275.00	Headache, flushing, angina pectoris, respiratory tract infection, AV block, migraine, syncope, tachycardia, postural hypotension, MI, cerebral thrombosis
25	TOLVAPTAN	NATRISE - 30MG	Tablet	15-60 mgOD	756.00	polyuria, xerostomia, increased thirst, and polyuria
26	TORSEMIDE	DYTOR - 10MG	Tablet	2.5-5 mg once daily. Max: 5 mg/day	55.00	Electrolyte disturbances e.g. hypokalaemia, dehydration, dry mouth, headache, dizziness
		DYTOR - 2 ML			13.00	

		DYTOR - 5 MG			33.75	mouth, headache, dizziness, hypotension
27	TRIMETAZIDINE	FLAVIDON-MR	Tablet	40-60 mg/day	99.50	GI disturbances, vomiting, nausea.
28	VASOPRESSIN	CPRESSIN - P (VASOPRESSIN)	Injection	Adult: IV Initial control of 20 u. variceal bleeding As argipressin:	167.00	Pallor, vomiting, nausea, belching, abdominal cramps, tremour, pounding headache, vertigo, fluid retention, hyponatraemia, hypersensitivity
				IM/SC Cranial diabetes insipidus As argipressin: 5-20 u 4 hrly.		
29	LABETALOL	LABEBET-100mg TABLET	Tablet	Initially 100 mg twice daily, dose to be increased at intervals of 14 days; usual dose 200 mg twice daily, increased if necessary up to 800 mg daily in 2 divided doses, to be taken with food, higher doses to be given in 3-4 divided doses; maximum 2.4 g per day	136.00	Excessive tiredness, Tingling scalp or skin, Dizziness, Lightheadedness, Upset stomach, Stuffy nose, Fatigue.
		ALPHADOPA-L-TABLET			12.20	Dizziness, Allergic reaction, Ejaculation disorder, Increased liver enzymes, Erectile dysfunction, Difficulty in urination.
30	AZILSARTAN MEDOXOMIL	AZILDAY-80mg TABLET	Tablet	20-40mg once daily increased upto 80mg once daily	200.00	Dizziness, Back pain, Sinus inflammation, Increased potassium level in blood.
		AZTRIC-40mg TABLET			99.90	
31	PROPRANOLOL	BETACAP-TR-20mg CAPSULE	Capsule	40-80 mg per day	19.48	Tiredness, Cold extremities, Nightmare, Stomach pain, Diarrhoea, Slow heart rate.
		BETACAP-TR-40mg CAPSULE			37.79	
32	CLIDINIUM + CHLORDIAZEPOXIDE + DICYCLOMINE	NORMOXIN-TABLET	Tablet	1-2 tabs 3-4 times /day	19.50	Nausea, Constipation, Nervousness, Dryness in mouth, Difficulty in urination, Confusion, Sleepiness, Weakness, Dizziness, Blurred vision, Coordination impaired.
33	TOLVAPTAN (15MG)	RESODIUM-15mg TABLET	Tablet	Initial dose: 15 mg orally once a day, Maintenance dose: Increase the dose to 30 mg once a day, maximum of 60 mg once a day,	575.55	Headache, Dizziness, Frequent urge to urinate, Nocturia (increased urge to urinate during the night), Dryness in mouth, Fatigue, Polyuria, Diarrhoea, Increased thirst.
34	ISOSORBIDE DINITRATE (10MG)	SORBITRATE-10mg TABLET		10-60mg per 4-6hrs	0.78	Headache, dizziness, lightheadedness, nausea, and flushing.

35	TAMSULOSIN + DUTASTERIDE	TAMGRESS-D-0.4mg TABLET	Capsule	0.4mg once daily increased to 0.8mg	180.00	Impotence, Decreased libido, Breast pain, Ejaculation disorder, Dizziness.
36	TAMSULOSIN (0.4MG)	TAMMAX-0.4mg CAPSULE	Tablet	0.4mg once daily increased to 0.8mg	63.00	headache, dizziness, runny nose, diarrhea, cough, decreased semen, loss of energy or muscle strength, sore throat or trouble swallowing.
37	TAMSULOSIN (0.4MG)	TASULIN-0.4mg TABLET	Tablet	0.4mg once daily increased to 0.8mg	9.50	Retrograde ejaculation, Dizziness, Nasal congestion.
38	TOLTERODINE (4MG)	TOLU-XR-4mg TABLET	Tablet	2mg bid	203.50	Dryness in mouth, Constipation, Headache, Dizziness, Sleepiness, Blurred vision, Dry skin.
39	OXYBUTYNIN (5MG)	TROPEN-5mg TABLET	Tablet	5-10mg QID	125.00	ry mouth, dry eyes, blurred vision, mild constipation, diarrhea, nausea, mild stomach pain or upset, dizziness, drowsiness, weakness, headache, sleep problems.
40	TADALAFIL (10MG) + DAPOXETINE (30MG)	UPHOLD-TABLET	Tablet	1-2 tab/day	256.00	Nausea, Vomiting, Indigestion, Headache, Flushing (sense of warmth in the face, ears, neck and trunk), Dizziness, Nose bleed, Dryness in mouth.
41	LEVOSIMENDAN	VIMENDAN-12.5mg INJECTION	Injection	12.5mg	384.00	Decrease in blood pressure, Headache, Nausea, Insomnia (difficulty in sleeping), Hypokalemic (decreased potassium level in blood) alkalosis, Increased heart rate, Constipation, Irregular heart rate.
42	AMIODARONE (100MG)	CORDARONE-100mg TABLET	Tablet	100-200mg/day	58.00	Tiredness, Blurred vision, Photosensitivity, Nausea, Coordination impaired, Peripheral neuropathy (tingling and numbness of feet and hand), Alterations in thyroid hormone levels, Altered taste, Constipation, Abnormal vision, Lung damage, Tremor.

HORMONES & THEIR ANTAGONIST

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	BETAMETHASONE DIPROPIONATE	STERNON - 30ML LOTION	Tablet, injection, lotion	Adult: 0.5 - 9 mg	75.00	peptic ulceration, acne eruptions, glaucoma, osteoporosis
		BETSONE - 1ML		Paed: 17.5 mcg/kg	4.24	

		BETNESOL - 1 ML		(Adrenocortical insufficiency)	4.40	
		BETNESOL			8.10	
2	BETNOVATE VALERATE CREAM I,P	BETNOVATE - 20GM CREAM	Cream	apply to affected area 2-3 times per day	20.00	Itching, redness
3	CABERGOLINE 0.25 MG	CABGOLIN - 0.25 MG	Tablet	0.25 mg orally twice a week. Dosage may be increased by 0.25 mg twice a week.	132.00	Decrease in BP, dizziness, vertigo, headache, nausea, sleeplessness, abdominal pain, dyspepsia, gastritis, weakness, fatigue, constipation, vomiting, breast pain, hot flushes
4	CARBIMAZOLE	NEO - MERCAZOLE - 10 MG	Tablet	Adult: Initial: 15-40 mg/day. Maintenance: 5-15 mg/day. Paed: 1-2 mg/kg./day	397.95	Mild leucopaenia, sore throat, mouth ulcers, fever, bruising, malaise; rarely cholestatic jaundice; nephrotic syndrome
		NEO - MERCAZOLE - 5 MG			204.75	
5	CARBOPROST TROMETHAMINE	PROSTODIN - 1 ML	Injection	Adult: IM Pregnancy termination in the 2nd trimester 250 mcg, repeat at 1.5-3.5-hrly depending on response. Max: 12 mg. Postpartumhaemorrhage 250mcg at 15-90 min intervals. Max:2mg.	146.70	Severe CVS disorders including fatal hypotension, tachypnea, pyrexia and MI (in patients with high risk factors) following intra-amniotic or intravaginal admin. Convulsions and ECG changes, amniotic fluid embolism and uterine rupture, foetal distress and rarely foetal death during induction of labour.
6	CHORIONIC GONADOTROPHIN 5000 I.U	HUCOG - 5000	Injection	500-4,000 u 3 times/wk.	501.00	Headache, irritability, restlessness, fatigue, depression, oedema; precocious puberty, gynaecomastia
7	CLOMIFENE	FERTYL-M -25MG	Tablet	50 mg/day for 5 days starting on the 5th day of menstrual cycle	4.24	Ovarian enlargement; abdominal pain and bloating; blurred vision; hot flushes; breast discomfort; depression; multiple or ectopic pregnancies;
		FERTYL-M -25MG			127.08	
		SIPHENE 50 MG			45.75	
		SIPHENE-100 MG			70.40	
		CCQ-100mg TABLET			186.00	
		CCQ-50mg TABLET			192.50	
8	CYPROTERONE & ETHINYLOESTRADIOL	KRIMSON-35	Tablet	Hirsutism 2 mg/day (w/ ethinylestradiol)for 21 days of the menstrual cycle. Palliative treatment for prostatic carcinoma Initial: 300 mg/day in 2-3 divided	260.00	Inhibits spermatogenesis, reduces volume of ejaculate, causes infertility, produces abnormal spermatozoa, gynecomastia and enlargement of mammary glands.

				doses. Maintenance: 200-300 mg/day.		
9	DEFLAZOCORT	DEZERT 6 MG	Tablet	120 mg/day. Maintenance: 3-18 mg/day.	87.00	GI disturbances, musculoskeletal, endocrine, neuropsychiatric, ophthalmic, fluid and electrolyte disturbances; susceptible to infection, impaired healing
		DFZ - 6 MG			85.00	
10	DEXAMETHASONE	DECDAK - 0.5 MG	Tablet, Injection	0.75-9 mg/day in 2-4 divided doses.	2.10	peptic ulceration, acne eruptions, glaucoma, osteoporosis
		DEXOREN - S 5ML	Drops		11.91	
		DEXONA - 2ML			5.61	
		DECMAX-8mg TABLET(DEXAMETHASONE)	Tablet		64.00	
11	DINOPROSTONE	PRIMIGYN - 0.5MG GEL	Gel	500 mcg, may repeat 6 hr later if needed. Max: 1.5 mg/24 hr.	259.00	Hypersensitivity reactions, nausea, vomiting, diarrhoea, abdominal pain; flushing, shivering, headache, dizziness; hypertension; convulsions, ECG changes
		CERVIPRIME GEL - 30 GM			260.50	
12	ESTRADIOL VALERATE	PROGYNOVA - 1 MG	Tablet	10-20 mg OD-TDS	178.00	GI disturbances, genitourinary changes, haematologic disorders, CV and CNS effects, endocrine and metabolic disorders, cholestatic jaundice, local skin reactions, chorea
13	ESTRIOL	EVALON - 15GM CREAM	CREAM	1 applicator upto the ring /day for 2-3 weeks;then 1 applicator upto the ring twice a week	289.00	
14	FLUDROCORTISONE ACETATE	FLORICOT	Tablet	50-300 mcg/day	115.00	Hypertension, sodium and water retention, potassium loss, dizziness, itching, skin rash, headache, convulsions, CHF, muscle weakness, hyperglycaemia
15	FLUNARIZINE DIHYDROCHLORIDE	SIBELIUM 10MG	Tablet	5-10 mg/day at bedtime	89.00	Drowsiness. Rarely wt gain, headache, depression, gastric pain, dry mouth, insomnia, extrapyramidal reactions, galactorrhoea.
		SIBELIUM 5MG			52.50	
16	ISOPHANE INSULIN	HUMAN INSULATARD	Injection	0.3-1 IU/kg/day; may be given in 1-2 divided doses	149.40	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision.
17	IBANDRONIC ACID	IDROFOS - 150 MG	Tablet	2 -6 mg	220.00	Fever; influenza-like syndrome; electrolyte disturbances;GI disturbances;musculoskeletal pain; headache; rash, allergic dermatitis; anaemia; decreased alkaline
	IBANDRONIC ACID + CALCIUM CARBONATE	ROCKBON KIT			320.00	
18	LETROZOLE	LETROZ - 2.5 MG	Tablet	2.5 mg once daily	172.00	"Hot flushes, arthralgia, nausea,

						vomiting, fatigue, dizziness, headache, dyspepsia, constipation,
						increased sweating, rash, peripheral oedema, osteoporosis, musculoskeletal pain, vaginal irritation.
		CHEMLET-2.5mg TABLET (LETROZOLE)	Tablet	2.5 mg once daily	51.00	"
19	LEVONORGESTREL & ETHINYLOGESTRADIOL	OVRAL - L	Tablet	Adult:PO Monophasic combined OC: As tab containing levonogestrel 150-250 mcg and and ethinylestradiol 30 mcg: 1tab once daily. Triphasic combined OC: Levonorgestrel 50-125mcg+ ethinylestradiol 30-40 mcg: 1 tab once daily.	70.56	Menstrual irregularities; headache, dizziness; breast discomfort; gynaecomastia; depression; disturbance of appetite; wt changes; fluid retention; oedema; changes in libido; hair loss or hirsutism; GI disturbances
20	MEDROXY PROGESTERONE ACETATE	MEPRATE - 10MG	Tablet	5 to 10 mg orally once a day	60.05	Depression, fluid retention. Fatigue, insomnia, dizziness, headache, nausea;breasttenderness; wt gain/loss, anorexia; cholestatic jaundice; pain at Inj site.
		DEVIRY - 10 MG			60.59	
21	CARBOPROST	CABOPROST-1ml INJECTION	Injection	Initial: 250 mcg. Alternatively, initiate w/ test dose of 100 mcg. May be repeated at 1.5- to 3.5-hr intervals depending on uterine response. May be increased to 500 mcg if uterine contractility is inadequate. Max: 12 mg. Max duration: 2 days.	192.60	Severe pelvic pain, cramping, or vaginal bleeding, high fever, feeling light-headed or short of breath, severe nausea, vomiting, or diarrhea.
22	EVENING PRIMOSE OIL	PRIMOSA-1gm CAPSULE	Capsule	1 cap /day	242.54	Headache, Diarrhea, Cephalalgia, Upset stomach, Nausea.
		PRIMOSA-500mg CAPSULE		2cap/day	194.81	
23	MEGESTEROL ACETATE	ENDACE-40mg TABLET(MEGESTEROL ACETATE)	Tablet	40 mg 4 times daily or 160 mg once daily.	311.60	hair loss, increased appetite, mild blood pressure increase, mood changes, nausea or vomiting, swelling of the face, ankles, or feet, weight gain.
24	METHYL PREDNISOLONE	IVEPRED -16 MG	Tablet, Injection	2-60 mg/day in 1-4 divided	99.00	peptic ulceration, acne eruptions, glaucoma, osteoporosis
		IVEPRED -8 MG	Syrup, Drops	doses	54.00	
		MEDROL 4 MG			41.00	
		IVEPRED -125MG			288.00	

		IVEPRED -40MG			150.00	
		PREDMET -4MG			33.50	
		DEPO-MEDROL - 2 ML			1271.00	
		NEODROL - 500 MG			270.00	
		SOLU - MEDROL - 1 GM			1271.00	
		PREDNISOLONE - 10 MG			15.98	
		PREDNISOLONE - 5 MG			9.45	
		PREDMET - 10ML			18.90	
		KIDPRED - 60ML			27.17	
25	MISOPROSTOL	MISOPROST- 200 MG	Tablet	200-800 mcg/day in 2-4 divided doses	87.00	Diarrhoea, abdominal pain, dyspepsia, constipation, flatulence, nausea and vomiting; abnormal vaginal bleeding, cramps, increased uterine contractility, headache.
		ZITOTEC - 200MCG			78.50	
26	MIXTURE OF INSULIN & PROTAMINECRYSTALLISED	NOVOMIX FLEXPEN	Injection	0.2 microgram/kg body wt/day. Titrate as reqd.	611.31	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision.
27	NANDROLONE DECANOATE	DECA DURABOLIN - 25 MG	Injection	25-100 mg 3-4 wkly	157.00	Male: Postpubertal: Acne, gynaecomastia, bladder irritability, priapism, insomnia, chills, decreased libido, hepatic dysfunction, nausea, diarrhoea, prostatic hyperplasia. Prepubertal: Acne, virilism, chills, insomnia, diarrhoea, hyperpigmentation, nausea. Female: Virilism, insomnia, iron deficiency anaemia, hepatic hypercalcaemia, nausea, diarrhoea, chills, dysfunction.
		DECA DURABOLIN - 50 MG			256.00	
28	NORETHINDRONE ACETATE	REGESTRONE - 5 MG	Tablet	2.5-10 mg/day	58.60	Mental depression, cholestatic jaundice, porphyria, epilepsy, migraine, headache, breast discomfort, dizziness, nausea and vomiting, changes in libido
		PRIMOLUT - N			58.73	
29	OCTREOTIDE	OCTRIDE 50 MCG - 1 ML	Injection	25 mcg/hr for 48 hr	305.00	Local pain, stinging, tingling at site of inj; anorexia, nausea, vomiting, abdominal pain, bloating, flatulence, loose stools, steatorrhoea
30	OXYTOCIN	EVATOCIN - 1 ML	Injection	10-40 u by infusion at a rate to control uterine atony	17.97	transient hypotension, reflex tachycardia; nasal irritation, rhinorrhoea, lachrymation
31	PROGESTERONE	SUSTEN - 100 MG	Capsule	200 - 400 mg/day	108.00	GI disturbances, appetite/wt change, fluid retention, oedema, acne, skin rash, urticaria, depression, headache, fever, fatigue, breast changes, hirsutism, changes in libido, altered
		SUSTEN - 200 MG			201.00	

						menstrual cycles or irregular menstrual bleeding (rare).
32	SOLUBLE INSULIN	HUMAN ACTRAPID	Injection	20 u, then 6 microgram/hr until blood glucose drops to 10 mmol/L, when the dose is given 2 hrly	149.40	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred
33	THYROXINE SODIUM	ELTROXIN - 50 MCG	Tablet	25-100 mcg	68.94	Vomiting & Diarrhoea
		ELTROXIN- 100 MCG			121.60	
		THYRONORM - 100 MCG			128.65	
		THYRONORM - 150 MCG			187.00	
		THYRONORM - 25 MCG			164.00	
		THYRONORM - 50 MCG			116.55	
		THYRONORM - 75 MCG			126.00	
34	TRIAMCINOLONE ACETONIDE	KENACORT - 40 MG	Injection ,Paste	Apply 2-3 times daily. Rub in gently.	99.50	peptic ulceration, acne eruptions, glaucoma, osteoporosis
		KENACORT - 5GM ORAL PASTE			48.00	
		KENACORT INJECTION - 10 MG			49.00	
35	NATURAL MICRONISED PROGESTERONE	HYPERTON 6-5gm OINTMENT	Ointment	max 5mg per oz not exceeding 2oz per month	87.00	Fatigue, Sleepiness, Headache, Abdominal pain.
36	LEVONORGESTREL	MIRENA IUS-52mg	IUD	Must be removed at the end of 5th year	3509.00	missed periods (amenorrhea), bleeding and spotting between periods, heavier bleeding during the first few weeks after device insertion, abdominal/pelvic pain, ovarian cysts, back pain, headache/migraine, nervousness, dizziness, nausea, vomiting, bloating, breast tenderness or pain, weight gain, changes in hair growth, acne, depression, changes in mood, loss of interest in sex, itching or skin rash, and puffiness in the face, hands, ankles, or feet.
37	METHYLPREDNISOLONE (4MG)	NEODROL-40mg INJECTION	Injection	10-500 mg/day	48.75	Nausea, vomiting, heartburn, headache, dizziness, trouble sleeping, appetite changes, increased sweating, acne, or pain/redness/swelling at the injectionsite
38	PREDNISOLONE (TOPICAL) (1% W/V)	PRED FORTE-10ml DROPS	Drops	Adult: PO Allergic and inflammatory disorders 5-60 mg/day, in 2-4 divided doses	34.18	Eye irritation, Watery eyes, Burning sensation.
39	CONJUGATED ESTROGENS (0.625MG)	PREMARIN-0.625mg TABLET(28'S)	Tablet	.45 mg/day, up to 1.25 mg/day. Attempt to discontinue medication at 3-6-mth intervals.	709.21	Headache, Vaginal bleeding, Fluid retention, Vomiting, Nausea, Stomach cramp, Breast pain, Hair loss.

40	PROGESTERONE (NATURAL MICRONIZED) (200MG)	PRENONE-200mg SOFT GELATIN CAPSULE	Capsule	400 mg daily for 10 days.	28.88	Fatigue, Sleepiness, Headache, Abdominal pain.
41	ESTRADIOL (2MG)	PROGYNOVA-2mg TABLET	Tablet	1-2mg per day	384.00	Headache, Nausea, Breast tenderness, Stomach pain.
42	MIFEPRISTONE + MISOPROSTOL	UNWANTED KIT-TABLET	Tablet	600 mg as a single dose, followed by misoprostol 400 mcg orally or gemeprost 1 mg vaginally 36-48 hr later	385.89	Nausea, Vomiting, Diarrhoea, Stomach cramp, Uterine contractions, Heavy menstrual periods.
43	TERIPARATIDE-250MCG/ML	BONISTA CATRIDGE INJECTION	Injection	20mcg /day	7117.50	Joint pain, Injection site reactions (pain, swelling, redness), Nausea, Leg cramps.
44	CORTICOTROPIN (60IU)	ACTON PROLONGTON 60I.U/ml-5ml INJECTION(ACTH)	Injection	15-30IU (0.25-0.5ml) SC every 2 days	1794.03	Fluid retention, Increased appetite, Glucose intolerance, Weight gain, Increased blood pressure, Mood changes, Behavioural changes.

HYPOLIPIDEMIC DRUGS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ATORVASTATIN	ATOCOR - 10 MG	Tablet	Adult: 10-80 mg /day children > 6 years same as adults	70.75	GI disturbance , flu syndrome, back pain, allergic reactions, asthenia
		AVAS - 10 MG			65.90	
		AZTOR - 10 MG			68.43	
		AVAS - 20 MG			187.00	
		AZTOR - 20 MG			180.00	
		TG-TOR - 40MG			211.35	
		STORVAS - 10 MG			105.93	
		AVAS-80mg TABLET			347.00	
2	FENOFIBRATE	FENOLIP - 145MG	Tablet	Adult: PO Standard micronised formulation: Initial: 67 mg tid or 200 mg once daily, may adjust to 67 mg 2-4 times/day or 267 mg once daily. Non-micronised formulation: Initial: 200-300 mg/day in divided doses. Usual range: 200-400 mg/day. Formulation w/ improved	121.00	Abnormal LFTs , increased creatini kinase concentrations, flu syndrome Mild to moderate decreases in Hb, haematocrit and leukocyte counts. Potentially Fatal: Hepatitis, cholecystitis, pancreatitis, rhabdomyolysis.
	ATORVASTATIN + FENOFIBRATE	TG-GOAL -145 MG			130.50	
3	ROSUVASTATIN	ROZAVEL - 10MG	Tablet	Adult: 5-20mg OD	128.00	Headache dizziness, Gi upset, peripheral edema, insomnia, proteinuria
		ROZAVEL - 20MG			223.00	
		ROZAVEL - 40MG			345.00	
		ROZAVEL - 5MG			68.00	

		ROZAVEL-F--5MG			10.50	
	ROSUVASTATIN + FENOFIBRATE	ROZAVEL-F			168.00	
4	ASPIRIN / ACETYLSALICYLIC ACID + ATORVASTATIN + CLOPIDOGREL	CLAVIX GOLD-10mg CAPSULE	Capsule	Loading dose: 4 tab; maintenance: 1 tab/day	57.83	Diarrhoea, Stomach pain, Constipation, Indigestion, Weakness, Increased glucose level in blood, Headache, Dizziness, Muscle pain,
		CLAVIX GOLD-20mg CAPSULE	Capsule	Loading dose: 4 tab; maintenance: 1 tab/day	78.05	
5	ATORVASTATIN (10MG) + CLOPIDOGREL	AVAS C.V-10/75mg TABLET	Capsule	10 or 20 mg once daily, may be adjusted at 4-wk interval	94.01	Stomach pain, Diarrhoea, Indigestion, Constipation, Headache, Dizziness, Muscle pain, Weakness, Increased glucose level in blood, Increased liver enzymes, Increased bleeding tendency.
6	ATORVASTATIN-20MG+CLOPIDOGREL-75MG+ASPIRIN	ECOSPRIN GOLD-10mg TABLET	Tablet	1tab/day may be increased according to requirements	81.94	Diarrhoea, Stomach pain, Constipation, Indigestion, Weakness, Increased glucose level in blood, Headache, Dizziness, Muscle pain, Increased liver enzymes, Increased bleeding tendency.
		ECOSPRIN GOLD-20mg TABLET		1tab/day may be increased according to requirements	110.50	
		ECOSPRIN-325mg TABLET		1tab/day may be increased according to requirements	11.13	

MUSCLE RELAXANTS, ANAESTHETIC & EMERGENCY

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ADRENALINE	ADRENALINE - 1 ML	Injection	0.2-0.5 ml of 1: 1000 solution (0.2-0.5 mg)	15.00	Cardiac arrhythmias, rise in BP, angina, tachycardia, anxiety
		VASOCON (ADRENALINE)			20.00	
2	ATROPINE SULPHATE	TROPINE - 1 ML	Injection	Adult: IM/IV 0.6-m 2mg Paed: < 5kg: 0.2 mg/kg > 5kg : 0.1-0.2 mg/kg	4.60	Rapid pulse, flushing, shortness of breath, urinary retension, GI upset
		ATROPINE SULPHATE -100ML			98.00	
3	BUPIVACAINE HYDROCHLORIDE	ANAWIN - 0.25% - 20 ML	Injection	Adult: 225mg with adrenaline & 175 mg without adrenaline by infiltration	51.00	Bradycardia, hypotension, HTN, nausea, vomiting, diplopia, neuropathy, peripheral nerve injury
		ANAWIN - 0.5% - 20 ML			77.00	
		ANAWIN HEAVY - 4ML			24.99	
4	DEXMEDITOMIDINE	DEXTOMID 100MCG - 1ML	Injection	Adult: a loading infusion of one mcg/kg over 10 minutes. a maintainance infusion of 0.2 to 0.7 mcg/kg/hour.	326.00	bradycardia, hypotension, nausea, agitation, tachycardia, hypertension,xerostomia.
		DEXTOMID 50MCG - 0.5ML			195.00	
5	GLYCOPYRROLATE	PYROLATE - 1 ML	Injection	IM/IV 200- 400 mcg every 3-4 hrs	17.80	Xerostomia; loss of taste, nausea, vomiting, constipation, reduced sweating; urinary hesitancy and retention
6	LIDOCAINE	LOX - 10 % - SPRAY	Spray, Injection,	Adult: 1-1.5 mg/kg	393.00	Arrhythmia, bradycardia, arterial

		LOX - 4% TOPICAL - 30 ML	Gel, Cream	Paed: 0.5-1mg/kg	34.00	spasms, CV collapse, oedema, flushing, heart block, hypotension, sinusnodesuppression, agitation
		ANORELIFE - 30GM GEL			107.00	
		TOPLAP-10GM CREAM			210.00	
7	LIGNOCAINE HYDROCHLORIDE	LOX - 2% JELLY 30 GM	Jelly, injection	Adult: 300mg Paed: 3mg/kg	40.00	Arrhythmia, bradycardia, arterial spasms, CV collapse, oedema, flushing, heart block, hypotension, sinusnodesuppression, agitation
		LOX-2% - 30ML			33.50	
		LOXICARD - 2% - 50ML			55.00	
		LOX-2% AND ADRENALINE-30 ML			29.75	
8	MAGNESIUM SULPHATE	MAGNEON - 2 ML	Injection	Adult: im/iv 1-2 G	11.00	Flushing, sweating, hypotension, hypothermia
		MAGNESIUM (SULP) PACKET - 20GM		Paed: IM/IV 25-100mg/kg/ dose	10.00	
9	MIDAZOLAM	MEZOLAM - 1 ML	Injection	0.05 - 0.15 mg/kg	27.70	Decreased respiratory rate, hypotension, bradycardia, GI upset
		MEZOLAM - 5 ML			32.50	
10	NEOSTIGMINE METHYLSULPHATE	MYOSTIGMIN - 1 ML	Injection	Adult:IM/IV/SC: 0.04-0.05 mg/kg Paed: IM/IV/SC- 0.01-0.04 mg/kg	4.45	Salivation, GI upset, drowsiness, visual changes, arrythmia
11	POTASSIUM CHLORIDE	POTCL 10 ML		Injection, syrup	Adult: IV infusion 10-20 meq/hr	23.60
		POTKLOR - 200 ML	Paed: IV infusion 10-20 meq/hr		68.68	
12	DI SODIUM HYDROGEN CITR	CITRALKA-100ml SYRUP	Syrup	PO 15-30 mL bid-tid.	65.14	Mild diuresis, stomach cramps, flatulence, diarrhoea, GI ulceration, hypokalaemia, metabolic alkalosis, tiredness, dyspnoea, anxiety, mood swings.
13	MAGNESIUM CITRATE, AND PYRIDOXINE.	ALKASTON B6-200ml SYRUP	Oral solution	less than 6yrs 0.5 mL /Kg orally upto max/; 200mL repeat every 4-6hrs util stool are clear ; 6-12yrs - 100-150mL/DAY once;Adult -240mL once/day	137.00	bowel movement within 6 hours after taking the medicine; pain with bowel movements, rectal bleeding; watery diarrhea, nausea, vomiting, severe stomach pain;painful or difficult urination;
14	MAGNESIUM CITRATE 375 MG+POTASSIUM CITRATE	KMG-CIT WITH B6-200ml SYRUP	Oral solution	less than 6yrs 0.5 mL /Kg orally upto max/; 200mL repeat every 4-6hrs util stool are clear ; 6-12yrs - 100-150mL/DAY once;Adult -240mL once/day		Bloating Confusion Anxiety Uneven heart rate
15	POTASSIUM CITRATE- 1100MG/5ML+MAGNESIUM CITRATE	KMAC-200ml SYRUP	Oral solution	less than 6yrs 0.5 mL /Kg orally upto max/; 200mL repeat every 4-6hrs util stool are clear ; 6-12yrs - 100-150mL/DAY once;Adult -240mL once/day	134.00	Bloating Confusion Anxiety Uneven heart rate

16	PROPOFOL	NEOROF 10MG -10ML	Injection	Adult: 1.5 - 2.5 mg/kg IV upto 15 mg/kg/hr Paed: 3 - 3.5 mg/kg upto 12.5 mg/kg/hr	98.00	Nausea, vomiting, hiccough, ophthalmoplegia, anaphylaxis
17	SODIUM BICARBONATE	AUXISODA-IV - 100ML	Injection, Tablet	Adult: 1 mEq/kg then 0.5 mEq/kg at 10 min interval Paed: 1-2 mEq/ kg/ dose	140.00	Metabolic alkalosis, hypernatremia
		AUXISODA-500MG			27.50	
		SODAE (8.4 %)			35.00	
		SODIUM - BI-CARBONATE - 10ML			20.00	
18	SUCCINYLMCHOLINE	SUCOL - 10 ML	Injection	Adult: IV 0.6 mg/kg Paed: IV 1-2 mg Maintenance: 0.3 -0.6 mg/kg	67.00	Bradycardia, tachycardia, hypotension, hyperpertenstion, raised intraocular pressure hyperkalaemia, excessive salivation.
19	VECURONIUM BROMIDE	NEOVAC - 4 MG \2 ML	Injection	Adult: 80 - 100 mcg/kg maintenance: 20 -30 mcg/kg Paed: < 4mth : 10-20mcg/kg > 4 mth: as in adults	135.00	Muscle weakness, paralysis, muscleatrophy (after long term use), hypersensitivity reactions e.g urticaria and erythema.
20	THIOLCHICOSIDE + ACECLOFENAC	THIOPAS - 4 MG	Tablet	Muscle spasms Adult: Initially, 16 mg daily. Intramuscular Muscle spasms Adult: Up to 8 mg daily.	132.00	Photosensitivity reactions.
		THIOPAS-A - 4MG			153.00	
		NEORELAX-A - 4MG			128.00	
		THIOPAS-A - 8MG			233.00	
		NEORELAX-MR			140.00	
		ZERDOL - TH			130.00	
21	THIOPENTONE SODIUM	THIOSOL - 500 MG	Injection	100-150 mg of a 2.5 or 5% solution injected over 10-15 sec repeated every 30-60 sec according to response or as a continuous infusion of a 0.2 or 0.4% solution. Max: 500mg.	47.00	Delirium, GI upset, apnea, laryngospasm
22	ANTI SNAKE VENOM	ANTI SNAKE VENOM - 10 ML	Injection	cobra antivenom 0.6 mg/1 mL, bungarus caeruleus venom antiserum 0.45 mg, daboia russellii antivenin 0.6 mg, echis carinatus venom antiserum 0.45 mg.	950.00	HS reaction
23	ARTARICURARIUM	ARTACIL - 2.5 MG	Injection	Adults: 3-6 mcg/kg Paed: 150 mcg/kg	128.00	Allergic reactions, hypotension, tachycardia, bronchospasm
24	ACETYLCYSTEINE	MUCINAC - 2 ML	Respules, injection, Tablet, Capsule	PO As a mucolytic As loz, granules or effervescent tab: 600 mg daily as a single or in 3 divided doses. Paracetamol poisoning As 5% soln: Initial: 140 mg/kg followed by	52.11	Bronchospasm, angioedema, rash, pruritus, hypotension, HTN, flushing, nausea, vomiting
		MUCOMIX - 2ML			49.95	
		MUCOMIX - 1ML			25.98	
		MUCOMIX			320.00	
		LUMENAC			169.00	

		MUCINAC - 600 MG		/0mg/kg 4 hrly for additional 1 / days	179.00	
25	BACLOFEN	RICLOFEN - 10 MG	Tablet	5 mg 3 times/day for 3 days, increased to 10 mg 3 times /day for 3 days	51.00	Sedation, drowsiness, ataxia, dizziness, headache, confusion, hallucinations, skin reactions, GI
26	BENZOCAINE	MUCOPAIN - 15 GM GEL	Gel	Apply up to 4 times/day	47.00	Skin sensitivity; irritation of conjunctiva, mucosal irritation; reversible brown staining of the teeth; tongue discoloration and burning sensation; transient taste disturbance; parotid gland swelling. Potentially Fatal: Anaphylaxis.
27	CHLORZOXAZONE + PARACETAMOL	MYOSPAS	Tablet	250 - 750 mg QID	84.50	GI upset, hypersensitivity reactions, headache
	CHLORZOXAZONE + PARACETAMOL+ DICLO	MYOSPAZ FORTE			166.50	
28	DICYCLOMINE	SPASMO - PROXYVON - 2ML	Injection, Tablet	10-20 mg 3 times/day. <i>IM</i> 20 mg 4 times/day for 1-2 days.	6.84	Difficulty in accommodation, exacerbation of glaucoma; tachycardia, palpitations, arrhythmias; urinary retention; restlessness; confusion, excitement, hallucination and delirium.
		COLIMEX - SYRUP	Syrup, Drops		29.65	
		CYCLOPAM - 10ML DROPS			31.90	
		CYCLOPAM - 30ML SYRUP			36.30	
		CYCLOPAM			36.90	
29	DROTAVERINE	DROTIKIND - 2 ML	Injection, Tablet	40-80 mg 3 times/day.	16.50	Vertigo, nausea, vomiting, dry mouth.
		DROTIN			16.21	
		DROTIN DS			70.03	
		DROTIN			40.69	
		DROTIN - M			99.73	
30	HYOSCINE BUTYLBROMIDE	HYOCIMAX - (HYOSCINE)	Injection, Tablet	PO Genito-urinary spasm; GI tract spasm As butylbromide: 20 mg 4 times/day. Motion sickness As hydrobromide: 300 mcg 30 mins before journey, then 300 mcg 6 hrly if needed.	11.39	Flushing, postural hypotension, tachycardia, fibrillation. Rarely psychotic reactions. Dizziness, drowsiness, fatigue, headache, memory loss.
		HYOCIMAX - (HYOSCINE)			32.87	
31	MEBEVERINE HCL	COLOSPA - 135 MG	Tablet	100 or 135 mg 3 times/day.	111.87	GI disturbances, dizziness, headache, insomnia, anorexia, decreased heart rate, hypersensitivity reactions.
32	PRALIDOXIME	ALDOPAM	Injection	Adult: IV/IM Used in combination w/ atropine. As Cl,iodide or mesilate: Upon atropinisation, administer 1-2	170.00	Drowsiness, dizziness, visual disturbances, nausea, tachycardia, headache, hyperventilation and muscle weakness.

				g.repeat dose after 1 hr, then 8-12 hrly if needed. Severe: 200-500 mg/hr via continuous infusion, titrate according to response.		
33	TIZANIDINE	TIZAN - 2 MG	Tablet	2 mg once daily	94.00	Drowsiness, fatigue, dizziness, insomnia, headache, anxiety, nausea, dryness of mouth, GI disturbances, hypotension
34	VALETHAMATE BROMIDE	VALETHAMATE - 1 ML	Injection	8-16 mg/day.	15.00	Dry mouth, blurring of vision, difficulty in swallowing and talking, constipation, urinary retention, mydriasis, arrhythmias.
35	GLYCOPYRROLATE, NEOSTIGMINE.	MYO-PYROLATE-5ml INJECTION	Injection	5-10 ml IV over a period of 10-30 secs.	70.20	dryness of the mouth, blurred vision, headaches, clouded thoughts, nervousness, fatigue, dizziness, stomach problems, vomiting and constipation.
36	BOVINE LIPID EXTRACT SURFACTANT SUSPENSION	NEOSURF-5ml	Suspension	adult and children more than 12 yrs: 2cap/3times daily	9202.00	Drowsiness, dizziness, hypotension or a headache
37	LIDOCAINE TOPICAL (25MG/GM) + PRILOCAINE TOPICAL (25MG/GM)	PRILOX-5gm CREAM	cream	Max: 20 g in 24 hr	99.00	Dizziness, drowsiness, vomiting, blurred vision and slurred speech.
38	SEVOFLURANE (99.97%)	SEVORANE-250ml LIQUID	Liquid	0.5-3% with or without Nitrous oxide		Vomiting, Nausea, Hallucination, Abnormal dreams, Nightmare, Confusion, Agitation, Generalized tonic-clonic seizure, Double vision, Increased blood pressure, Tachycardia.
39	BENZOCAINE + CHLORIBUTOL + PARADICHLOROBENZENE + TURPENTINE OIL	SOLIWAX-10ml EAR DROPS	Drops	Instill 5-10 drops into the affected ear 2-3 times a day and allow contact for 10-20 minutes and overnight. Continue for 2-3 days, if necessary.	111.00	Burning/stinging/redness/new pain in or around the ears.
40	ETOMIDATE (20MG/10ML)	TROYMIDATE-10ml INJECTION	Injection	IV 300 mcg/kg over 30-60 sec.	467.50	Nausea, Rash, Vomiting, Abnormality of voluntary movements, Apnea (absence of breathing), Hyperventilation, Thrombophlebitis, Stridor, Myoclonus, Decreased blood pressure.

41	HYDROXYETHYL STARCH(HES) (6GM) + SODIUM CHLORIDE (0.9GM)	VOLUVEN FREEFLEX-6%-500ml(HYD.ETH.STARCH)	I.v fluid	IV Blood loss & hematocrit up to approx 20 mL/kg/day equiv to 1.2 g hydroxyethyl starch/kg/day Patient weighing 70 kg 1,500 mL. Septic shock due to burns Adult Infusion rate approx 25 mL/kg/min & 12.5-25 mL/min or 1,000 mL in 40-80 min.	531.18	Nausea, vomiting, constipation, lightheadedness, dizziness, drowsiness, increased sweating, urinary retention, headache or dry mouth.
		MORPHINE SULPHATE-10mg TABLET		5-20 mg 4 hrly. Extended-release: 5-20 mg 12 hrly.	5.00	
		MORPHINE-10mg/ml-1ml INJECTION(Vermor)		5-10 mg at 1-2 mg/min followed by a further 5-10 mg as necessary	22.00	
		PETHIDINE-50mg/ml-1ml INJECTION(Verpat)		Moderate to severe acute pain 25-100 mg IM/SC inj or 25-50 mg by slow IV inj repeated after 4 hr	49.35	
		FENTANYL-50Mcg/ml-2ml INJECTION(Verfen)		Initial: 50-200 mcg followed by supplements of 50 mcg. Max: 200 mcg. Admin lower infusion rates of 0.05-0.08 mcg /kg/min. Patients w/ assisted ventilation: Initial: 300-3,500 mcg (up to 50 mcg/kg) followed by supplements of 100-200 mcg depending on response. Loading dose (alternatively via bolus): Approx 1 mcg/kg/min given for the 1st 10 min followed by infusion of approx 100 ng/kg/min	46.00	
		ANEKET-250mg-5ml INJECTION		Adult: IV 1-4.5 mg/kg as inj or infuse 0.5-2 mg/kg. IM 6.5-13 mg/kg.	52.65	
42	BUPRENORPHINE	BUPREGESIC-1ml INJECTION	Injection	IM Anaesth premed 300 mcg. IV Perioperative analgesia 300-450 mcg. IM/IV Moderate to severe pain 300-600 mcg 6-8 hrly.	21.00	Nausea, vomiting, dizziness, sweating and drowsiness.

NSAIDS & GOUT

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	PARACETAMOL + ACECLOFENAC	ICONAC - P	Tablets	Adults: 100mg	38.00	Nausea, Vomiting, Perforation, GI ulceration, Bleeding, Bronchospasm, Coagulopathy
		MOVON - P			37.00	
		HIFENAC - 100 MG			31.50	
		HIFENAC - P			54.50	
		HIFENAC -D			68.50	
		DOLOKIN PLUS			23.47	
		ALTRAFLAM - P			47.00	
		ACECLOFENAC + DROTAVERIN			DROLGAN	
ACECLOFENAC+RABEPRAZ OLE CISSUS QUADRANGULARIS EXTRACT MESALAMINE	ALTRADAY-CAPSULE	Capsule	100mg bd	89.00	Nausea, Flatulence, Indigestion, Diarrhoea, Constipation, Flu-like symptoms.	
	CISSACT-TABLET	Tablet	2tabs/bd per day	154.00	headache, intestinal gas, dry mouth, diarrhea, and insomnia.	
	MESACOL-400mg TABLET	Tablet	2.4 g/day in divided doses. Maintenance of remission: 1.2- 2.4 g/day in divided doses	75.93	mild nausea, vomiting, stomach cramps, diarrhea, gas, fever, sore throat, or other flu symptoms, constipation, headache or dizziness, tired feeling or skin rash.	
2	ACETAMINOPHEN	DOLOPAR - 500 MG	Tablet, Syrup,	Adult dose : 500, 650 mg; Paedaitric : 10-15 mg/kg	35.50	Thrombocytopenia, Pancytopenia, Neutropenia
		DOLO 650 MG	Injections, Drops		33.46	
		NEOMOL - 170 MG	Suppositories		33.50	
		NEOMOL - 250 MG			58.00	
		NEOMOL - 80 MG			27.00	
		NEOMOL I.V - 100ML			297.00	
		P-250 - 60ML			35.10	
		FEVASTIN - 2 ML			9.69	
		CALPOL- 15 ML			24.00	
		CALPOL - 500 MG			16.30	
		FEBRAX PLUS			33.20	
		WIKORYL			40.50	
		SINAREST			38.20	
		SINAREST			50.33	
		DOLO-100ml INFUSION			268.00	
		CROCIN-60ml SUSPENSION	suspension		32.25	

	ACETAMINOPHEN + DOMPERIDON	GRENIL		2 tab at the onset of attack, may repeat 4 hrly if needed. Max: 4 g paracetamol/day.	42.50	Irregularity in heart beat A feeling of dizziness or faintness Difficulty in breathing, swelling of the throat or the face Menstrual irregularities Discharge of breast milk from the nipple Swelling of breasts in males
	ACETAMINOPHEN +	CALPOL -T		Adult: PO 0.5-1 g 4-6 hrly. Max: 4 g/day. Rectal As supp: 0.5-1 g 4-6 hrly. Max: 4 g/day. IV 33-50 kg: 15 mg/kg as a single dose, at least 4 hrly. Max: 60 mg/kg/day up to 3 g/day; >50 kg: 1 g as a single dose, at least 4 hrly. Max: 4 g/day.	38.65	Acid or sour stomach Major Common Heartburn Major Common Nausea and Vomiting Major Common Abdominal discomfort Major Less Common Constipation Major Rare Decreased urine output Major Rare Yellow colored eyes or skin Major Rare Skin rash Major Rare
	TRAMADOL	ULTRACET		2 tab 6 hrly. Max: 8 tab/day. Max duration: 5 days.	130.00	dizziness, drowsiness, tired feeling; stomach pain, nausea, loss of appetite; diarrhea, constipation; or sweating
3	DICLOFENAC SODIUM	VOVERAN EMULGEL - 30 GM	Tablet, Injection Gel, Sprays, suppositories, liniment	Adult dose : 50mg	78.90	Odema, Syncope, Nausea, Diarrhoea
		VOVERAN TPM GEL - 30 GM			105.45	
		OXALGIN NANOGEL - 30GM			90.00	
		NOBEL GEL - 30 GM			50.60	
		VOLITRA PLUS			83.00	
		VOVERAN - 50 MG			74.21	
		JUSTIN SUPPOSITORIES - 12.5MG			35.00	
		JUSTIN SUPPOSITORIES - 25MG			37.00	
		JUSTIN SUPPOSITOTIES - 100 MG			52.90	
		VOVERAN - 1 ML			19.47	
		DICLOGESIC			32.85	
		LYSER - D			154.00	
		FENLONG MR LINIMENT - 30ML			50.00	
	FENLONG LINIMENT-30ml	54.00				
DICLOFENAC SODIUM + METAXALONE	FLEXURA - D	Liniment			115.00	

	DICLOFENAC+PARACETAMOL+ACETAMINOPHEN	CLOFENAC-P-TABLET			4.00	nausea, upset stomach, stomach pain, and dizziness.
4	ASPIRIN	ECOSPRIN - 150 MG	Tablets	Adult dose : 75- 150 mg ; 300 - 900 mg, 4- 6 hours	7.00	Prolonged bleeding time, GI distrubances,
		ECOSPRIN - 75 MG			4.26	
		ECOSPRIN AV - 75 MG			34.30	
		DISPRIN-TABLET			4.70	
5	IBUPROFEN + PARA	COMBIFLAM	Tablets, Syrup	Adult : 200 - 600 mg	11.80	GI Irritation, Nausea, Diarrhoea
		COMBIFLAM			21.90	
	IBUPROFEN	BRUFEN - 200 MG			4.30	
		BRUFEN - 400 MG			7.80	
	IBUGESIC - 100 ML	22.50				
6	PIROXICAM	DOLONEX - 20 MG - 1 ML	Tablets, Injection	Adult: 10 -30 mg	18.00	Upset stomach, Constipation,
		DOLONEX - DT - 20 MG			79.66	
7	ETORICOXIB	ETOCOX - 90 MG	Tablet	Adult: 60-90 mg	66.30	Constipation, Nausea, Vomiting
		ETOSHINE - 60MG			76.00	
		ETORIFLEX-MR-4mgTABLET			165.00	
		ETORIFLEX-MR-8mgTABLET			209.00	
	ETORICOXIB + THICOLCHICOSIDE	EZACT-MR		Adult: 60- 4 mg	169.00	Constipation, Nausea, Vomiting
		NUCOXIA-MR			175.00	
		ETOSHINE - MR			169.00	
8	ETODOLAC	ETOVA - 400MG	Tablet	600-1,000 mg/day	58.50	Constipation, Nausea, Vomiting
		ETOVA - P - 400 MG			78.65	
	ETODOLAC + THICOLCHICOSIDE	ETOVA-MR			157.00	
9	FEBUXOSTAT	FEBUSTAT - 40MG	Tablet	40 - 80 mg OD	102.00	GI Distrubances, Arthritis, Anemia, Hepatitis, Heptomegaly Eczema
10	FLUPIRITINE MALEATE	KATADOL - 100 MG	Capsule	1cap/day	78.50	Constipation, Drownness.
11	FLURBIPROFEN SODIUM	FLUR DROPS - 10 ML	Drops	Instill 1 drop into the eye 4 times/day for 1-3 wk starting at 24 hr after surgery.	111.00	irritation,itching,redness of eyes
12	INDOMETACIN	MICROCID - 25 MG	Capsule	Adult : 25mg bid ro tid - 200mg/day	33.00	CHF, Tachycardia, Chest pain, Arrhythmia, Palpitations, HTN, Pulmanary HTN, Odema, GI disrubances
		MICROCID-SR - 75MG			85.00	
13	KETOROLAC	KETANOV	Tablets , Injection	Tablet 20- 40 mg/day 10-30 mg 4-6 hrly	24.00	Drowsiness, Dizziness, Headache, Mental and
		KETOROL - 1 ML			20.00	

		KETOROL-10MG		IM or IV	40.30	Sensory changes, Psychotic reaction, Sweeting, Dry mouth, Thirst, Fever, Convulsions, Myalgia, Aseptic meningitis, HTN, Dyspnoea, Pulmonary odema, Bradycardia, Chest pain, Palpitations, Fluid retention
14	LEFLUNOMIDE	SUPRIDOL - 1 ML	Tablet	100 mg once daily for 3 days	123.38	HTN, Diarrhoea, Upper respiratory tract infection, Thrombocytopenia, Leukopenia, Pancytopenia, Agranulocytosis, Headache, Alopecia, Dry skin, Eczema, Erythema multiforme
		LEFRA - 10MG			101.81	
15	LORNOXICAM	LORSAID SD - 8 MG	Tablet	Adult: 8-16 mg/day	82.00	Abdominal pain, Diarrhea, Indigestion, Headache, Drowsiness, Visual disturbance, Ringing in ear, High blood pressure, palpitations, Rashes and kidney damage
16	MEFENAMIC ACID	MEFTAL - P	Tablet, suspension	Adult: PO 500 mg tid.	31.00	CHF, HTN, tachycardia,
		MEFTAL - P			21.00	Syncope, Arrhythmia,
	MEFENAMIC ACID +PARA	MEFTAL SPAS			31.00	Vasculitis, Hypotension, Palpitations, Abdominal pain, Vomiting, Dyspepsia, Constipation, Diarrhoea, Nausea, Heartburn
17	MELOXICAM	MUVERA - 15MG	Tablet	7.5 - 15 mg OD	65.00	Dyspepsia, Headache, Nausea, Diarrhoea, upper resp tract infection, Abdominal pain, Dizziness, Oedema, Flatulence, Influenza-like symptoms, Musculoskeletal and Connective tissue signs
18	NAPROXEN	NAPROSYN-250MG	Tablet	0.5-1 g/day in 1-2 divided doses	20.50	GI disturbances, arthritis, Anemia, Hepatitis, Heptomegaly eczema
19	TRAMADOL	SUPRIDOL - 1 ML	Injection, Capsule	PO 50-100 mg 4-6	13.20	GI disturbances, arthritis, Anemia,

		DOMADOL	Tablet	hrly,	69.89	Hepatitis, Heptomegaly eczema
	TRAMADOL + DICLOFENAC SODIUM	DURAPAIN		IV/IM 100 mg, then 50 mg every 1 10-20 mins if needed, up to 250 mg for the 1st hour	119.00	
20	SERRATIOPEPTIDASE	LYSER FORTE	Tablet	Adult: PO 5-10 mg 3 times/day	77.00	Skin rash, diarrhoea, anorexia, GI disturbances, epistaxis (rare).
		BIOSUGANRIL			112.00	
21	TAPENTADOL	TYDOL - 50MG	Tablet	Adult: PO 5-10 mg 3 times/day	118.00	Bladder pain bloody or cloudy urine, body aches , chills
22	TRYPsin	CHYMORAL FORTE	Tablet	2-3tab/ 3times daily	265.00	Tightness in the chest, Swelling of the mouth, Difficulty breathing, Temporary stinging sensation, Burning sensation.
		RUTOHEAL			154.00	
23	TRYPsin,CHYMOTRYPsin and SERRATIOPEPTIDASE	CHYMOCIP-TABLET	Tablet	2-3tab/ 3times daily	274.00	Tightness in the chest, Swelling of the mouth, Difficulty breathing, Temporary stinging sensation, Burning sensation.
24	SINUS INFLAMMATION	FINEHEAL-TABLET	Tablet	1-2 tabs with meals without chewing	17.13	Tightness in the chest, Swelling of the mouth, Difficulty breathing, Temporary stinging sensation, Burning sensation.
25	DICLOFENAC,LINSEED OIL,METHYL SALICYLATE,MENTHOL	LIMUNIGEL-30gm	gel	100-200mg in divided dose	85.00	Nausea, vomiting, dry mouth, abdominal cramps, unpleasant taste, diarrhoea, constipation.
26	NEPAFENAC (0.1% W/V)	NEPAFLAM-5ml EYE DROPS	Drops	1drop, 0.3% should be applied to the affected eye one-time- daily beginning 1 day prior to cataract surgery, continued on the day of surgery and through the first 2 weeks of the postoperative period. An additional drop should be administered 30 to 120 minutes prior to surgery.	137.00	Application site irritation.
27	ADEMETIONINE / S- ADENOSYL METHIONINE	NUSAM-200mg TABLET	Tablet	1tab OD/day	285.00	Headache, Feeling anxious or nervous, Vomiting, Abdominal pain, Diarrhea, Insomnia, Increased sweating, Nausea, Dizziness.
		NUSAM-400mg TABLET		Adults: 1-2 tablets/day	450.00	
28	CAPSAICIN, DICLOFENAC, MENTHOL, METHYL SALICYLATE	SUPAMOVE-30gm CREAM	gel	2 times a day		Nausea, Vomiting, Dizziness, Dry mouth, Drowsiness, Weakness.

29	PIROXICAM (20MG)	UGESIC-20mg TABLET	Tablet	20 mg/day as a single or in divided doses as necessary	106.50	Nausea, vomiting, constipation, lightheadedness, dizziness, or drowsiness.
30	MECOBALAMIN,ALA,INOSITOL,FOLIC ACID,CHROMIMUM,SELENIUM,ZINC	VITASURGE-TABLET	Tablet	1-2 tab/day	159.00	Sleepiness, Colic pain, Injection site related side effects, Pain in abdomen, Abdomen pain, Erosion, Headache, Nausea, Allergic reactions, Vomiting.

SEDATIVES AND HYPNOTICS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ALPRAZOLAM	ALPRAX - 0.25 MG	Tablet	Adult: 0.25 - 0.5 mg BID	16.50	Anorexia, loss of memory & coordination, tachycardia, hypotension
		ALPRAX - 0.5 MG			35.00	
2	CLONAZEPAM	PETRIL MD - 0.5 MG	Tablet	Adult: 1.5 mg/day Paed: 0.01 - 0.03 mg/kg/day	35.00	Anorexia, loss of memory & coordination, tachycardia, hypotension
3	CHLORDIAZEPOXIDE	LIBRIUM - 10 MG	Tablet	Adult:20 - 40 mg OD	45.00	Anorexia, loss of memory & coordination, tachycardia, hypotension
		LIBRAX		Paed: 5-10 mg OD	54.15	
4	CLOBAZAM	FRISIUM 10MG	Tablet	Adult: 20-30 mg/day, up to 60 mg/day Paed: 3-12 yr: 125 mcg/kg bid increased every 5 days. Usual maintenance dose: 250 mcg/kg bid. Max: 500 mcg/kg bid.	139.67	Anorexia, loss of memory & coordination, tachycardia, hypotension
		FRISIUM 5MG			70.16	
5	DIAZEPAM	VALIUM - 5 MG	Tablet	Adult: 2-20 mg BID to QID	15.23	Anorexia, loss of memory & coordination, tachycardia, hypotension
6	LORAZEPAM	LARPOSE - 2 MG	Tablet, injection	Adult: 2-3 mg BD to TID 4 mg IV over 2-5 min Paed: 0.1 mg/ kg IV over 2-5 min	23.50	Anorexia, loss of memory & coordination, tachycardia, hypotension
		LOPEZ - 2ML			17.12	
		NEULORA(LORAZEPAM)-2MG			15.50	
		ATIVAN - 1MG			82.50	
7	PHENOBARBITONE	GARDENAL - 30 MG	Tablet, Syrup	Adult: 30-60 mg OD- TDS	46.62	Sedation, HS reaction, dependence
		GARDENAL - 100ML			46.20	
8	ZOLPIDEM	NITREST - 5 MG	Tablet	Adult: 5-20 mgOD	59.00	Anorexia, loss of memory & coordination, tachycardia, hypotension

9	DOXYLAMINE SUCCINATE, PYRIDOXINE HYDROCHLORIDE	DOXINATE	Tablet	25 mg every 4-6 hr. Max: 150 mg daily.	4.32	Acute dystonic reactions and long-lasting impaired consciousness in child. CNS depression including slight drowsiness to deep sleep, lassitude, dizziness, incoordination. Headache,
10	TRICLOFOS (500MG/5ML)	PEDICLORYL-30ml SYRUP	Syrup	Adult: 1 - 2 g at night Children: up to 1 year: 25 - 30 mg / kg 1 - 5 year: 0.25 - 0.5 g 6 - 12 year: 0.5 - 1 g	75.00	dizziness, irritability, and vomiting.
11	PHENOBARBITONE	PHENOBARBITONE-1ml INJECTION*	Injection	100-200 mg 60-90 min	19.45	Injection site reactions (pain, swelling, redness), Drowsiness, Skin rash, Nausea, Vomiting, Vertigo, Diarrhoea.
12	CHLORDIAZEPOXIDE+MEBE VERINE	COLOSPA-X-5mg TABLET	Tablet	Adult: 5 mg or 10 mg, 3 or 4 times daily;Geriatric Patients: 5 mg, 2 to 4 times daily; pediatric patients: 5 mg, 2 to 4 times daily (may be increased in some pediatric patients to 10 mg, 2 to 3 times daily)	208.50	Sleepiness, Drowsiness, Fatigue, Confusion, Uncoordinated body movement, Abnormality of voluntary movements, Skin rash, Slurred speech.

VACCINE

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	INACTIVATED INFLUENZA VACCINE	INFLUGEN	Injection	0.5ML	720.00	Aches pain, head ache, fever
2	PNEUMOCOCCAL POLYSACCHARIDE VACCINE	PNEUMOVAX 23 - 0.5ML	Injection	0.5 ML	1365.00	Aches pain, Increased temperature Tenderness
3	TETANUS TOXOID	T.T - 0.5 ML	Injection	0.5 ML	10.00	Aches pain, Increased temperature Tenderness
4	TETANUS IMMUNOGLOBULIN	TETGLOB-250MG	Injection	Prophylaxis: 250 Units IM (single dose) Active tetanus: 3000-6000 Units	1200.00	Aches pain, Increased temperature Tenderness
		TETGLOB-500MG			1474.90	

MISCELLANEOUS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ACAMPROSTATE	ACAMPROL - 333 MG	Tablet	PO ≥60 kg: 666 mg 3 times/ day; <60 kg: 666 mg at	57.00	GI disturbances (diarrhoea, nausea, vomiting, abdominal pain, asthenia),

				breakfast, 333 mg at midday and 333 mg at night. Usual duration: 1 yr.		dermatological (pruritus, rash).
2	ADAPALANE	ADIFF-15GM DERIVA CMS - 15 GM	Gel	Adult: Apply thinly onto affected areas once daily at night after cleansing.	130.00	Burning , Itching, Redness
3	ALOE VERA GEL, CALAMINE ,LT LIQ PARAFFIN	CALAPURE-A-50ML CALOSOFT-AF--50 ML	Lotion	apply 1-4 times/day	85.00 89.00	Allergic skin reaction
4	ALOE XTRACT,VITAMIN E ACETATE	ELOVERA - 75GM AHAGLOW-50gm FACE WASH	Cream Gel	Apply regularly to the affected areas as reqd, specially recommended for use after thrice daily	160.00 225.00	Itching, redness
5	ALPHA-CALCIDOL	ALPHA-D3 - 0.25MCG	Tablet	Initial: 1 mcg/day. Maintenance: 0.25-1 mcg/day.	82.40	Rash (or) Hives, Itching
6	ALUMINIUM CHLOROHYDRATE	ALDRY - 100ML ALDRY-150gm LOTION	Lotion	twice/thrice daily or sos twice/thrice daily or sos	175.00 300.00	Itching, redness
7	AMINO ACIDS & VITAMINS CAPSULES	OVABLESS ASTYMIN FORTE ASTYMIN - 3 - 200 ML I.V XTRAGLO ASTYMIN-3-20ml INJECTION KETOADD-TABLET	Tablet, Capsules, Injection	1-2 cap/day 1-2 cap/day adult:200-800mL iv once/daily;children:0.2-0.25 grams of nitrogen per kg body weight; drip drate 15-20 drops initially followed by 20-40 drops per min 5mL/daily adult:200-800mL iv once/daily;children:0.2-0.25 grams of nitrogen per kg body weight; drip drate 15-20 drops initially followed by 20-40 drops per min 200mg bd or tds/day	12.00 96.00 416.50 100.00 98.65 299.00	Nausia, Pain, Head ache
8	ANTI- D IMMUNO GLOBULIN(MONOCLONAL)	RHOCLONE - 300MCG (ANTI-D)	Injection	300 mcg	2466.90	Pain, Discomfort
9	AZATHIOPRINE TABLETS	AZORAN - 50MG	Tablets	1-3 mg/kg/day.	114.50	Fever, chills; bone marrow characterised by leucopenia, depression thrombocytopenia or anaemia; anorexia, nausea, diarrhoea; arthralgias; secondary infections
10	AZELAIC ACID	AZIDERM-10% - 15GM AZAC SOAP - 75 GM	Cream, Soap	Apply thinly into the affected areas bid (morning and evening) after cleansing for 6months	159.00 70.00	Itching, redness

11	BARIUM SULPHATE ORAL SUSPENSION 1 LIT	BARIUM SULPHATE SUSPENSION 1 LIT ESKAY	Suspension	5-10 g dissolved in 250 mL of water	260.00	Bloating, Constipation
12	B-COMP FORTE VIT-C	BECOSULES	Capsules	1tab/BD	25.80	Itching, redness
		BECOSULES - Z			25.96	
13	B12-DEFICIENCY	POLYBION-CZS-TABLET	Tablet	1tab/BD	63.00	Nausea, vomiting, nail changes, loss of energy, and irritability.
14	BENFOTIAMINE 100 MG	BENALGIS - 100 MG TABLET	Tablet	1tab/ tid	65.85	Nausia, Vomiting, GI Irritation
15	BENZALKONIUM & ZINC	HAPPY NAP - 15GM CREAM	Cream	Apply twice or thrice daily	65.00	Itching, redness
16	BENZOYL PEROXIDE 5%	BENZAC AC - 20 GM GEL	Gel	Apply once daily or twice to cover affected areas after washing	125.95	Itching, redness
17	BETAHISTINE	VERTIN - 16 MG	Tablet	Adult: 8-16 mg TDS	109.00	GI upset, dizziness, insomnia
		VERTIN - 8 MG			58.00	
18	BIOTIN, AMINOACIDS, VITAMINS, MINERALS, NATURALEXTRACTS	FOLLIHAIR TABLET		1-2cap/day	155.00	Nausia, Vomiting, Head ache
19	BIOTIN 10 MCG+DL-METHION	MACTOTAL-200ml SUSPENSION	Suspension	1tab/bd	106.00	Constipation, diarrhea, or upset stomach.
20	BISACODYL	DULCOFLEX	Tablet, Suppository	5-10 mg/day at night po As supp/enema: 10 mg/day in the morning.	11.21	Abdominal discomfort (colic, cramps). Suppositories may cause irritation and proctitis.
		DULCOFLEX - ADULT			19.05	
21	BRIMONIDINE TARTARATE, TIMOLOL MALEATE	BRITIBLU-5ML DROP	Eye drops	Instill 1 drop bid or tid (approx 8 or 12 hrly).	108.50	Eye irritation, Burning sensation
22	CALAMINE	LINICAL- 100ML LOTION	Lotion	Topical Apply 1-4 times/day.	110.00	Itching, redness
		VENUSIA SOFT-75ML LOTION			275.00	
23	CALCITONIN	OSTOSPRAY	Spray	200 -400 IU daily	1020.00	Nasal Irritation
24	CALCITRIOL	ROCALTROL - 0.25mcg	Capsule	0.015-0.02 mcg/kg/day. Maintenance: 0.03-0.06 mcg/kg/day.	184.78	Weakness; headache; somnolence; nausea; vomiting; dry mouth; constipation; muscle pain; bone pain; metallic taste; polyuria; polydipsia; anorexia
25	CALCIUM CARBONATE	OSTONATE	Syrup, Capsule, Tablet,	10mcg	82.50	Constipation, bloating, flatulence
		SHELCAL - 500 MG	Powder, Suspension	Initial: 1 mcg/day. Maintenance: 0.25-1 mcg/day	62.00	
26	CALICIUM CITRATE	CALCINOS		1200 mg/day.	88.50	
		ROCKBON			160.00	
		PHOSTAT - 667MG			34.15	
		CCM			194.70	
		SHELCAL CT			178.00	
		CALDIKIND - 200ML			100.00	

27	CALCIUM & VITAMIN D	JUSDEE 400-30ml DROPS	Drops	10mg/day	74.26	Nausea/vomiting, constipation, loss of appetite, increased thirst, increased urination, mental/mood changes, unusual tiredness.
28	CALCITRIOL +CALCIUM CITRATE	BON-K2 HD-TABLET		Initial: 0.25 mcg/day	207.00	Constipation, wind, feeling sick, stomach ache, diarrhoea.
29	DENOSUMAB	XGEVA-TM-120mg/1.7ml-INJECTION(DENOSUMAB)		120 mg single SC inj once every 4 wk	28787	Tiredness, weakness, headache, back pain, joint pain, diarrhea, or nausea.
30	POTASSIUM BINDER	K-BIND - 15GM		used once/daily	112.00	Vomiting, Stomach irritation, Nausea, Constipation, Loss of appetite.
31	CALCIUM DOBESILATE	OSIL	Capsule, Injection,	Adult: PO 0.5-1 g/day in divided doses.	130.00	Skin rash, fever, vertigo
		CALCIUM GULCONATE - 10 ML	Suspension		20.00	
		CALCIMAX-P - 200ML			99.00	
32	ELEMENTAL CALCIUM	MAXICAL-500mg TABLET	Tablet	1000mg/day	71.50	Nausea/vomiting, loss of appetite, unusual weight loss, mental/mood changes, bone/muscle pain, headache, increased thirst/urination, weakness, unusual tiredness.
		CALCIMAX FORTE-TABLET	Tablet		192.00	
33	CARBONYL IRON WITH ZINC AND FOLIC ACID	FEFOL-Z - CAPSULE	Capsule	1 cap od	78.70	GI irritation, Nausea, Vomiting
34	CARBOXY METHYL CELLULOSE	JUST TEARS - 10 ML EYE	Eye drops	1-4 applications/day	81.00	Eye irritation, Burning sensation
		LACRIMOS 0.5% - 10ML			79.00	
		LUBREX - 10ML EYE			89.89	
		AQUALUBE - 10ML EYE			153.00	
35	CHLORHEXIDINE	HEXIDINE - 160ML	Mouth wash	Rinse w/ 10 mL for about 1 min bid. As spray: Apply bid. Max: 12 actuations bid. As gel: Brush teeth w/ 1 inch of gel once or	69.00	Dry mouth, Mouth irritation
		REXIDIN - 120ML			72.00	
		CLOHEX PLUS - 150 ML			86.50	
36	CHOLECALCIFEROL	DVION PEARL - 60K	Capsule	10 mcg/day	109.75	Hyperphosphataemia or hypercalcaemia (in excessive intake). Associated effects of hypercalcaemia include hypercalciuria, ectopic calcification,& renal &CVdamage.
37	CHOLINE SALICYLATE	ZYTEE - 10 ML GEL	Gel	Apply gel on affected area once/day	70.00	Dry mouth, Mouth irritation
		DOLO-CT GEL - 10GM			49.00	
38	CLINDINUM BROMIDE, CHLORDIAZEPOXIDE & DICYCLOMINE HYDROCHLORIDE	NARMOXIN	Tablet	PO Per cap contains chlordiazepoxide 5 mg and elidinium 2.5 mg: 1-2 cap 3-4 times/day	18.00	mouth dryness, blurring, hesitancy and urgency in micturition, constipation, rise in body temperature, heat stroke, mydriasis, tachycardia, headache, nervousness

39	CLOBETASOL PROPIONATE & FUSIDIC ACID CREAM	CLONATE - F - CREAM	Cream	Apply thinly onto the affected area once daily or bid, reduced if necessary. Max dose: 50 g wkly. Max duration of therapy: 4 wk.	145.00	Irritation, Itching, Burning
	CLOBETASOL PROPIONATE & SALISYLIC ACID	TOPISOL 3% LOTION - 30ML			74.00	
40	CLOBETASOL PROPIONATE	TENOVATE - 15 GM	Ointment	Apply thinly onto the affected area once daily or bid, reduced if necessary. Max dose: 50 g wkly. Max duration of therapy: 4 wk.	56.30	Irritation, Itching, Burning
41	COAL TAR & SALICYLIC ACID & BENZALKONIUM CHLORIDE	IONEX-T - 60 ML LOTION	Lotion	Massage the solution into wet hair. Rinse, then apply again and massage in hair for about 5 mins before rinsing. May be used daily	193.65	Itching, redness
42	COLLAGEN PARTICLES	BIO-SEAL - 5 ML	Powder	make a paste/solution of BIO-SEAL in sterile saline/DM water and use with the help of a syringe. Cover the wound with any conventional absorbent dressing of choice	320.00	Itching, redness
43	COLLOIDAL IRON, LYSINE, FOLIC ACID, VIT B12	TONOFERON - 15 ML DROPS	Drops, Syrup	Infants less than 1 yr: 5 drops b.i.d. Children above 1 yr: 10 drops b.i.d.	63.00	GI Irritation, Vomiting, Nausea
		TONOFERON SYRUP - (PAED)			78.00	
44	COROVERINE	TINNICAR	Tablet	20-40 mg 3-4 times daily. Max: 200 mg/day.	185.00	
45	CRANBERRY & D-MANNOSE	CRANPAC- D	Tablet	Stir evenly & drink 1 sachet diluted in 100 mL water every morning for 10 days	210.00	
46	DESONIDE	DESOWEN - 10GM	Cream	Apply as thin film to the affected area bid/tid depending on the severity of the condition.	126.99	Itching, redness
47	DIACEREIN	VOKMOR -50MG	Tablet , Capsules	50 mg bid.	49.00	Diarrhoea, epigastric pain, nausea, vomiting, intense yellow colouring of urine.
48	DIACETYL BOLDINE, JOJOBA OIL, ASCORBYL PALMITATE	AVARTA-10GM UNDER EYE CREA	Eye Cream	apply night under eye once /day	490.00	

49	DIATRIZOATE MEGLUMINE	UROGRAFFIN - 76 %	Injection	Radiographic Examination of Segments of the Gastrointestinal Tract Oral Administration: Adult oral dosage may range from 30 to 90 mL (11 to 33 g iodine), depending on the nature of the examination and the size of the patient. For infants and	206.00	Stomach cramps, Rash, Heart burn
	& DIATRIZOATE SODIUM			children less than 5 years of age, 30 mL (11 g iodine) are usually adequate; for children 5 to 10 years of age, the suggested dose is 60 mL (22 g iodine). These pediatric doses may be diluted 1:1, if desired, with water, carbonated beverage, milk, or mineral oil. When used in infants, the solution may be		
50	DIATRIZOATE SODIUM	GASTROPAQUE - 30ML	Oral Contrast	Oral Administration: Adult oral dosage range 30 to 90 mL (11 to 33 g iodine), For infants & children < 5 years of age, 30 mL (11 g iodine); for children 5 to 10 years of age, the suggested dose is 60 mL (22 g iodine). These pediatric doses may be diluted 1:1, if desired, with water, carbonated beverage, milk, or mineral oil. When used in infants, the solution may be given in a nursing bottle. Pediatric doses may also be used	310.00	
51	DIFLUPREDNATE	DIFLUCOR DROPS-5ML	Drops	Instill one drop into the conjunctival sac of the affected eye(s) 4 times daily beginning 24 hours after surgery and continuing throughout the	104.00	Skin thinning when applied under occlusion. Folliculitis, hypertrichosis, skin atrophy, superinfection, stretch mark, loss of skin collagen and SC atrophy,
		UVEFROST-5ML EYE DROP			102.00	
52	ESCIN ASCORBIC ACID & PROANTHOCYANIDIN	ELIP	Tablet	one tablet daily	148.50	GI Irritation, Vomiting, Nausea
53	ESSENTIAL AMINO ACIDS AND VITAMINS	ASTYMIN - 200 ML	SYRUP	Single Dose : 15 ml	152.50	Nausia, Vomiting, Haed ache

54	FERRIC PREPARATIONS	HEMSI-PD - 15ML	Drops & Injection	IRON DEFICIENCY IN INFANTS AND TODDLERS : Breastfed preterm infants - elemental iron of 2 mg/kg per day from 1 month of age through 12 months ,Formula-fed preterm infants, 2 mg/kg per day of iron from a standard infant formula	37.30	Constipation
		FERIUM - 500MG		ferium dose : 5 ml bid	1845.00	
		DEXORANGE-200ml SYRUP	Syrup	dexorange syrup -Adult: 15-30ml; child: 2-5 yrs: 5ml; child 5-12 yrs: 10ml. All bid after meals.	109.25	Nausea, Vomiting, Indigestion, Diarrhoea, Bitter taste in mouth, Dark coloured stool.
		DEXORANGE-CAPSULE	Capsule	dexorange capsule -1 cap bid after meals.	106.56	
55	FERROUS PREPARATION	HEMFER-XT	Tablet , Capsules, Injections, Syrup	Hemifer - XT tab :1 tab daily.	77.00	Constipation, GI Irritation, Upset Stomach
		LIVOGEN-XT		Livogen XT Adult Dose for Megaloblastic Anemia :1 mg orally, intramuscularly, subcutaneously or IV once a day. Adult Dose for Folic acid (Livogen-XT) Deficiency :400 to 800 mcg orally, intramuscularly, subcutaneously or IV once a day. Women of childbearing age,	99.50	
		FERIUM-XT		Ferium XT :1 tab daily.	102.40	
		LIVOGEN		Livogen capsule : pregnancy anemia :1 cap bid-tid. As a supplement & maint dose: 1 cap/day.	28.90	
		LIVOGEN-250ml SYRUP		Livogen syrup :Adult: PO – The recommended dose is 10-25 mg/day in single or divided doses.	129.00	
		AUTRIN		Autrin :1 cap daily	2.82	
		LIVOGEN - Z		Livogen - Z :1 cap daily	31.00	
		FEFOL		Fefol: 1 cap od	67.00	
		IMFERON - 2ML		Imferon :Adult: IM- 250-1,000 mcg on alternate days for 1-2 week, then 250 mcg weekly. Maintenance: 1,000-mcg monthly.	30.62	

		REFIL	Refil :1 – 2 capsules a day	66.00		
		JECTOCOS - 1.5 ML	Jectocos :Adult: IM- 250-1,000 mcg on alternate days for 1-2 week, then 250 mcg weekly. Maintenance: 1,000-mcg monthly.	34.26		
		FERIKIND-S -100MG - 5ML	Ferrikind S :5 ml bid or as reqd.	242.00		
		FERRI	Ferri :Adult 1-2 amp/time (100 mg/5 mL amp) 2-3 times wkly. Max: 7 mg/kg body wt as Fe (0.35 mL inj) infuse over 3½ hr. Childn 3 mg/kg body wt as Fe (0.15 mL inj).	30.00		
		IV CROSS - 100 MG - 5ML		336.00		
		ZINCOFER	Zincofer tablet :PO Usual range: Up to 600 mg/day, up to 1.2 g/day if needed.	31.90		
		ZINCOLIFE-50ml SYRUP	Diarrhoea – Children aged < 6 month: 2.5 ml once a day for 14 days. Children aged 6 months and above): 5 ml once a day for 14 days.	51.23	Diarrhea, Abdominal cramping, Vomit, Loss of appetite, Stomach cramps, Headache.	
		ZINCOVIT SYRUP-200ml	Zincovit syrup :Adult dose : 5ml.	106.00	nausea, vomiting, diarrhea, metallic taste, kidney and stomach damage	
		ZINCOVIT TABLET	Zincovit tablet :1 tablet daily	83.00		
		ZINCOVIT-15ml DROPS**		50.00		
		Z&D-DS-20-15ml DRY SYRUP	Diarrhoea – Children aged < 6 month: 2.5 ml once a day for 14 days. Children aged 6 months and above): 5 ml once a day for 14 days.	67.00	Convulsions, Vomiting, Loss of appetite, Stomach cramps, Diarrhea, Headache.	
		ALOHA XT - 150 ML	1 tab daily.	140.00	Allergic reactions, Anorexia, Nausea, Abdominal distention, Flatulence, Bitter or bad taste, Altered sleep patterns, Difficulty in concentrating, Irritability, Overactivity.	
		C-PINK-150ml SUSPENSION	Suspension	Elemental iron 30 MG + Folic acid 500 MCG /5ML	133.00	Diarrhea, nausea, vomiting, abdominal cramps/pain, or heartburn
		CPINK-15ml DROPS	Drops	Elemental iron 10 MG + Folic acid 100 MCG /5ML	74.70	

		ASCAZIN-TABLET	Tablet	Adults: 25 to 50mg zinc daily or 1 to 2 tablets daily. Children: 5mg/kg 1 to 3 times daily.	54.00	nausea, vomiting, diarrhea, metallic taste, kidney and stomach damage.
		LIV-52 TABLET	Tablet	Drug-induced hepatitis: 1 tablets b.i.d. with anti TB/statin/chemotherapy/antiretroviral treatment. Jaundice and anorexia during	95.00	Burning of skin, Skin irritation, Irritation of skin.
		LIV-52-100ml SYRUP	Syrup	Acute viral hepatitis: children 5 ml tds, adult 10 ml tds/Alcohol-induced liver damage: Syrup: 10 to 15 ml t.i.d./Anorexia and suboptimal growth: children 5 ml tds/bd, adult 10 ml tds/b.d. Drug-induced hepatitis: 1 to 2 teaspoonfuls/b.i.d. Jaundice and anorexia during pregnancy: 1 teaspoonful b.i.d. for 8 weeks. Chronic active hepatitis: 2 teaspoonfuls b.i.d. for 6 months Early cirrhotic conditions: 1 teaspoonful t.i.d. for 6 to 12 months	65.00	
		APTIVATE-175ml SYRUP	Syrup	Children under age 6 years are given a dosage of 12mg daily on spilted basis and children above 6 years of age are given a dosage of 16mg each day.	84.50	Nausea, Stomach Upset, Skin Rash, Acute Toxicity.
56	DIASTASE+PEPSIN	ARISTOZYME-SYRUP	Syrup	take 5 ml of Diastase + Pepsin, immediately after meals (for adults)	81.00	Irritation in Stomach, Upset stomach, Irritation in throat, Damage in throat, Blisters, Allergic reactions.
57	FLAVOXATE HCL	URISPAS - 200 MG	Tablet	200 mg tid.	234.91	Increased intraocular pressure; difficulty in ocular accommodation; blurred vision; dry mouth & throat GI disturbances; nervousness; vertigo headache

58	FLUOROMETHOLONE	FML EYE DROPS - 5ML	Eye drops	TO THE EYE ► Child 2–17 years: Apply every 1 hour for 24–48 hours, then reduced to 2–4 times a day ► Adult: Apply every 1 hour for 24–48 hours, then reduced to 2–4 times a day	87.17	
59	FRAMYCETIN SULPHATE	SOFRAMYCIN	Cream	Adult Dose Antibacterial (ophthalmic) Topically, to the conjunctiva, a thin strip of ointment to the affected eye 2 or 3 times daily, or at bedtime if the drops have been used during the day.	31.29	Itching, redness
60	FUCIDIC ACID & BETA METHASONE VALERATE	FUCIBET - 15 GM	Cream	Adult : Topical Apply to the affected area(s) bid. Max duration: 2 wk	43.70	Itching, redness
61	FOLIC ACID, METHYLCOBALAMIN, VITAMIN A, VITAMIN B1, VITAMIN B2, VITAMIN B3, VITAMIN B5, VITAMIN B6, VITAMIN C, VITAMIN E	A TO Z NS-200ml SYRUP	Syrup	For adults: 3 - 4 teaspoonfuls daily; For children of 4 - 12 years: 2 - 3 teaspoonfuls daily;	121.00	Constipation, diarrhea, or upset stomach
		A TO Z-15ml DROPS	Drops		32.71	Feeling tired or weak (fatigue), stomach pain, loose stools (diarrhea), headache, nausea, or blurred vision.
62	FOLIC ACID(VITAMIN B9), VITAMIN B12, VITAMIN D3, VITAMIN B1, NIACINAMIDE, VITAMIN B2, VITAMIN B6, CALCIUM PANTOTHENATE, VITAMIN A, VITAMIN C, VITAMIN E, ZINC	A TO Z-TABLET	Tablet	once daily	79.45	Tooth staining, increased urination, stomach bleeding, uneven heart rate, confusion, and muscle weakness or limp feeling.
63	GLUCOSAMINE & MSM & DIACEREIN	JOINTACE-DN	Tablet	500 mg 3-4 times daily	233.00	Heart burn, epigastric pain/tenderness, diarrhea, nausea, dyspepsia, constipation, abdominal pain, palpitations, drowsiness, skin reaction, headache, indigestion.
		TOVOK			150.50	
		REJOINT			173.00	
		CARTISAFE - D			142.50	
64	GLYCOLIC ACID 12%	GLYCO - A - 30 GM	Cream	The first night leave the Glycolic Cream® on for 10 minutes then wash off in cool	208.00	Itching, redness
		GLYCO-6% - 30GM			130.00	
65	HALOBETASOL PR0PIONATE	HALOVATE - 15GM	Lotion, Ointment	Adult Dose/ 12 years older .Apply a thin layer to affected area once or twice a dayTreatment should be limited to two weeks.	149.00	Itching, redness

		HALOVATE - 30ML			169.50	
		HALOVATE-F - 10GM			130.50	
		HALOVATE-S - 15GM			179.00	
		HALOX-S - 30ML			189.00	
		ULTRAVEX - S3			127.00	
		ULTRAVEX - S6			169.50	
66	HUMAN PLASMA PROTEIN	HUMAN ALBUMIN 5 % -250 ML	Injection	minimum effective dose in adults is 250-500 mL.	6350.00	
67	HUMAN NORMAL IMMUNOGLOBULIN	HISTOGLOB	Injection	Adult: IV Prevention of infections after bone marrow transplantation 500 mg/kg/wk.	91.00	
68	HYALURONIC ACID	HYALGAN PFS 20MG - 2ML	Injection	20 mg	1450.00	
		HYNIDASE			114.24	
69	HYDROCORTISONE ACETATE CREAM 1%W/W	CUTISOFT-10GM	Cream, Injection	0.1-2.5% oint/cream/lotion: Apply as directed.Intra-articular Joint inflammations As acetate: 5-50 mg depending on size of affected joint	71.00	odium and fluid retention. Potassium and calcium depletion. Muscle wasting, weakness, osteoporosis. GI disturbances and bleeding.
		CORT - S - 100 MG			41.00	
		CLEARZ MAX - 15GM			185.00	
70	HYDROXYPROPYL	HICOOL - 5 ML EYE DROPS	Eye drops	drops may be used as often as needed.over the eye and squeeze out 1 or 2 drops as needed. Look down and gently close your eye for 1 or 2 minutes,Ointments are usually used 1 to 2 times daily	69.50	
	METHYLCELLULOSE	APPAVISC SOLUTION - 5 ML			60.00	

71	IODIXANOL	VISIPAQUE-320MG -- 100ML	Injection	<p>Intra-arterial administration (arteriography): Carotid arteries: 10-14 mL Vertebral arteries: 10-12 mL Right coronary artery: 3-8 mL Left coronary artery: 3-10 mL Left ventricle: 20-45 mL Renal arteries: 8-18 mL Aortography: 30-70 mL Major aorta branch: 10-70 mL Peripheral arteries: 15-30 mL Aortofermoral runoffs: 20-90 mL</p> <p style="text-align: center;">Intra-arterial administration (IA-DSA)</p> <p style="text-align: center;">Carotid or vertebral arteries: 5-8 mL Aortography: 10-50 mL Major aorta branch: 2-10 mL Aortofemoral runoffs: 6-15 mL Peripheral arteries: 3-15 mL</p> <p>IV administration CECT of head or body: 75-150 mL bolus, then 100-150 mL infusion not to exceed 150 mL Excretory urography: 1 mL/kg,</p>	2750.00	
72	IOHEXOL	OMNIPAQUE - 300--50ML	Injection	<p>Adult: Inj As iohexol soln containing 6-350 mg iodine/ml:</p> <p>Dose and strength used dependson the procedure and route of administration</p>	742.00	Flushing or a sensation of heat; pain, extravasation, thrombophlebitis at the inj site; nausea, vomiting, headache, and dizziness; urticaria, pruritus, pallor, sweating, metallic taste, weakness, coughing, rhinitis, sneezing, lachrymation, visual disturbances
73	ISOTRETINOIN	TUFCANE 10 MG TUFCANE 20 MG	Capsules	<p>Adult: PO Initial: 0.5-1 mg/kg/ day in 2 divided doses</p>	114.30 142.00	Dryness of mucous membranes, dryness of skin with scaling, fragility, erythema, cheilitis, pruritus, epistaxis, conjunctivitis, dry sore mouth and palmo-plantar exfoliation
74	LACTITOL MONOHYDRATE	LACTIHEP - 200ML	Syrup	<p>the usual recommended dosage in adults is about 15-30 ml per day. The medication should be taken with meals. If no adverse reactions show then the dosage</p>	196.00	Abdominaldistension, Flatulence, Nausea, Vomiting, Dyspepsia, Cramps, Epigastric pain, Abdominal discomfort, Anal pruritus, dyspepsia

				maybe increased to 60 ml per day.		
75	LACTULOSE	DUPHALAC - 200 ML LIVO LUK - 60 ML LOOZ ENEMA - 275ML	Solution, Syrup, Enema	Adult: PO Constipation Initial: 10-20 g (15-30 mL)/day Max: 45 mL (or 40g of the reconstituted oral formulation)/day. Hepatic encephalopathy 60-100 g (90-150 mL)/day in 3 divided doses; adjust accordingly. Rectal Hepaticencephalopathy Mix 200 g (300 mL w/ 700 mL water or 0.9% NaCl as a retention enema. Retain enema for 30-60 mins; repeat 4-6 hrly until oral medication can be administered.	223.50 68.90 235.00	Diarrhoea, Nausia, Vomiting
76	L-AEGININE, CARNITINE, PIPERINE,OMEGA 3 LYCOPENE	K.Q-100	Capsule	once daily	390.00	Nausea, Vomiting
77	L-ARGININE , PROANTHOCYANIDIN	ARG9 - 5GM SACHET ARGIPREG - 6.5GM SACHET	Powder	Mix the contents of the sachet in half a glass of water and one sachet daily	30.00 29.46	Stomach burn
78	MOMETASONE 0.1% W/W MOMETASONE 0.1% W/W FUSIDIC ACID 2.0 W/W MOMETASONE 0.1% W/W TERBINAFINE	MOMATE - 30ML MOMATE -15GM MOMEFLO - NASAL SPRAY METASSPRAY - NASAL SPRAY HH SONE - 10 GM HH FUDIC -10 GM MOMATE-F -10GM TYZA-M - 10GM	Cream, Lotion, Nasal spray	Prophylaxis and treatment of allergic rhinitis ► BY INTRANASAL ADMINISTRATION ► Child 6–11 years: 50 micrograms daily, dose to be sprayed into each nostril ► Child 12–17 years: 100 micrograms daily, increased if necessary up to 200 micrograms daily, dose to be	250.00 171.50 202.50 245.50 116.00 208.00 145.00 63.00	Itching, redness
79	MULTIVITAMIN PREPARATIONS	RICONIA L.P POLYBION LC - 100ML 5ml single d FOURTS - B 1 tab daily AQUASOL - A NEUROBION FORTE LIMCEE - 500 MG JUSDEE - 50ML	Capsule, Syrup, Tablet, Drops, Injections	One capsule daily, POLYBION LC - 100ML 5ml single dose, FOURTS - B 1 tab daily ,Aquasol -A Vitamin A Deficiency Malabsorption or oral administration not feasible: 100,000 U/day IM for 3 days; then 50,000 U/day for 2 weeks; follow with oral therapy.Oral therapy: Take oral therapeutic	67.00 47.50 108.00 24.55 7.22 15.34 72.60	Increased Urination

	JUSDEE 400 - 15 ML		multivitamin containing 10,000-20,000 U/day vitamin A for 2 months	61.71	
	D-RISE--60K		Deficiency prophylaxis: 10,000-50,000 U PO qDay	142.75	
	ARACHITOL- 3 I.U -1 ML		Xerophthalmia (Off-label).Recommended dose except for females of reproductive age: 200,000 units PO qDay for 2 days; repeat dose again after 2 weeks.Females of	27.50	
	FOURTS - B			108.00	
	BEPLEX FORTE-TABLET	Tablet	once daily	29.22	Constipation, diarrhea, or upset stomach.
	BECOZINC-CAPSULE	Capsule	1 cap OD	24.00	Swelling of lips, Swelling of face and tongue, Difficulty in breathing, Diarrhea, Swelling, Rapid weight gain, Itching or mild rash, Nausea, Stomach upset, Heartburn.
	HEALTH OK-15ml DROPS	Drops	once daily	50.00	Constipation, diarrhea, or upset stomach.
	HEALTH OK-TABLET	Tablet	once daily	150.00	Tooth staining, increased urination, stomach bleeding, uneven heart rate, confusion, and muscle weakness or limp feeling.
	MULTI VITAMIN-10ml INJECTION	Injection	daily dosage volume is 10 mL. One daily dose (10 mL) is diluted by adding directly to a specified volume of an intravenous fluid	25.00	Severe allergic reaction (anaphylaxis), abnormally high levels of vitamin A, rash, redness, itching, headache, dizziness, agitation, anxiety, and double vision
	COBADEX CZS-TABLET	Tablet	Adult: 200-1000 mcg daily.	70.35	loss of appetite, diarrhea, vomiting, abdominal cramps, breast soreness, photosensitivity, sleep disturbances, mental changes

80	THIAMINE	THIAMIN-2ml INJECTION	Injection	<p>beriberi :10 mg to 20 mg of thiamine hydrochloride are given IM 3 times daily for as long as 2 weeks.</p> <p>Infantile beriberi :doses of 25 mg may cautiously be given IV.Neuritis of pregnancy :5 mg to 10 mg of thiamine hydrochloride IM daily.Wernicke-Korsakoff syndrome, thiamine hydrochloride has been administered IV in an initial dose of 100 mg, followed by IM doses of 50 mg to 100 mg daily.Patients with marginal thiamine status to whom dextrose is being administered should receive 100 mg thiamine hydrochloride in each of the first few liters of IV fluid to avoid precipitating heart failure.</p>	50.40	Pain/redness/soreness at the injection site
		AMINTOZ-100mg-1ml (THIAMINE) INJECTION			25.20	
81	BENFOTIAMINE 100 MG	AISA-TABLET	Tablet	twice with divided doses. It is taken orally with a dosage between 300-600mg.	144.00	Nausea, Vomiting.
82	BENFOTIAMINE, BIOTIN, CALCIUM ASCORBATE, CHOLINE	BECOSTAR-G-TABLET	Tablet	One capsule to taken once daily after meal	160.00	Constipation, diarrhea, or upset stomach
83	UBIDECARENONE+L-CARNITINE+L-TARTRATE+CHOLINE BITARTRATE+ SELENIUM	ASTHOQUE-TABLET	Tablet	2 tablets a day	59.00	Drowsiness, dizziness, headache, excitement, nausea, vomiting, and diarrhea
84	LIQUID PARAFFIN	LIQUID PARAFFIN - 100 ML	Suspension, Syrup	adults is 10-30ml when required. children over the age of 3 years old is 5-20ml	60.00	No common side effects seen.
		CREMAFFIN - 200 ML			106.37	
		CREMAFFIN PLUS			123.85	
85	LOPERAMIDE HYDROCHLORIDE	LOPERMIDE	Tablet	6-8 mg/day	16.87	Abdominal pain, distention, and discomfort; paralytic ileus; constipation, dry mouth, drowsiness, dizziness, fatigue, rash.
86	L-ORNITHINE &	HEPACOR - 10 ML	Injection & Tablet	Adult: 1 - 2tablets 3 times daily	213.00	Vomiting,Transient nausea.

	L-ASPARTATE	HEPAMERZ - 10ML		each tablet contain 150 mg of L-Ornithine L-aspartate ,Adult	270.00	
		HEPAMERZ		Dosage	122.00	
87	MECOBALAMIN	NUROKIND - 500MG	Injection & Tablet	Oral :Peripheral neuropathies Adult: 1500 mcg/day in 3 divided doses. Parenteral :Peripheral neuropathies Adult: 500 mcg daily IM/IV 3 times/wk. Parenteral :Megaloblastic anaemia caused by vitamin B12	32.40	GI Irritation, Nausea, Vomiting
		NUROKIND			12.00	
		RENERVE			83.50	
	METHYLCOBALAMIN & ALPHA LIPOIC ACID, FOLIC ACID& PYRIDOXINE HCL	117.00				
88	MENADIONE	VITAMIN - K - 1 ML	Injection	IV Vit K Deficiency 0.03 mcg/kg/day. IM Haemorrhage or threatened haemorrhage during severe hypoprothrombinaemia 2.5-10 mg/day.	6.00	Facial flushing, sweating, chest constriction/pain, dyspnoea, cyanosis, CV collapse. IM: Severe shock-like reactions, phlebitis, inj site reactions
89	MEPHENESIN METHYL NICOTINE & CAPSICUM OLEORESIN	RELAXYL - 30GM	Ointment	Apply half inch ointment over affected area gently 3-4 times a day.	50.25	Skin irritation & rash
90	METHYLERGEOMETRINE MALEATE	MEM - 1 ML	Injection	200 mcg 3-4 times/day	14.50	Headache, dizziness, hallucinations; tinnitus; nausea, vomiting, foul taste, diarrhoea; hypertension, temporary chest pain, palpitations, bradycardia
91	MINODOXIL	MINTOP 2% - 60 ML LOTION	Lotion	apply 1 ml to the scalp bd	233.00	Itching, redness
		MINTOP FORTE 5% - LOTION			528.00	
92	MODIFIED LANOLIN	NIP CARE OINTMENT	Ointment		118.00	Itching, redness
93	NAPHAZOLINE & CPM & METHYLCELLULOSE	NEFACOOOL - 10ML EYE DROP	Drops	Instill one or two drops into the conjunctival sac(s) every three to four hours as needed	38.00	Eye irritation
94	OCTINOXATE & ZINC OXIDE	SUNSTOP 30 - 60GM LOTION	Lotion	apply liberally 15 minutes before sun exposure reapply at least every 2 hours use a water resistant sunscreen if swimming or sweating	298.00	Itching, redness
95	OXYMETAZOLINE HCL	NASIVION (ADULT) - 0.05 %	Drops	Adults and children 6 years of age: 2 to 3 drops of in each nostril twice daily, in the morning and evening.	63.00	Nasal irritation
		NASIVION (PAED) - 0.025 %			46.00	
96	PAPAIN-UREA DEPRIDING	DEBRIDACE - 15GM	Ointment	1 or 2 times daily. After cleansing, apply ointment directly to lesion and cover with appropriate dressing.	145.00	Itching, redness

97	PAPAVERINE HYDROCHLORIDE	REPAVERINE - 2ML	Injection	1 to 4 mL	19.80	general discomfort, nausea, abdominal discomfort, anorexia, constipation or diarrhea, skin rash, malaise, vertigo, headache, intensive flushing of the face, perspiration,
98	PARADICHLOROBENZENE BENZOCAINE, CHLORBUTOL	OTOREX - 10 ML EAR DROPS	Drops	2 or 3 drops should be instilled into the ear three or four times daily; alternatively, a gauze wick kept saturated with the drops may be inserted into the external auditory meatus	68.72	
99	PERMETHRIN	PERMITE - 30 MG CREAM	Cream, Soap	Apply Perlice Creme/Rinse 1% Cream to the hair, or skin, depending on the location of infection. Avoid its contact with eyes. The same treatment can be repeated after seven days if required	61.34	Itching, redness
		SCABELICE MEDICATED-75GM			48.40	
		PERLICE - 60GM CREAM			119.15	
100	POLIDOCANOL	ASKLEROL 3% - 2ML	Injection	0.1 to 0.3 mL per injection	111.00	Bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation
101	PYRIDOXINE HCL	BENADON - 40 MG	Injection & Tablet	150 mg per day	13.28	Severe peripheral neuropathies (with long-term admin of large doses)
102	RACECADOTRIL	REDOTIL	Capsule	Adult: PO 100 mg 3 times/day for ≤7 days.	134.00	Vomiting, nausea, constipation, thirst, abdominal pain, vertigo and headache.
103	RECOMBINANT TISSUES PLASMINOGEN	ELAXIM - 30 MG	Injection	Usual Adult Dose for Myocardial Infarction less than 60 kg: 30 mg IV bolus	30074.00	
104	SALICYLIC ACID	SALICYLIX - SF - 6 % -50GM	Ointment, Shampoo	1 to 2 times daily for up to 2 weeks for corns and calluses and 12 weeks for warts.use as a shampoo, wet hair and scalp. Lather well and leave on as you have been told. Rinse fully	94.00	Itching, redness
		DERSOL - 24% -25GM			99.00	
		SALISIA KT - 60 ML			197.00	
		AKRALIC - 12 %			65.00	
		AKRALIC - 6 %			45.00	
105	SILODOSIN	RAPILIF - 8MG	Tablet	4 - 8mg OD	19.90	Retrograde ejaculation ,Prostate- specific antigen increased, Nasopharyngitis, nasal congestion, sinusitis, , priapism, pharyngeal edema
106	SILVER NITRATE	SILVEREX - 25 GM	Cream	on the affected area 2-3 times per week for 2-3 weeks	83.50	Itching, redness
107	SILYMARIN	SILYBON - 140 MG	Tablet	adult dosage is 140 mg 2-3	107.00	

		SILYBON - 70 MG		times daily.	87.00	
108	SLS, GLYCERIN, LANOLIN, DISODIUM EDTA	AQUASOFT-75GM SOAP	Soap			Itching, redness
109	SODIUM ACID PHOSPHATE & SODIUM PHOSPHATE	PROCTOCLYSIS ENEMIA - 100 ML		Constipation (using Phosphates Enema BP Formula B) Do not administer before	48.00	
110	SODIUM FUSIDATE	CHITOMESH-SF - 10GM	Cream, Ointment	Apply a small amount to the affected area 3 or 4 times daily	100.00	Itching, redness
		FUCIDIN - 15 GM			136.50	
111	SULFASALAZINE DELAYED - RELEASE	SAZO - 500 MG	Tablet	1-2 g 4 times/day	42.25	Headache, anorexia, nausea, vomiting, diarrhoea, abdominal discomfort, photosensitivity, crystalluria, reversible oligospermia
112	TAMSULOSIN	DYNAPRES-0.4MG	Tablet, Capsule	400 mcg once daily.	242.00	orthostatic hypotension, syncope, priapism, headache, infection, asthenia, back pain, chest pain, dizziness, somnolence, insomnia
		URIMAX - 0.4 MG			137.00	
	TAMSULOSIN + FINASTEROII	URIMAX - F			169.00	
	TAMSULOSIN + DUTASTEROI	URIMAX - D			328.50	
113	TOBRAMYCIN	TOBAREN - 5ML EYE DROPS	Drops, Respules	Local treatment of infections ▶ TO THE EYE ▶ Child 1–17 years: Apply twice daily for 6–8 days ▶ Adult: Apply twice daily for 6–8 days Local treatment of infections (severe infection) ▶ TO THE EYE ▶ Child 1–17 years: Apply 4 times a day for first day, then apply twice daily for 5–7 days ▶ Adult: Apply 4 times a day for first day, then apply twice daily for 5–7 days	61.78	
		TOBAMIST - 5ML RESPULES			301.50	
114	TOLTERODINE	ROLITEN - 2 MG	Tablet, Capsule	2 mg twice daily	134.25	Dry mouth, chest pain, headache, somnolence, fatigue, dry skin, abdominal pain, constipation, dyspepsia, diarrhoea, dizziness
		TEROL LA-4			194.50	
115	TRAVOPROST	LUPITROS-3ML DROP	Drops	once a day in the evening.	180.50	Eye irritation
116	TRETINOIN	RETINO - A - 20 GM CREAM	Cream	usually once daily at bedtime	135.00	Itching, redness
117	TROXERUTIN & CALCIUM DOBESILATE	OXERUTE CD	Tablet, Cream	600-1,200 mg/day.	163.00	GI disturbances, headache and flushing.
		OXERUTE - 30 GM			118.00	
118	URSODEOXYCHLolic ACID	UDILIV - 150 MG	Tablet	6 - 15 mg/kg/day in 2-3 divided doses	136.00	Diarrhoea, pruritus, nausea, vomiting gallstone calcification.
		UDILIV - 300MG			238.00	

119	DAPSONE	DAPSONE	Tablet	Multibacillary leprosy 100 mg/day for 12 mth. Paucibacillary leprosy 100 mg/day for 6 mth.	241.50	Anaemia, peripheral neuropathy, haemolysis and methaemoglobinemia (dose-related), nephrotic syndrome, psychological changes, hepatitis.
120	CYCLOSPORINE	C-PSORIN-50mg CAPSULE	Capsule	once daily.	390.00	Chapped lips, Peeling fingertips, palms, and soles, Itching, Scaly skin, Weak nails, Fragile skin, Runny or dry nose, Nosebleeds, Dry mouth, Joint pain.
121	SODIUM PICOSULPHATE	CREMALAX-10mg TABLET	Tablet	Constipation :Child 1 month–3 years: 2.5–10 mg once daily, adjusted according to response ► Child 4–17 years: 2.5–20 mg once daily, adjusted according to response ► Adult: 5–10 mg once daily, dose to be taken at night	104.25	Vomiting, Headache, Nausea.
122	CALCIUM DOBESILATE	DOBESIL-CAPSULE	Capsule	Diabetic Microangiopathy: Initial Treatment: 1.5-2 g/day according to the severity of the case. Maintenance Treatment: 1 g/day. Other Vascular Disorders: 0.5-1 g/day	153.00	Dobesil is Allergic reaction.
123	CALCIUM DOBESILATE + HYDROCORTISONE	DOBESIL-H-30gm CREAM	Cream	about 2-3 times a day	69.5	Application site reactions (burning, irritation, itching and redness).
124	LACTULOSE	DUPHALAC-250ml SOLUTION	solution	once daily for constipation	262.31	Gas, bloating, burping, stomach rumbling/pain, nausea, and cramps.
125	PIRFENIDONE (200MG)	FIBERDONE-200mg TABLET	Tablet	adults for treating Idiopathic Pulmonary Fibrosis is 267 mg orally thrice a day over a period of 1-7 days and 534 mg from days 8-14.	192.95	Unusual tiredness, weakness, dizziness upon standing, diarrhea, weight loss, menstrual period changes, decreased sexual interest or ability, enlarged/tender breasts in men.
126	FINASTERIDE 1 MG	FINAX-TABLET	Tablet	Benign prostatic hyperplasia :► Adult: 5 mg daily, review treatment at 3–6 months and then every 6–12 months (may require several months treatment before benefit is obtained) Androgenetic alopecia in men :► Adult: 1 mg daily	195.00	Impotence, loss of interest in sex, or trouble having an orgasm, abnormal ejaculation, swelling in your hands or feet, swelling or tenderness in your breasts, dizziness, weakness, feeling like you might pass out, headache, runny nose.

127	PHENOXYETHANOL, HYDROXYPROPYL OXIDIZED STARCH PG- TRIMONIUM CHLORIDE, PERFUME, SODIUM BENZOATE, CITRIC ACID MONOHYDRATE AND SODIUM CITRATE	FOLIGLO-100ml SHAMPOO	shampoo	everyday shampoo	207.00	Irritation / redness / itching / burning.
128	LACTATION PREPARATION	GALACT GRANULES-200gm	Granules	Ayurvedic preparation containing asparagus racemosus 15 g, pueraria tuberosa 1 g, anethum sowa 1 g, tribulus terrestris 2.5 g, glycyrrhiza glabra 1 g, cuminum cyminum 1 g, lepidium sativum 1 g. Recommended dosage of medicine is 2 teaspoonfuls or 10 grams twice a day taken with milk.	290.00	Occasional stomach upset, Diarrhea, Tiredness, Absence of a menstrual period in women, Headache, Water and sodium retention, Decreased sexual interest in men.
129		GLYCEP SOAP-75gm	Soap	apply soap to wet body and facetill lather comes,then rinse with water	60.00	No common side effects seen.
		GLYCEROL-100ml	liquid	adult laxative for constipation: The common dose of glycerol is a 2-3 grams in suppository form or a 5-15 mL enema. For children younger than six years old, the dose is a 1-1.7 grams as a suppository or a 2-5 mL enema.	40.00	Headaches,dizziness, bloating, nausea, vomiting, thirst, and diarrhea.
130	HALOBETASOL PROPIONATE	HABICCOR-10gm OINMENT	Ointment	one to two times daily	102.00	Itchiness, dryness, redness, burning and stinging sensation, stretch marks, increased hair growth, folliculitis, hives, warm skin, acne and skin discoloration.
131	HALOBETASOL PROPIONATE + SALICYLIC ACID	HABICCOR-S-15gm OINMENT*	Ointment	once or twice daily		Skin thinning, Dry skin, Application site reactions (burning, irritation, itching and redness).
132	HERBAL PREPARATION	HADJOD-TABLET	Tablet	2 tablets twice daily	147.00	headache, intestinal gas, dry mouth, diarrhea, and insomnia.
133	HALOBETASOL PROPIONATE	HALOX-30ml LOTION	Lotion.	one to two times daily	199.00	itchiness, dryness, redness, burning and stinging sensation, stretch marks, increased hair growth, folliculitis, hives, warm skin, acne and skin discoloration.

134	CALCIUM PANTOTHENATE 10 MG+ELEMENTAL ZINC 10 MG + SILYMARIN 70MG	HEPTOGON-TABLET	Tablet	Recommended dose of Silymarin is 140 mg 2-3 times/day.	159.00	Constipation, diarrhea, or upset stomach.
135	HOMATROPINE HBR	HOMIDE DROPS-5ml	Drops	apply it usually 2 to 3 times daily, up to every 3 to 4 hours,	30.61	Burning/stinging/redness of the eye, eye irritation, or temporary blurred.
136	ALBUMIN (HUMAN)	HUMAN ALBUMIN-20%-100ml INJE	Injection	Hypovolemia :In adults, an intravenous infusion of 25 g should be given. If adequate response (stabilization of circulation) is not achieved within 15 to 30 minutes, an additional dose may be given.In spite of limited information about the efficacy in pediatric subjects, an intravenous infusion of 2.5 to 12.5 g or 0.5 to 1 g/kg body weight may be given. If adequate response (stabilization of circulation) is not achieved within 15 to 30 minutes, an additional dose may be given.Hemodilution may follow administration of Albumin (Human) 20%. If hemorrhage has occurred, this may result in relative anemia. This condition should be controlled by the supplemental administration of compatible red blood cells or compatible whole blood. Hypoalbuminemia :In adults, intravenous infusion of 50 to 75	4332.00	Flush, urticaria, fever and nausea.
137	COLLOIDAL SILVER	HYDROHEAL-AM-15gm GEL	Gel	Gel to the effected area of skin 2 to 3 times daily.	96.00	Drowsiness, dizziness, hypotension or a headache.

138	LACTIC ACID, SORBITOL, COCAMIDOPROPYL BETAINE, POLYQUATERNIUM-7, TEA TREE OIL, HIPPOPHAE RHAMNOIDES.	INTIWASH-50ml	Liquid	during bathing daily, it should be used every time when there is any chance of vaginal pH imbalance, like every time after urination, after unprotected sex, during periods and even during pregnancy. Any woman under the age group of 15-55 can use Intiwash.	90.00	Drowsiness, dizziness, hypotension or a headache.
139	TIMOLOL MALEATE	IOTIM-0.5%-5ml EYE DROPS	Drops	Reduction of intra-ocular pressure in primary open-angle glaucoma Adult: Apply twice daily	55.18	Burning eyes, Stinging in the eyes.
140	PHENYLEPHRINE +TROPICAMIDE	ITROP PLUS-DROPS	Drops	The normal adult dose is 1 drop followed by a second drop after an interval of 5 minutes. A further 1 drop may be instilled after 30 minutes, if required	68.20	Eye pain, Stinging in the eyes, Blurred vision, Photophobia, Allergic reaction.
141		JOHNSON'S COTTON BUDS-15'S			20.00	
142		JOHNSON'S COTTON BUDS-30"S			36.00	
143	CALCIUM-D-PANTOTHENATE	KERABLACK-TABLET	Tablet	consume one tablet every day after a healthy meal. It can either be consumed with a glass of water or juice for best results.	108.90	Constipation, diarrhea, or upset stomach.
144	HYPROMELLOSE+SODIUM CHLORIDE+POTASSIUM CHLORIDE-+SODIUM ACETATE+CALCIUM CHLORIDE-+SODIUM CITRATE +MAGNESIUM CHLORIDE	LACRIGEL-5gm OINTMENT		The usual dose is 1 or 2 drops in each affected eye 3 times daily	72.30	Mild eye burning or irritation, itching or redness of your eyes, watery eyes, blurred vision or unpleasant taste in your mouth.
145	CLOBETASOL , LACTIC ACID, SALICYLIC ACID AND UREA	LOBATE-S-30gm OINTMENT	Ointment	Apply Lobate S Cream to the affected area, usually twice everyday in the morning and evening	125.50	Application site reactions (burning, irritation, itching and redness), Skin thinning.
146	SODIUM CARBOXYMETHYL CELLULOSE	LUBI-FRESH-10ml EYE DROPS	Drops	drops can be used as often as required.	98.00	Stinging in the eyes, Burning eyes, Eye redness, Allergic reaction in eye.
147	OXYBENZONE, HYDROQUINONE, METHOXYCINNAMATE	MELALITE-15-30gm CREAM	Cream	Apply the cream on the affected area twice daily	128.03	Allergies, Burning, Depigmentation, Allergic reaction, Stinging of skin, Itchy, Redness, Acute dermatitis, Exogenous ochronosis.

148	BIOTIN, CALCIUM PANTOTHENATE, ELEMENTAL BORON, ELEMENTAL COPPER, ELEMENTAL IRON, ELEMENTAL MAGNESIUM, ELEMENTAL MANGANESE, ELEMENTAL ZINC, FOLIC ACID, IODINE, NIACINAMIDE, PARA-AMINOBENZOIC ACID, PYRIDOXINE, RIBOFLAVINE, SOY ISOFLAVONES, THIAMINE MONONITRATE, VITAMIN A, VITAMIN C, VITAMIN D2 AND VITAMIN E	MENOPACE ISO-TABLET	Tablet	Adult: One capsule daily,	220.00	Allergic reaction, allergic rejection, allergic reactions, anorexia, nausea, and abdominal distention.
149	MINOXIDIL	MINICHEK-5%-60ml SOLUTION	Solution	Clean and dry the scalp area before applying the medication. You may apply this product to damp hair. To use the solution, fill the applicator with 1 milliliter of medication (to the 1 milliliter line), or use 20 drops. Part your hair in the area of thinning and apply the solution evenly to the affected area of the scalp. Gently rub in. Allow the solution to dry completely before using other styling products (e.g., gels, mousse) or before going to bed.	650.00	Application site irritation, Itching, Headache.
150	UREA+LACTIC ACID +PROPYLENE GLYCOL+LIQUID PARAFFIN	MOISTUREX-100gm CREAM	Cream	A thick layer Cream is applied twice daily after washing the affected area. The cream is left on the skin for 3-5 minutes and then rubbed lightly in.	200.00	Rash, diarrhoea, constipation, abdominal pain, headache, sleeping disorders, dizziness, galactorrhea, gynecomastia.
151	MOMETASONE FUROATE	MOMATE-40gm OINMENT	Ointment	thin film of medication to the affected area once daily.	317.50	Rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

152	MOMETASONE FUROATE	MOTIMESH-5gm CREAM	Cream	Apply a thin film of Mometasone Furoate Cream to the affected skin areas once daily.may be used in pediatric patients 2 years of age or older.	75.00	Application site reactions (burning, irritation, itching and redness).
153	ALPHA LIPOIC ACID, FOLIC ACID, MECOBALAMIN, PYRIDOXINE, THIAMINE MONONITRATE	MYLAMIN-TABLET	Tablet	Adult- PO- The recommended dose range is 150 to 400mg/day	46.75	Allergic reactions, Altered sleep patterns, Nausea, Itch, Dysfunction of nerves resulting in numbness, Vomiting, Vertigo, Anorexia, Abdominal distention, Flatulence.
154	SODIUM CHLORIDE	NASIVION-S-10ml DROPS	Drops	Instil 2-3 drops in each nostril 3-4 times daily	36.19	Allergic contact dermatitis, Nausea, Vomiting, Diarrhea, Muscular twitching, Inflammation of the gastrointestinal tract, Congestion, Eye irritation, Restlessness, Convulsions.
155	SODIUM CHLORIDE \$ SALINE 0.9	NASOCLEAR MIST SPRAY-100ml	Spray	Adults and Children: Apply 2-4 sprays in each nostril 3 times a day.	276.00	Confusion, constipation, dizziness (mild), drowsiness (mild), headache, nausea or vomiting, stomach pain.
156	SODIUM CHLORIDE	NASOCLEAR-20ml DROPS	Drops	2 drops thrice daily for 1 week	40.00	Emesis, Collapse, Nausea, Vomiting, Diarrhea, Muscular twitching, Inflammation of the gastrointestinal tract, Congestion, Eye irritation, Restlessness.
157		NEW MASK-BABY(CIPLA)			225.00	Dry/irritated throat, hoarseness, voice changes, bad taste in the mouth, runny nose, or nosebleeds.
158		NOVOFINE NEEDLE			12.50	Redness, swelling, or itching at injection sites.
159	AMINO ACID	NUTRIVEN 10% I.V-200ml INJECTION*(AMINO ACID)	Injection	Total Parenteral Nutrition (Central Infusion) :In unstressed adult patients with no unusual nitrogen losses, a minimum dosage of 0.1 gram nitrogen (4.2 mL of 15% Amino Acids Injection)plus 4.4 grams (15 calories) of dextrose per kilogram of body weight per day are required to achieve nitrogen balance and weight stability. For patients stressed by surgery, trauma or sepsis, and those with unusual nitrogen losses, the dosage required for maintenance may be as high as 0.3 to 0.4	450.00	Nausea, diarrhea, and stomach pain.

160	SODIUM LACTATE TOPICAL. SODIUM PYRROLIDONE	OLESOFT-200ml LOTION**	Lotion.	twice a day and thrice a day	357.50	Drowsiness, dizziness, hypotension or a headache
		OLESOFT-60gm CREAM**			130.00	
161	OXYMETAZOLINE HCL	OTRIVIN-ADULT NASAL SPRAY	Spray	Adult Dose for Nasal Congestion:2 or 3 sprays in each nostril not more often than every 10 to 12 hours Pediatric Dose for Nasal Congestion: 6 years or older :2 or 3 sprays in each nostril not more often than every 10 to 12 hours	66.50	Temporary burning, stinging, dryness i n the nose, runny nose, and sneezing.
162	POLYETHYLENE GLYCOL 3350 AND ISPAGHULA HUSK GRANULES	PEGFIBER-154.812gm GRANULES	Granules	Bowel preparation:Oral: Adults: 1-2 lts of dissolved solution. Max 4 lts(Peglec soln is prepared by dissolving 1 pack of content in 2 ltrs of boiled and cooled water).	210.00	Nausea, abdominal cramping.
163	POLYETHYLENE GLYCOL ELECTROLYTES FOR ORAL SOLUTION	PEGLEC POWDER-137.15gm	powder	Adults: 1-2 lts of dissolved solution. Max 4 lts (Peglec soln is prepared by dissolving 1 pack of content in 2 ltrs of boiled and cooled water).	316.80	Vomiting, Abdominal cramps, Stomach pain or swelling, Unusual tiredness or weakness, Frequent urge to urinate, Swelling of feet or lower legs, Muscle pain or twitching, Rapid breathing, Unpleasant taste, Mood or mental changes.
164	POLYETHYLENE GLYCOL 3350 POWDER ORAL SOLUTION	PEGMOVE-POWDER	powder	Mix in 1/2 to 1 cup (120 to 240 ml) of liquid like water, juice, soda, coffee, or tea and drink it right away.	173.00	Stomach pain, Nausea, Diarrhoea.

165	LACTULOSE	LAXOPEG SYRUP-200ml	Syrup	Constipation ▶ BY MOUTH ▶ Child 1–11 months: 2.5 mL twice daily, adjusted according to response ▶ Child 1–4 years: 2.5–10 mL twice daily, adjusted according to response ▶ Child 5–17 years: 5–20 mL twice daily, adjusted according to response ▶ Adult: Initially 15 mL twice daily, adjusted according to response	205.80	Diarrhea, Nausea, Abdominal pain, Vomiting, Gas, Electrolyte imbalance due to diarrhea.
166	SILDENAFIL (25MG)	PENEGRA-25mg TABLET	Tablet	Pulmonary arterial hypertension ▶ BY MOUTH:Adult: 20 mg 3 times a day Erectile dysfunction ▶ BY MOUTH ▶ Adult: Initially 50 mg, to be taken approximately 1 hour before sexual activity, adjusted according to response to 25–100 mg (max. per dose 100 mg) as required, to be taken as a single dose; maximum 1 dose per day	109.80	Headache. flushing of the face.
167	PERMETHRIN TOPICAL & 50MG/GM	PERLICE-120gm CREAM	Cream	Apply Perlice Creme/Rinse 1% Cream to the hair, or skin, depending on the location of infection. Avoid its contact with eyes. The same treatment can be repeated after seven days if required	196.22	Mild burning or stinging.

		PERMITE-60gm CREAM		Apply the medicine from your head to the soles of your feet, including under your nails and in skin folds such as between the toes, as directed. Massage the cream into the skin. Do not use more medication than prescribed. Wash off the cream after 8-14 hours by showering or taking a bath. Demonstrable living mites after 14 days indicate that retreatment is necessary. Permethrin cream is safe and effective in pediatric patients two months of age and older.	106.17	
168	SUNSCREEN LOTION	PHOTOBAN-30-100gm AQUAGEL	Gel	apply over the face and neck, half an hour before going out into sunlight	339.50	Dermatitis, Contact dermatitis, Burning of skin, Irritation of skin, Skin irritation.
169	PIMECROLIMUS TOPICAL	PICON-1%-10gm CREAM	Cream	Apply a thin layer to the affected areas of skin, usually twice daily	400.00	Itching, rash, and redness.
170	TROXERUTIN, CALCIUM DOBESILATE, ZINC, PHENYLEP	PILORUTE-30gm CREAM	Cream	Apply on the affected area of the skin after cleaning and drying it well. It is usually recommended to use about 2-3 times a day.	115.50	Burning or stinging sensation, dizziness, drowsiness, changing body temperatures, blurred vision or a pounding/ringing in your ears.
171	PODOPHYLLUM RESIN BENZOIN ALOES SOLUTION	PODOWART-10ml PAINT	Solution	twice a day and once a week, Clean and dry the skin area to be treated. do not wash the treated area after immediately applying Podowart Paint Lotion.	154.00	severe burning, stinging, itching, redness, swelling, or skin changes where the medicine was applied, numbness, burning, pain, or tingly feeling, easy bruising or bleeding, weakness, fever or severe constipation or stomach cramps.

172	MALTODEXTRIN, SUNFLOWER SEED OIL, CALCIUM CAEINATE, WHEY PROTEIN ISOLATE, ISOMALTULOSE, FRUCTOSE, SOY PROTEIN ISOLATE, RAPESEED OIL, GUM ARABIC, OLIGOFRUCTOSE, MINERALS, ANTIOXIDANT, VITAMINS, CARNITINE, TAURINE AS ACTIVE INGREDIENTS.	PROHANCE-D VANILLA-400gm PO	powder	50 gm Daily or as per requirement	600.00	Nausea, or unusual or unpleasant taste in your mouth.
173	LACTOBACILLUS ACIDOPHILUS FIDOBACTERIUM	PROVIDAC-CAPSULE	Capsule	daily in 3-4 divided doses.ADULTS: BY MOUTH: For hayfever: At least 2 billion colony-forming units of lactobacillus daily for 7 weeks has been used along with 10 mg of loratadine once daily for 5 weeks. For preventing diarrhea caused by antibiotics: Many different lactobacillus species have been studied. In most cases, lactobacillus is given in daily doses that provide 10-100 billion colony-forming units daily. Lower doses of 100 million colony-forming units have also been used. Usually treatment is started within 2	275.60	Stomach pain and discomfort, bloating, and incomplete bowel movements.
174	CARBOXYMETHYLCELLULOSE	REFRESH-10ml LIQUIGEL	Drops	drops may be used as often as needed. squeeze out 1 or 2 drops as needed. Look down and gently close your eye for 1 or 2 minutes.	180.07	Eye pain, change in vision, continued eye redness/irritation.

175	FOLIC ACID(0.8 MG), MECOBALAMIN(0.5 MG), PYRIDOXINE(20 MG)	ALZIVIT-TABLET	Tablet	<p>Adult- Treatment of folate-deficiency, megaloblastic anaemia: 5 mg daily for 4 months (up to 15 mg daily may be necessary in malabsorption states).</p> <p>Prevention of first occurrence of neural tube defect: 400 to 500 µg daily before conception and during the first twelve weeks of pregnancy.</p> <p>Prevention of recurrence of neural tube defect: 5 mg daily (reduced to 4 mg daily, if suitable preparation available) from at least 4 weeks before conception until twelfth week of pregnancy</p>	8.25	Vomiting, Allergic reactions, Anorexia, Nausea, Abdominal distention, Flatulence, Bitter or bad taste, Altered sleep patterns, Difficulty in concentrating, Irritability.
176	SALICYLIC ACID AND LACTIC ACID	SALACTIN-10ml LOTION	Lotion.	<p>First wash the affected area. You may soak it in warm water for about 5 minutes to soften it. Dry thoroughly. Removing the dead skin helps the medication to work better. Do not try to rub the wart, corn, or callus off.</p> <p>Using the applicator provided, apply a thin coat of the medication (one drop at a time) to cover the entire wart, corn, or callus. To minimize irritation, be careful not to get the liquid on surrounding skin. Let dry.</p> <p>Repeat this procedure 1 to 2 times daily as needed until the wart, corn, or callus is removed.</p> <p>For warts, you may use this product for up to 12 weeks. For corns and calluses, you may use it for up to 2 weeks. Wash hands after use.</p>	271.00	Rash, diarrhoea, constipation, abdominal pain, headache, sleeping disorders, dizziness, galactorrhea, gynecomastia.

177	FLUOCINOLONE TOPICAL	SEBOWASH-100ml SHAMPOO	shampoo	Apply about one ounce of the shampoo to the scalp once daily by working into a lather. Allow it to remain on the scalp for 5 minutes.	175.00	Application site reactions (burning, irritation, itching and redness).
178	POTASSIUM NITRATE 3 %+SODIUM FLUORIDE	SENQUEL-AD-100ml MOUTH WASH	Mouthwash	recommend that you swish the product around in your mouth for 30 seconds to one minute, then spit it out.	55.50	Dizziness, laboured breathing, abdominal pain, blue fingernails, lips or skin, diarrhoea. a headache, convulsions, confusion, nausea, unconsciousness.
179	POTASSIUM NITRATE 5 %W/W+SODIUM MONOFLUOROPHOSPHATE	SENQUEL-F-PASTE-50gm*	Paste	for external use .once daily	71.00	Abdominal bloating, Dental fluorosis.
180	SOLUTION OF SORBITOL AND TRICHOLINE CITRATE	SORBILINE-100ml SYRUP	Solution	Hepatitis: 15ml t.i.d. following meals. CHILDREN: 5-10ml. Constipation: 10-20ml before breakfast. CHILDREN: 5-10ml before breakfast.	69.87	Gas, Diarrhea, Debilitation, Urge to vomit, Dizziness, Anal irritation, Abdominal cramping, Abdominal bloating, Dizzines, Acute allergic reaction.
181		STAY FREE (DRY WINGS)			30.00	
182		STREPSILS			2.92	
183	MULTIVITAMIN, MINERALS	SUPRADYN-TABLET	tablet	1-2 tabs daily	28.05	Nausea, Vomiting, Diarrhea, Constipation, Upset stomach, Black or dark-colored stools or Temporary staining of the teeth.
184	TACROLIMUS TOPICAL	TACORYZA-0.01-10gm CREAM	Cream	Apply a thin layer to the affected areas of skin, usually twice daily	388.70	Application site reactions (burning, irritation, itching and redness).
		TACROTOR-0.03%-10gm OINTMEN	Ointment	Apply a thin layer to the affected areas of skin, usually twice daily	154.40	
		TACROZ FORTE-0.1%-10gm OINTM	Ointment	Apply a thin layer to the affected areas of skin, usually twice daily	461.50	
		TACROZ-0.03%-10gm OINTMENT**	Ointment	Apply a thin layer to the affected areas of skin, usually twice daily	177.00	
185	SODIUM TALLOWATE, SODIUM COCOATE, WATER, GYLCEIN, FRAGRANCE,SODIUM CHLORIDE, SODIUM HYDROXIDE, TETRASODIUM ETIDRONATE, O-TOLYL BIGUANIDE,TITANIUM DIOXIDE	TETMOSOL-100gm SOAP	Soap	For external application. Apply soap till lather comes,rinse with water	61.00	itching, skin irritation, inflammation, rashes, redness on skin.

186	URSODEOXYCHOLIC ACID / URSODIOL (300MG)	URSOCOL-300mg TABLET	Tablet	13-15 mg/kg/day administered in two to four divided doses with food.	240.00	Upset stomach, indigestion, or vomiting. Dizziness.
187	ACESULFAME POTASSIUM, CALCIUM SODIUM PHOSPHOSILICATE, CARBOMER, GLYCERIN, PEG 400, SILICA, SODIUM LAURYL SULPHATE AND TITANIUM DIOXIDE	VANTEJ-50gm PASTE	Paste	Calcium sodium phosphosilicate in Vantej toothpaste imparts significant level of dentinal occlusion with durable occlusive deposits following four days of twice daily brushing.	116.00	Puffiness of face, Cough syrup, Urge to vomit, Abdominal cramps, Wind, Diarrhea, Rash, Allergic reaction, Glycerin laxative-anal, Bowel irritation.
188	ALMOND BUTTER TOPICAL, CETYL ALCOHOL TOPICAL, COCOA BUTTER TOPICAL, CYCLOMETHICON E TOPICAL, DIMETHICONE TOPICAL, DISODIUM EDETATE TOPICAL, EMULSIFYING WAX TOPICAL, FRAGRANCE TOPICAL, GLYCERIN TOPICAL, MANGO BUTTER TOPICAL, METHYL PARABEN TOPICAL, PROPYLENE GLYCOL TOPICAL, PURIFIED WATER TOPICAL, SHEA BUTTER TOPICAL, STEARIC ACID TOPICAL AND ZINC OXIDE TOPICAL	VENUSIA MAX-150gm CREAM	cream	topical applicaltion all over the body after bath apply twice a day	355.00	Burning of skin, Redness of skin, Inflammation of chinchilla middle ears, Contact dermatitis, Mild lactic acidosis, Burning, Stinging, Erythema, Skin irritation, Pruritus.

	BUTTER TOPICAL, METHYL PARABEN TOPICAL, PROPYLENE GLYCOL TOPICAL, PURIFIED WATER TOPICAL, SHEA BUTTER TOPICAL, STEARIC ACID TOPICAL AND ZINC OXIDE TOPICAL	VENUSIA MAX-300gm LOTION	Lotion.	topical applicaltion all over the body after bath apply twice a day	468.00	
189	ALOE VERA TOPICAL, SQUALENE TOPICAL AND VITAMIN E	VENUSIA-75gm CREAM	cream	apply over the body both morning and evening	265.00	Erythema multiforme, No known side effects.
190	BETAHISTINE (24MG)	VERTIREST-OD-TABLET	Tablet	Vertigo, tinnitus and hearing loss associated with Ménière's disease :Adult: Initially 16 mg 3 times a day, dose preferably taken with food; maintenance 24-48 mg daily	135.00	Headache, Nausea, Dyspepsia.
191	MENTHOL,METHYL SALICYLATE,DICLOFENAC DIETHYLAMMONIUM,LINSEED OIL	VOLINI-40gm SPRAY	spray	spray over the affected area local application whenever needed	125.00	Rash, No side effects reported, Drowsiness, Anaphylactic reactions, Allergic contact dermatitis, Pruritus, Dermatitis.
192	MINOXIDIL	ZENOXIDIL-2%-60ml GEL	Topical solution	FOAM men Androgenetic alopecia ▶ TO THE SKIN ▶ Adult: Apply 0.5 capful twice daily, to be applied to the affected areas of scalp; discontinue if no improvement after 16 weeks SOLUTION Androgenetic alopecia ▶ TO THE SKIN ▶ Adult: Apply 1 mL twice daily, to be applied to the affected areas of scalp; discontinue if no improvement after 1 year FOR	225.00	Burning, stinging, or redness at the application site.

193	CYNOCOBALAMINE	CYNOCOBALAMINE INJECTION-3	Injection	Pernicious Anemia :Adult Dosages 100 mcg intramuscularly or subcutaneously once daily for 6-7 days, then every other day for 7 doses, then 3-4 days for 2-3 weeks, then monthly /Alternative parenteral dosing: 1,000 mcg intramuscularly or subcutaneously once daily for 7 days, then weekly for 1 month, then monthly Pediatric Dosages :30-50 mcg intramuscularly or	30.00	Pain/redness at the injection site, mild diarrhea, itching, or a feeling of swelling all over the body.
194	CICLOPIROX +ZINC PYRITHI	8X - 100ML SHAMPOO	Shampoo	wet your hair, take approximately 1 teaspoon of 8x shampoo (or 5ml) apply onto the scalp. Or u can use for long and thick hair 2 teaspoons of 8x shampoo (10ml).Apply it equally onto the affected area and massage into scalp, Leave it for 3 minutes. The eyes must be protected and rinse off with water.The procedure should be repeated twice in a week for 1 month. There should be minimum 3 days of gap per each cycle.	273.35	Burning sensation, redness, swelling, irregular heartbeat, rash, dandruff, headache and skin problems.
195	ALOE VERA +D-PANTHENOL	KERAGLO AD-75ml SHAMPOO	shampoo	Apply to wet hair, massage into scalp, leave on hair for 5 mins and rinse thoroughly with water.	273.90	Nausea, headache, diarrhea, stomach pain and dizziness.
196	MELANOSTALINE	ADVAN-10gm CREAM	Cream	_once a day and twice a day	446.00	Mild burning, redness, stinging and dryness.
197	CARICA PAPAYA	CARIPILL-TABLET	Tablet	Adult :One Tablet to be taken 3 times daily for 5 days. Children aged 6 to 12 years : 10 ml 3 times daily for 5 days	410.50	Severe stomach/abdominal pain, nausea/vomiting, slow heartbeat, severe drowsiness, inability to move.
198	KOJIC ACID	KOJIC-25gm CREAM**	Cream	Wash your face and neck thoroughly before each use Then, lightly massage the Kojic cream onto the surface of the skin Use twice a day on your face and neck	325.00	Redness, irritation, itchiness, rashes, swollen skin, or pain and discomfort.

199	KOJIC ACID(12%W /W) , OCTII	KOJIVIT PLUS-15gm GEL	Gel	Take small quantity of cream and rub on face in circular motion till it disappears and dry completely.Use this cream every time you wash your face to notice visible effects.	325.00	Feeling tired or weak (fatigue), stomach pain, loose stools (diarrhea), headache, nausea, or blurred vision.
200	CETEARETH,PURIFIED WATER, GLYCERIN, HYDROGENATED POLYISOBUTENE, CETEARYL ALCOHOL,	CETAPHIL DAM-30gm LOTION	Lotion.	<p><u>Cetaphil Lotion Intense gel</u> :Apply & gently rub into the scar once a day for 8 wk (on new scars) & 3-6 mth (on existing scars).</p> <p><u>Cetaphil Lotion Proactive gel</u> : Apply & gently massage into the scar at least bid for 8 wk (on new scars) & 3-6 mth (on existing scars).</p> <p><u>Cetaphil Lotion PM Overnight cream</u> : Apply & gently rub into scar once nightly for 8 wk (on new scars) & 3-6 mth (on existing scars).</p> <p><u>Cetaphil Lotion Kids cream Scars</u> : Apply a thin coat to the scar & gently massage in tid for 8 wk (on new scars) & tid for 3-6 mth (on existing scars).</p> <p>Sunscreen : Apply liberally 15 min before sun exposure. Reapply at least every 2 h</p>	152.00	Stinging, contact dermatitis, and burning.

201	HYALURONIDASE 1500IU	ENTODASE-INJECTION	Injection	<p><u>Subcutaneous Fluid Administration (Hypodermoclysis)</u> :For infants and children less than 3 years old, the volume of a single clysis should be limited to 200 mL; and in premature infants or during the neonatal period, the daily dosage should not exceed 25 mL/kg of body weight; the rate of administration should not be greater than 2 mL per minute. For older patients, the rate and volume of administration should not exceed those employed for intravenous infusion.</p> <p><u>Subcutaneous Urography</u> : 75 Units of Entodase (hyaluronidase injection) is injected subcutaneously over each scapula, followed by injection of the contrast medium at the same sites</p>	131.00	Acute urticaria, Hormone imbalance, Allergic reaction, Slow growth in children and teenagers, Angioedema (swelling of deeper layers of skin), Dizziness, Injection site reaction, Edema (swelling).
202	VITAMIN E 400 MG	EVION-400mg CAPSULE	Capsule	one capsule daily	24.58	Nausea, diarrhea, stomach cramps, tiredness, dizziness, blurred vision, and headache may rarely occur.
203	LEVOCARNITINE 150 MG+VITAMIN E 200 MG	EVION-LC-TABLET	Tablet	1 to 2 tablets in a day.	35.42	Rare incidences ofgastrointestinal distrubances such as nausea, vomiting, diarrhoea, abdominalcramps,

204	CALCIUM ASCORBATE 75 MG+COENZYME Q10 100 MG+DOCOSAHEXAENOIC ACID 60 MG+EICOSAPENTAENOIC ACID (EPA) 90 MG+L-ARGININE 100 MG+LEVOCARNITINE 50 MG+LYCOPENE 5000 MCG+MIXED CAROTENE 10.33 MG+PIPERINE 5 MG+SELENIUM DIOXIDE 100 MCG+VITAMIN B12 5 MCG+WHEAT GERM OIL 25 MG+ZINC SULPHATE 6.5 MG	KQ-100-CAPSULE	Capsule	once a day	425.00	Very upset stomach or throwing up. Very loose stools (diarrhea).
205	CYANOCOBALAMIN 2.5 MCG+ELEMENTAL IRON 30 MG+FOLIC ACID 0.5 MG	RUBIRED-150ml SUSPENSION	Suspension	<p><u>Adult Dose for Megaloblastic Anemia</u> :1 mg orally, intramuscularly, subcutaneously or IV once a day.</p> <p><u>Adult Dose for Folic acid (Rubired SUSP) Deficiency</u> 400 to 800 mcg orally, intramuscularly, subcutaneously or IV once a day. Women of childbearing age, pregnant, and lactating women: 800 mcg orally, intramuscularly, subcutaneously or IV once a day.</p> <p><u>Pediatric Dose for Folic acid (Rubired SUSP) Deficiency</u> Infant: 0.1 mg orally, intramuscularly, subcutaneously or IV once a day. Child: Less than 4 years: up to</p>	139.00	Pulmonary edema, Congestive heart failure, Polycythemia vera, Diarrhea, Itching, Feeling of swelling of entire body, Pain in upper central abdomen, Darky stool may occur, Vomiting, Urge to vomit.

206	PROTEIN	PROSURE-400gm POWDER	powder	The recommended daily intake is 2 servings/day. 2 servings a day should be consumed for at least 3 weeks to see results. ProSure tastes best if served chilled. The supplement should be shaken well. Tube Feeding: Patients on tube feeding can receive the benefits of ProSure by providing 2 servings (240 mL) in conjunction with a standard tube feeding formula. In a tube feeding regimen, providing 2 servings of ProSure a day is suggested to achieve the recommended level of 2 g EPA,	789.00	Asthma, Allergic rhinitis with hives, Eczema, Increased bowel movements, Nausea, Thirst, Bloating, Cramps, Reduced appetite, Tiredness.
		PROTINEX-250gm POWDER	powder	1 serving of 25 gm (2 to 3 heaped tablespoons) per day in 200 ml of milk.	315.00	
		ENSURE VANILLA-400gm POWDER	powder	Directions for use: to prepare a 230 ml feeding, put 195 ml of cold water in a glass. While stirring, gradually add 6 level scoops (scoop enclosed) of Ensure powder and mix until dissolved.	480.00	
207	FIBER	FYBOGEL-100gm POWDER	powder	Adults and children over 12yrs: 1 measureful (5.4gm) in morning and at bedtime or 2 measureful at bedtime. Children between 6-12yrs: Half to one measureful in the morning and at bedtime or 1-2 measureful at bedtime. Children below 6yrs: Not recommended except under medical advice.	130.00	wind and bloating.
208	ACOTIAMIDE (100MG)	ACOTRUST-100mg TABLET	Tablet	Dose:100 mg tid	190.00	Headache, Diarrhoea.
209	CETRIMIDE	CETRILAK-100ml SOLUTION	solution	Topical/Cutaneous- wound cleansing,Adult :Apply 0.1-1% aqueous solution or 0.5% cream onto affected area.Seborrhoeic dermatitis Adult: Apply 10% shampoo solution onto the scalp.	30.00	Drowsiness, dizziness, hypotension or a headache.