


Sri
MANAKULA VINAYAGAR
Medical College and Hospital

Kalitheerthalkuppam, Madagadipet, Puducherry – 605107.

Department of Pharmacy
DRUG FORMULARY



Edition-II

Supplementary - 01 / 2019

PREFACE

This Drug Formulary of Sri Manakula Vinayagar Medical College and Hospital is a concise and handy book designed for quick reference of current essential drugs in medical practice. As it is not possible to include all drugs, commonly used drugs in our hospital with their recommended dosages, both in adults and children, adverse drug reaction and cost have been included. However, as the medical knowledge is vast and continuously being upgraded, changes may have occurred and may not have been reflected in the incorporated information. Clinical judgment and experience should be used while prescribing the drugs.

Note: Dose of drug varies with indications and severity of disease, the given doses is just for reference, and is for the most common indication. Where feasible, the dose ranges from lowest to maximal permissible doses have been given. Pediatric doses are for children more than 1 month of age i.e. neonatal doses may vary. The price given is for an estimation of the cost of therapy and not the exact selling price, which may vary.

“Let food be thy medicine and medicine be thy food.”

-Hippocrates

CONTENTS

SL. NO.	SECTION	PAGE NO.
1	ANTI-ASTHMA, DECONGESTANTS , OTHER DRUGS	1
2	ANTI-BACTERIALS	1
3	ANTI-CANCER	4
4	ANTI-COAGULANTS AND THROMBOLYTIC	6
5	ANTI-DEPRESSANTS	9
6	ANTI-EMETIC	10
7	ANTI-EPILEPTICS	10
8	ANTI-FUNGALS	11
9	ANTI-HISTAMINE	12
10	ANTI-HYPERTENSIVE	12
11	ANTI-PARASITIC	13
12	ANTI-PARKINSONISM	13
13	ANTI-PSYCHOTICS	13
14	ANTI-ULCER	14
15	ANTI-VIRAL	14
16	CNS - OTHERS	15
17	CVS - OTHERS	18

18	HORMONES & THEIR ANTAGONIST	18
19	MUSCLE RELAXANTS, ANAESTHETIC & EMERGENCY	20
20	NSAIDS & GOUT	21
21	SEDATIVES AND HYPNOTICS	22
22	VACCINE	23
23	MISCELLANEOUS	23

**“Writing prescriptions is easy, understanding people is
hard”**

- Franz kafker, 1883-1924

ANTI-ASTHMA, DECONGESTANTS , OTHER DRUGS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	LEVOCETRIZINE	LEVOCKAST Z-TABLET	Tablet	5 mg orally once a day in the evening	16.50	Dizziness, Fatigue
		LEVOCETNE Z TABLET	Tablet		8.90	
		XYZAL-5mg TABLET	Tablet		9.07	
2	SALMETEROL+ FLUTICASONE PROPIONATE	SEROFLO-250 SYNCHROBREATH	Inhaler, Rotacaps	1 inhalation orally twice a day, approximately 12 hours apart	587.40	Mouth & throat candidiasis, throat irritation
3	DOXOFYLLINE	DOXOLIN-SR-400mg TABLET	Tablet	The recommended dose is up to 1,200 mg/day.	2.89	Restlessness, tremor, palpitation, tachypnoea, flushing, hypotension
4	FLUTICASONE PROPIONATE	NEZAFLO-NASAL SPRAY	Nasal spray	The usual starting dose is 2 sprays into each nostril once a day, preferably in the morning.	262.00	Topical: Pruritus, hypertrichosis, dryness, numbness of fingers, burning, eruptions, hypopigmentation,
5	BEPOTASTINE	BEPOZIC-10mg TABLET	Tablet	Once a day	10.80	Headache Mild taste disturbances Skin rash Redness of the eye

ANTI-BACTERIALS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMOXYCILIN & POTASSIUM CLAVULANATE	SENSICLAV-375mg TABLET	Tablet	Amoxycillin Clavulanic Acid	18.83	Diarrhea, Alteration of
		SENSICLAV-625mg TABLET	Tablet	Adult: 250, 500 mg	14.86	Bacterial flora, rashes
				Paediatric dose: 20- 40 mg/kg/day, Clavulanic acid: 125mg		HS reactions
2	AZITHROMYCIN	AZITHR Z-TABLET	Tablet	Adult dose: 500 mg OD on day 1 then 250 mg OD Paediatric dose: 10mg/kg/day on day 1 then 5mg/kg/day	22.39	Mild GI upset, Headache
3	GATIFLOXACIN + PREDNISOLONE	GATE-P EYE DROPS	Eye Drop	One / two drop in the affected eyes	33.50	Fluid retention, alteration in glucose tolerance, increased appetite, high blood pressure

4	TRAVOPROST	TRAVO-Z-3ml EYE DROPS	Eye Drop	The recommended dosage is one drop in the affected eye(s) once daily in the evening.	540.00	Blurred vision Eye pain Dry eyes Redness of the eye Burning or itching of eyes Eye discharge Headache
5	MOXIFLOXACIN (0.5 %W/V) + DEXAMETHASONE (0.1 %W/V)	MOXIGRAM-D-5ml EYE DROPS	Eye Drop	One / two drop in the affected eyes	64.00	Blurred vision Dry eyes Headache Eye irritation and discomfort Dizziness Blurred vision
6	CLINDAMYCIN	FACECLIN-AT-15gm GEL	GEL	Adult dose: 600 -1800 mg per day in four divided doses i.m /i.v: 300 - 2400 mg/day in 2-4 divided doses child <1 mth: 15-20mg/kg /day in 3-4 divided doses child>1mth: 20-40mg/kg/day in 3-4 divided doses	199.50	Diarrhea, skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness
		CLAMIN-300mg/2ml INJECTION(CLINDAMYCIN)	Injection		108.00	
7	PIPERACILLIN & TAZOBACTAM	PIPTUM- 4.5gm-INJECTION(PIPERACILLIN&TAZ OBACTAM	Injection	Adult: 2.25- 4.5 g IV 6 hrly	398.00	GI disturbance, hypokalemia, hyponatremia, platelet dysfunction
8	ISONIAZID	SOLOX-300mg TABLET (ISONIAZID)	Tablet	Adult: PO Each tab contains rifampicin 120 mg, isoniazid 50 mg and pyrazinamide 300 mg: <40 kg: 3 tab/day; 40-49 kg: 4 tab/day; 50-64 kg: 5 tab/day; ≥65 kg: 6 tab/day.		GI symptoms,pseudomembranous colitis; flushing and itching w/ or w/o rash, urticaria, pruritus, erythema, rash; hepatitis; thrombocytopenia w/ or w/o purpura, eosinophilia, leucopenia, oedema, muscle weakness, loss of tendon reflexes, paraesthesia, myopathy; vertigo, polyneuritis; occasional menstrual cycle disturbances. Flu-syndrome
9	LINEZOLID	LINID-300ml INFUSION (LINEZOLID)	Injection	Adult: PO 400 mg 12 hrly PO/IV600 mg 12 hrly	504.70	Reversible myelosuppression (e.g. leucopenia, thrombocytopenia, anaemia, pancytopenia); headache, fever, convulsions and seizures, rashes, dizziness, nausea and vomiting, metallic taste, insomnia, constipation, diarrhoea, abdominal pain, skin disorder, cough, pharyngitis

10	LEVOFLOXACIN	LOXOF-500mg TABLET	Tablet	Once daily with or without food	8.47	Diarrhea Abdominal pain Confusion Fever Redness of skin Change in taste Constipation Dizziness Headache Nausea or vomiting
11	POVIDONE -IODINE	DREZ SPRAY-40gm	Spray	Topical: Clean affected area. Apply small amount topically to affected area 1—3 times daily. May cover with sterile bandage. Do not use for longer than 1 week. Pessary: one pessary for a period of 7 days Gargle: Adults, elderly and children over 6 years of age:	117.13	Skin irritation, Severe allergic reaction
		CIPLADINE OINTMENT-125GM-003	Ointment	Use undiluted with an equal volume of warm water. Gargle or rinse with up to 10ml for up to 30 seconds without swallowing. Repeat up to four times daily, for up to 14 consecutive days. Not to be used in children of 6 years and under.	69.90	
12	CEFPODOXIME, OFLOXACIN	CEPODEM-O TABLET	Tablet	Three times a day		Upset stomach or throwing up. Many small meals, good mouth care, sucking hard, sugar-free candy, or chewing sugar-free gum may help. Loose stools (diarrhea).
13	FOSFOMYCIN TROMETHAMINE	MONUROL-SACHET	Sachet	1 packet (sachet) as a single dose. Mix this drug with water before taking it. Pour the contents of 1 packet (sachet) into a half glass (4 ounces or 120 milliliters) of cold water and stir to dissolve. Do not use warm or hot water. Drink all of the mixture right away.	380.00	Nausea, diarrhea, headache, vaginal discomfort, or dizziness may occur

14	CEFUROXIME	CEROXIM-500mg TABLET	Tablet	The recommended dose range is 125mg to 3gms day	35.60	Diarrhea, Chills, Headache, Itching of vagina or genital area Chest pain Difficult or painful urination Shortness of breath
		CEROXIM-250mg TABLET	Tablet		19.80	

ANTI-CANCER

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	RITUXIMAB	CYTOMAB* 500mg/50ml-INJECTION (RITUXIMAB)	Injection	One Vial of 50ml of concentrate for solution for infusion For intravenous administration after dilution	28000.00	Low blood pressure Nausea and Vomiting Chills and fever Abdominal pain Decrease in white blood cells and platelets Itching Infections Headache Muscle aches and pain Cough
		ENFIERA-500mg INJECTION(RITUXIMAB)	Injection		28000.00	
2	CYLOPHOSPHAMIDE	ENDOXAN-50mg TABLET (CYCLOPHOSPHAMIDE)	Tablet	Malignancies Low dose regimen: 2-6 mg/kg/wk in divided doses. Moderate dose regimen: 10-15 mg/kg /wk; high dose regimen: 20-40mg/kg	4.00	Congestive heart failure; leucopenia poor wound healing; anorexia. Nausea, vomiting; alopecia; oral mucosal ulceration; anemia thrombocytopenia, nonhaemorrhagic cystitis.
3	VINBLASTINE	CHEMOBLAST-10MG/10ml-INJECTION (VINBLASTINE)	Injection	Once weekly. 1st dose: 3.7mg/ml as a single dose, continue to increase dose by increments	201.50	Nausea. Vomiting. Decreased appetite. Decreased white blood cell count. Diarrhoea.
4	ERLOTINIB	ERLIB-150mg(ERLOTINIB)-30S	Tablet	once daily	3380.10	Diarrhea, dry skin, muscle/joint pain, mouth sores, unusual eyelash growth, or eye dryness/redness/irritation may occur

5	ETOPOSIDE	ETOPLAST-100mg*/5ml-INJECTION (ETOPOSIDE)	Injection	Testicular cancer- Adult: IV- The recommended dose is 50-100 mgm 2day from days 1-5. Small cell lung cancer- The recommended dose is 35 mgm 2day for 4 days to 50 mgm 2day for 5 days. May repeat course 3-4 weekly.	131.00	Stay away from people with infections, colds, or flu. Anemia, low white blood cell count, and low platelet count. Feeling tired or weak. Upset stomach or throwing up. Many small meals, good mouth care, sucking hard, sugar-free candy, or chewing sugar-free gum may help. Loose stools (diarrhea). Irritation or swelling where the shot was given. Hair loss
		ETOPLAST-50mg CAPSULE(12s)-(ETOPOSIDE)	Capsule		585.00	
6	METHOTREXATE	ONCOTREX-7.5mg TABLET	Tablet	Tab: 2.5 - 30 mg/day	11.91	Ulceration of the mouth and GI disturbances (e.g. stomatitis and diarrhoea), bone marrow depression, hepatotoxicity, renal failure, skin reactions, alopecia, ocular irritation, arachnoiditis
7	CARBOPLATIN	KEMOCARB-450mg/45ml-INJECTION (CARBOPLATIN)	Injection	Adult(Previously untreated patient):400mg/m ² as single short term IV infusion over 15-60min. If previously treated reduce initial dosage by 20-25%(300-320mg/ml)	2433.82	Pain at the injection sites, Low WBC and Platelet counts, Fever and chills.
8	TRAZTUZUMAB	CANMAB-440mg INJECTION (TRAZTUZUMAB)	Injection	6 mg/kg as an intravenous infusion over 30-90 minutes every three weeks.	59981.60	Fever Headache Nausea and Vomiting Tiredness and weakness Shortness of breath Diarrhea Reduction in blood cell count Infections
9	DACARBAZINE	ZICARB-200mg- INJECTION (DACARBAZINE)	Injection	150 mg/m ² IV a Day for 5 days, OR 375 mg/m ² IV on Day 1; repeat every 15 Days	215.00	Nausea Vomiting Injection site pain Leukopenia Thrombocytopenia
		ZICARB-500mg- INJECTION (DACARBAZINE)	Injection		470.45	

10	IMATINIB	CHEMOTINIB-400mg TABLET(IMATINIB)	Tablet	400 mg a Day		Neutropenia Nausea Muscle cramps Musculoskeletal pain Thrombocytopenia, Rash
11	PACLITAXEL NANOPARTICLE	NANOXEL-300mg/15ml- INJECTION (PACLITAXEL NANOPARTICLE)	Injection	Adult:135mg/ml infused over 24 hr and repeat at 3 weeks intervals.	19621.55	Difficult or painful urination, Dizziness when getting up suddenly from a sitting position, Fever and chills.
12	ZOLEDRONIC ACID	OSTEOMET-4mg INJECTION (ZOLEDRONIC)	Injection	Adult: 4-5mg as a single infusion over at least 15 min. Retreatment after atleast 7 days if necessary.	387.50	Headache, Back pain, Musculoskeletal (bone, muscle or joint) pain, Indigestion, Heartburn, Diarrhoea.
		ZOBONE-4mg-INJECTION (ZOLEDRONIC ACID)	Injection		2500.00	
13	CURCUMA LONGA	ADCUMIN-CAPSULE(60S)	Capsule	12 g/day over 3 months		Nausea Gastric irritation Stomach upset
14	PROCARBAZINE	P-CARZINE-50mg- CAPSULE(PROCARBAZINE)	Capsule	2-4 mg/kg/day in single or divided doses for 7 days; increase dose by 4-6 mg/kg/day until maximum response obtained or WBC<4,000 cells/mm ³ or plateletes<100,000 cells/mm ³ ; reduce dosage to 1-2 mg/kg/day after maximum effect obtained Alternatively, 100 mg/m ² PO Day for 7-14 days of each 28 day cycle in combo with other antineoplastics	33.70	Constipation Hair loss Dry mouth Increased urination Chest pain Headache

ANTI-COAGULANTS AND THROMBOLYTIC

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	CILOSTAZOL	CILOBID-100mg TABLET	Tablet	100 mg bid	12.20	Dizziness, palpitations, diarrhoea, abnormal stools; pain, infection; peripheral oedema,
2	FONDAPARINUX SODIUM	FONDAFLO-2.5mg/0.5ml INJECTION (FONDAPARINUX SODIUM)	Injection	The usual dose is 2.5 mg once a day, injected at about the same time each day. If there is kidney disease, the dose may be reduced to 1.5 mg once a day.	784.50	Bleeding, Anemia (low number of red blood cells).

3	BOVINE-DERIVED GELATIN MATRIX	FLOSEAL-5ml	Injection	<p>The mixing accessories include a syringe, a Luer connector, a small bowl, and a 5 mL syringe with needle attached. The mixing accessories are included to facilitate the reconstitution and mixing of the Thrombin into the Gelatin Matrix</p>	14990.00	<p>Gelatin-based hemostatic agents may serve as a nidus for infection and abscess formation and have been reported to potentiate bacterial growth.</p> <ul style="list-style-type: none"> • Giant cell granulomas have been observed at implant sites when used in the brain. • Compression of the brain and spinal cord resulting from the accumulation of sterile fluid have been observed. • Multiple neurologic events were reported when absorbable gelatin-based hemostatic agents were used in laminectomy operations, including cauda equina syndrome, spinal stenosis, meningitis, arachnoiditis, headaches, paresthesias, pain, bladder and bowel dysfunction, and impotence. • The use of absorbable gelatin-based hemostatic agents during the repair of dural defects associated with laminectomy and craniotomy operations, has been associated with fever, infection, leg paresthesias, neck and back pain, bladder and bowel incontinence, cauda equina syndrome, neurogenic bladder, impotence, and paresis.
---	-------------------------------	-------------	-----------	---	----------	---

4	CEREBROPROTEIN HYDROLYSATE	CEREBION-INJECTION	Injection	10 to 20 days of continuous daily usage of the required dosage of cerebroprotein hydrolysate, based on the patient's age and illness. The usual dosage is 60 to 180 mg of cerebroprotein hydrolysate. For preparing the infusion, the first required dose of cerebroprotein hydrolysate should be dissolved in 10 ml of sterile Water for Injection, which can be further diluted in 250 ml of normal saline.	990.00	Injection site reactions (pain, swelling, redness), Sweating, Headache, Dizziness, Nausea
5	DABIGATRAN	DABIGO-110mg TABLET	Tablet	110 mg BID		Constipation Belching Nausea Black or tarry stools Vomiting Cough Dizziness Loss of consciousness Fast/irregular heart beat
6	TRANEXAMIC ACID	TRANESMA-15gm GEL	Gel	2 or 3 times a day	218.00	Headache. Back pain. Muscle pain. Nose stuffiness.
7	TENECTIPLASE	TENECTASE-20mg-INJECTION(TENECTIPLASE)	Injection	Tenecteplase is for intravenous administration only. The recommended total dose should not exceed 50 mg and is based upon patient weight	28900.00	Fever Headache Nausea and Vomiting Tiredness and weakness Shortness of breath Diarrhea Reduction in blood cell count Infections
		ELAXIM-40mg(TENECTEPLASE)-INJECTION	Injection	Less than 60 kg: 30 mg IV bolus administered over 5 seconds.	43900.00	Bleeding or bruising of any kind, especially around the place of injection collection of blood under skin

ANTI-DEPRESSANTS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	DULOXETINE HCL	DUZELA-20mg CAPSULE	Capsule	40-60 mg/day PO initially (in single daily dose or divided q12hr for 1 week if patient needs to adjust to therapy)		GI upset, CNS toxicity, agitation, insomnia, raise in BP
		DUZELA-30mg TABLET	Tablet	For some patients, consider starting at 30 mg day for 1 week, to allow patients to adjust to medication before increasing to 60 mg day		
		DUVANTA-20mg TABLET	Tablet	Target dosage: 60 mg/day PO; not to exceed 120 mg/day	8.00	
2	DULOXETINE & METHYLCOBALAMIN	DUZELA-M-20mg CAPSULE	Capsule	Usually 1 or 2 times a day with or without food	8.80	Dizziness, confusion, mood swings, headache, tiredness, diarrhea, sleep changes
3	DESVENLAFAXINE EXTENDED RELEASE	D-VENIZ-50mg TABLET	Tablet	Once daily	13.00	Headache, tiredness, sleep changes, and brief feelings similar to electric shock.
4	FLUPENTIXOL & MELITRALEN	FLITRA-TABLET	Tablet	Adults: Usually 2 tablets daily: morning and noon. In severe cases the morning dose may be increased to 2 tablets. Elderly patients: 1 tablet in the morning. Maintenance dose: Usually 1 tablet in the morning. In case of insomnia and severe restlessness additional treatment with sedative in the acute phase is recommended	5.50	Restlessness, insomnia, hypomania reported, rarely dizziness, tremor, visual disturbances, headache, hyperprolactinaemia
5	TOFISOPAM	TOFICALM-50mg TABLET	Tablet	50–300 mg per day divided into three doses.	12.80	Nausea stomach discomfort fever dry mouth skin rash sleeplessness headache vomiting drowsiness

ANTI-EMETIC

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	DOMPERIDONE & NAPROXEN SODIUM TAB	NAXDOM-500mg TABLET	Tablet	Usually 2 or 3 times a day	7.20	Upset stomach, nausea, heartburn, headache, drowsiness, or dizziness may occur
		NAXDOM-250mg TABLET	Tablet			
2	ONDANSETRON	VOMIKIND MD-4mg TABLET	Tablet	4 mg orally 3 times a day		Headache Constipation Fatigue Painful urination Fever Dry mouth Anxiet
		ZOFER-30ml SYRUP	Syrup	2mg syrup every 12 hours	35.16	Headache, feeling tired or weak, hard stools (constipation)

ANTI-EPILEPTICS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	LACOSAMIDE	LACOSSET-100mg TABLET	Tablet			Feeling lightheaded, sleepy, having blurred eyesight, or a change in thinking clearly
		LACOSA-10mg/20ml- INJECTION	Injection	Infuse IV over 15-60 min; 30-60 min preferable, and should be used when a 15 min administration is not required	201.00	
2	GABAPENTIN, METHYLCOBALAMIN	GABANEURON-100mg TAB	Tablet	Dose: 300mg to 600mg about three times a day.	8.14	Dizziness, Sleepiness, Tiredness, Uncoordinated body movements.
3	LEVETIRACETAM	LEVIPIL-1000mg TABLET	Tablet	Adult: 500 mg BD - 3g OD	25.00	Somnolence, asthenia, dizziness, vertigo, depression, emotional instability, hostility,
		LEVERA-500mg TABLET		500 mg orally or IV twice a day	12.96	
		LEVERA-750mg TABLET(Levetiracetam)		750 mg orally or IV twice a day		
		LEVEPSY 500 MG TABLET		500 mg orally or IV twice a day	9.91	
4	PREGABALIN	MAXGALIN-50mg CAPSULE	Capsule	Adult: PO Adjunct in partial	8.90	Somnolence, dizziness, headache, diplopia, blurred vision, vertigo, fatigue, irritability, arthralgia
		MAXGALIN-75mg TABLET	Tablet	Seizures Initial: 150 mg/day,	13.80	

5	PREGABALIN & NORTRIPTYLINE	GABAWIN-NT TABLET	Tablet	May increase to 300 mg/day after a week. Anxiety Initial:150mg/day, may increase in increments of 150 mg weekly. Max:600mg/day.Neuropathic pain Initial: 150 mg/day, may increase to 300 mg/day after 3-7 days. Max: 600 mg/day after a 7-day interval.	15.00	Drowsiness, dizziness, dry mouth, blurred vision, constipation, weight gain, or trouble urinating may occur
		AXOGURD-NT-TABLET	Tablet	Usually 1 to 4 times daily	15.59	
6	PREGABALIN, METHYLCOBALAMIN	MAXGALIN-M-75mg TABLET	Tablet	50 mg taken three times per day Adult: 75–150 mg taken two times per day, or 50–100 mg taken three times per day (for a total of 150–300 mg per day).	14.00	Difficulty in breathing Chest tightness Chills Severe diarrhea Swelling of face, lips, eyelids, tongue, hands and feet Blurred or double vision Dizziness
7	SODIUM VALPROATE	ENCORATE-200ml SYRUP	Syrup	Dosage should start at 600 mg daily increasing by 200 mg at three day intervals until control is achieved. This is generally within the dosage range 1000 mg – 2000 mg per day,	122.21	GI upset, drowsiness, ataxia & tremor Feeling lightheaded, sleepy, having blurred eyesight, or a change in thinking clearly. Avoid driving and doing other tasks or actions that call for you to be alert or have clear eyesight until you see how this drug affects you
		VALPROL-CR-300mg TABLET	Tablet		6.35	
		VALPROL-CR-200mg TABLET	Tablet		4.23	

ANTI-FUNGALS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	FLUCONAZOLE	FZHH-150mg TABLET	Tablet	Adult: 100 - 400 mg OD Paed: 6 - 12 mg/kg/day OD	12.40	GI upset, Headache Taste perversion, HS reaction. Headache, Cutaneous rash, Anaphylaxis, Dry mouth, Alopecia, Seizure, Dizziness, Tremors.
2	PERMETHRIN	HHMITE-75gm SOAP	Soap	Twice daily	138.00	Itching, Scalp irritation. Skin irritation.

ANTI-HISTAMINE

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	CETRIZINE HCL	CETZINE-10mg TABLET	Tablet	Adult: 10 mg OD, Paed: children >6 yrs 5-10mg	1.73	Dizziness, dry mouth, GI upset, disturbed coordination, confusion

ANTI-HYPERTENSIVE

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	BENIDIPINE	BENLONG-4mg TABLET	Tablet	2-4 mg once daily, up to 8 mg once daily if needed.	7.80	Headache, facial flushing, vomiting, palpitation, dizziness, lightheadedness and malaise.
2	MOXONIDINE	MOXON-0.2mg TABLET	Tablet	Take one tablet (200 micrograms) daily, in the morning.		Dryness in mouth, Headache, Rash, Dizziness, Back pain, Insomnia (difficulty in sleeping), Sleepiness, Vomiting, Weakness, Nausea, Dyspepsia, Diarrhea, Vertigo
3	TELMISARTAN	TELMIDUCE-20mg TABLET	Tablet	Initial dose: 40 mg orally once a day	1.50	Hypotension, tachycardia, bradycardia, muscle cramps
		TELMIGET-40mg TABLET	Tablet	Maintenance dose: 40 to 80 mg orally once a day	5.75	
4	AMLODIPINE+TELMISARTAN	TELMIDUCE-AM TABLET	Tablet	once daily doses of 20 to 80 mg while amlodipine is effective in doses of 2.5 to 10 mg.		Swelling of ankles or feet Difficulty in breathing Dizziness Back pain Redness of face, neck, arms and chest Muscle pain
5	PRAZOSIN	PRAZOPRESS-XL-5mg TABLET	Tablet	0.5 mg bid or tid	11.80	During cataract surgery, Intraoperative Floppy Iris Syndrome(IFIS) may occur. Postural hypotension, syncope, tachycardia, palpitations, lack of energy, dizziness,
6	PROPRANOLOL	CIPLAR-LA-20mg TABLET*	Tablet	40 - 80 mg BID	12.60	Bradycardia, hypotension, syncope, shock, angina pectoris. Lightheadedness, giddiness, ataxia, dizziness, irritability, sleepiness, hearing loss, and visual disturbances to vivid dreams, hallucinations, and confusion.

ANTI - PARASITIC

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	IVERMECTIN	IVREA-0.5%-30ml SHAMPOO	Shampoo	Apply ivermectin shampoo only to dry hair and scalp. Use enough lotion (up to 1 tube) to completely coat and cover all of your hair from the roots to the ends and all of your scalp. Leave it on for a full 10 minutes, then rinse off with water. Children should have an adult apply this medication for them. After using this medication, wait 24 hours before applying shampoo to hair and scalp.	95.00	Facial swelling Increased heart rate Orthostatic hypotension (sudden lowering of blood pressure on standing) Peripheral edema.

ANTI-PARKINSONISM

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	LEVODOPA, CARBIDOPA, ENTACAPONE	SYNCAPONE-50mg TABLET	Tablet	usually 3 to 4 times a day. Take only one tablet for each dose	7.75	Diarrhea, nausea, vomiting, dizziness, lightheadedness, drowsiness, stomach upset, loss of appetite, trouble sleeping, headache, or dry mouth may occur. If any of these effects persist or worsen, tell your doctor or pharmacist promptly.
		SYNCOPANE-100mg TABLET	Tablet		9.00	
		SYNCAPONE-150mg TABLET	Tablet			

ANTI-PSYCHOTICS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	RISPERIDONE ORAL SOLUTION 60ML	RISDONE-60ml SYRUP	Syrup	Adult: 2.-12 mg/day	117.60	Increase in prolactin levels , postural hypotension, parkinsonism.
2	ARIPIRAZOLE	ARPIZOL-2mg TABLET	Tablet	Adult: 10 mg or 15 mg once daily	4.90	GI disorders, headache, anxiety, wt gain, insomnia, lightheadedness, drowsiness.

ANTI-ULCER

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	PANTOPRAZOLE SODIUM	PANDIFF IV-40mg INJECTION	Injection, Tablet	40 mg orally once daily 40 mg via IV infusion once a day, given over at least 2 minutes OR over 15 minutes	46.78	GI upset, headache, dry mouth confusion, drowsiness
		PANZO-40mg TABLET (PANTOPRAZOLE)			8.80	
		RBP P I.V- 40mg- INJECTION (PANTOPRAZOLE)			48.00	
		ZEPOXIN-40mg INJECTION			46.00	
2	RABEPRAZOLE	RAB ZE-20mg TABLET(RABEPRAZOLE)	Tablet	Adult:20 mg OD , 20 mg orally 2 times a day	7.03	GI upset, headache, dry mouth confusion, drowsiness
3	RABEPRAZOLE + DOMPERIDONE	RAB Z DSR-TABLET (RABEPRAZOLE+DOMPERIDONE)	Tablet	usually 1 to 2 times daily	9.30	
4	DOMPERIDONE & ESOMEPRAZOLE	SOMPRAZ-D-20mg CAPSULE	Capsules	usually 1 to 2 times daily	8.30	Nausea, bloating, or gas may occur headache, diarrhea, nausea, flatulence, stomachpain, constipation, dry mouth
5	SODIUM BICARBONATE	UDOSIS-500mg TABLET(NAHCO3)	Tablet	Parenteral: 50 to 150 mEq sodium bicarbonate diluted in 1 L of D5W to be intravenously infused at a rate of 1 to 1.5 L/hour. Oral: 325 to 2000 mg orally 1 to 4 times a day. One gram provides 11.9 mEq (mmol) each of sodium and bicarbonate.	3.00	Frequent urge to urinate headache (continuing) loss of appetite (continuing) mood or mental changes muscle pain or twitching nausea or vomiting

ANTI-VIRAL

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	SOFOBUVIR & DACLATASVIR	HEPCINAT PLUS-TABLET(28s)	Tablets	60 mg orally once a day		Fatigue Headache Nausea Insomnia (difficulty in sleeping) Anemia (low number of red blood cells)

2	GANCICLOVIR	VIRSON GEL	Gel	five times a day	111.00	Chills, headache, shortness of breath diarrhea, abdominal and stomach pain, nausea and vomiting, back pain seizures, swelling of legs, itching, decreased urine output.
3	TENOFOVIR (300MG)	TERAVIR-300mg TABLET	Tablet	300mg-once daily paed: 8mg/kg daily	43.67	Nausea, stomach pain, diarrhea, depression, headache, dizziness, weakness, trouble sleeping, itching or rash.

CNS - OTHERS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	CEREBROPROTEIN HYDROLYSATE FOR INJECTION 60MG	CEREBION-INJECTION	Injection	It is given in a dose of 60 -180 mg once daily for 10 - 20 days. It should be slowly perfused in 250 ml saline in 60 - 120 minutes. Maintenance doses (30 mg) can be given by intramuscular route. Dose of antidepressants should be reduced if used with cerebroprotein hydrolysate	990.00	Headache Nausea and vomiting Fever Confusion Hallucinations Increased sweating Diarrhea
2	PRAMIPEXOLE DIHYDROCHLORIDE	PRAMIPEX-0.25mg TABLET	Tablet	Three times a day	7.30	Nausea, dizziness, drowsiness, lightheadedness, trouble sleeping, constipation, headache, or dry mouth may occur
		PRAMIPEX-0.5mg TABLET	Tablet		11.60	
		PRAMIPEX-1mg TABLET	Tablet		16.00	
		PRAMIPEX-0.125mg TABLET	Tablet		4.70	
3	ROPINIROLE	ROPARK-0.25mg TABLET	Tablet	With or without food, usually 3 times a day	5.60	Nausea, vomiting, constipation, dizziness, drowsiness, weakness, unusual sweating, headache, and dry mouth may occur
		ROPARK-0.5mg TABLET	Tablet		8.10	
		ROPARK-1mg TABLET	Tablet		12.50	
		ROPARK-2mg TABLET	Tablet		18.00	
4	TOPIRAMATE	TOPIROL-25mg TABLET	Tablet	25-50 mg/day PO initially; increase by 25-50 mg/day at weekly intervals to 100-200 mg q12hr	7.70	Somnolence Fever
		TOPIROL-50mg TABLET	Tablet		12.20	Cognitive problems
		TOPIROL-100mg TABLET	Tablet		21.40	Infection Flushing

5	RASAGILINE	RASALECT-1mg TABLET	Tablet	0.5 mg PO qDay initially, may increase to 1 mg/day if needed and tolerated	14.00	Dizziness, drowsiness; sleep problems (insomnia), strange dreams; involuntary muscle movements; loss of appetite, weight loss; indigestion, stomach pain, nausea, vomiting, constipation; joint pain or stiffness; swelling in your hands or feet; dry mouth, cough.
		RASALECT-0.5mg TABLET	Tablet		8.00	
6	EDARAVONE	ARAVON-20ml-INJECTION	Injection	Initial treatment cycle: Administer 60 mg as an IV infusion over 60 minutes qDay for 14 days, followed by a 14-day drug-free period Subsequent treatment cycles: Administer 60 mg as an IV infusion qDay over 60 minutes on 10 of 14 days, followed by a 14-day drug-free period	580.00	Headache red, itchy, or scaly rash
7	ALPHA LIPOIC ACID ,BENFOTIAMINE, METHYLCOBALAMIN	REJUNEX CD3-TABLET	Tablet	The best dose for neuropathy is 600 mg daily. Between 50 to 100 mg is sufficient for antioxidant purposes. You can buy it in doses of 30 to 100 mg tablets	21.50	Nausea; or skin rash
8	VITAMIN D3 ORAL SOLUTION	ARACHITOL-NANO-5ml ORAL SOLUTION	Oral Solution	0.25-1microgram/day	65.12	Arachitol is oral solution (pain, swelling, redness).
9	TETRABENAZINE TABLETS	REVOCON-TABLET	Tablet	12.5 mg PO qDay initially; after 1 week, the dose should be increased to 12.5 mg q12hr	26.00	Fatigue Depression Dizziness Headache Shortness of breath Anxiety

10	BOTULINUM TOXIN -A	BOTOX- 50 IU INJECTION	Injection	BOTOX should be administered within 24 hours after reconstitution. During this time period, unused reconstituted BOTOX should be stored in a refrigerator (2° to 8°C) for up to 24 hours until time of use. BOTOX vials are for single-dose only. Discard any unused portion	12622.00	Headache Injection site pain Tiredness Neck pain Dryness in mouth
11	CEREBROLYSIN	CEREBROLYSIN-INJECTION	Injection	Cerebrolysin was administered as intravenous infusion in daily dose of 15 ml during 21 days	880.00	Loss of appetite Headache Fever Lower back pain Shortness of breath Chills
12	PYRIDOSTIGMINE	DISTINON-TABLET	Tablet	Tablets/syrup: 600 mg/day spaced to provide maximum relief	7.78	Nausea and Vomiting Diarrhea Abdominal cramps Increased salivation Weakness Muscle cramps Skin Rash Dizziness and Drowsiness
13	KHELLIN EXTRACT L-PHENYLALANINE L-TYROSINE GARLIC FICUS HISPIDA PICORRHIZA KUORRA PIPER LONGUM ZINGIBER OFFICINALE ELA CUMINUM CYMINUM CHOCORI	VITANEXT-TABLET	Tablet	The usual adult dose is 1 to 2 (30 to 120 mg Pyridostigmine) to be taken five to six times daily		Feel sick, experience vomiting, diarrhoea or stomach cramps and you may produce more saliva than usual
14	GABAPENTIN, METHYLCOBALAMIN	GABANEURON-100mg TABLET	Tablet	Three times a day	8.14	Nausea and vomiting Diarrhea Headache Skin rash Chest pain and discomfort Dizziness

CVS - OTHERS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	FRUSEMIDE + POTASSIUM CHLORIDE	DIUCONTIN-K-20mg TABLET	Tablet	The usual dose is 2 to 3 tablets a day.	4.28	High blood levels of potassium (which can cause an abnormal heartbeat) itching and skin rashes (nettle rash).
2	TADALAFIL	TAZZLE-20mg TABLET	Tablet	20 mg once daily	53.80	Headache, flushing, angina pectoris, respiratory tract infection, AV block, migraine, syncope, tachycardia, postural hypotension, MI, cerebral thrombosis

HORMONES & THEIR ANTAGONIST

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	MENOTROPHIN	HUMOG-150IU-INJECTION	Injection	Three equal doses of HUMOG HP are given on alternate days	1272.00	In female, a local reaction at the injection site, fever and arthralgia have been observed in rare cases. In the male, a combined treatment with HUMOG HP
2	COENZYME Q10, DEHYDROEPIANDROSTERONE (MICRONIZED) & MELATONIN	OVARES PLUS-TABLET	Tablet	Heart failure Adult: 50-150 mg daily in 2-3 divided doses. Angina Adult: 150-600 mg daily in divided doses. Periodontal disease Adult: 25 mg bid.	47.00	Drowsiness, change in sleep pattern and headache
3	DYDROGESTERONE TAB	DUPHASTAN-10mg TABLET	Tablet	Duphaston 10 mg tablets once or twice daily with or without food	62.35	Depression Mood swings Difficulty in falling asleep Headache Dizziness Drowsiness
4	TESTOSTERONE ENANTHATE	TESTOVIRON-250mg INJECTION	Injection	Usually every 1 to 4 weeks	311.60	Nausea, vomiting, headache, skin color changes, increased/decreased sexual interest, oily skin, hair loss, and acne may occur. Pain and redness at the injection site may also occur

5	FSH	SITRODIN-75 IU INJECTION(FSH 75 IU)	Injection	The recommended dose is 150 IU/day for 5 days, increased up to 450 IU/day for up to 12 days.	1044.90	Headache, nausea, vomiting, mild stomach/abdominal pain, bloating, redness/pain at the injection site, or breast tenderness/pain may occur
6	DEXAMETHASONE	AURADEx-8mg TABLET (DEXAMETHASONE)	Tablet	0.75-9 mg/day in 2-4 divided doses	6.00	Peptic ulceration, acne eruptions, glaucoma, osteoporosis
7	DINOPROSTONE	DINOFIRST-2mg TABLET	Tablet	500 mcg, may repeat 6 hr later if needed. Max: 1.5 mg/24 hr.	54.79	Hypersensitivity reactions, nausea, vomiting, diarrhoea, abdominal pain; flushing, shivering, headache, dizziness; hypertension; convulsions, ECG changes
8	FLUNARIZINE	MIGRANEX-10mg TABLET	Tablet	5-10 mg/day at bedtime	4.97	Drowsiness. Rarely wt gain, headache, depression, gastric pain, dry mouth, insomnia, extrapyramidal reactions, galactorrhoea.
	DIHYDROCHLORIDE					
9	MEDROXY PROGESTERONE ACETATE	EMILY IUS	Single insertion	20 µg per day	2664.00	Depression, fluid retention. Fatigue, insomnia, dizziness, headache, nausea; breast tenderness; wt gain/loss, anorexia; cholestatic jaundice; pain at Inj site.
10	MISOPROSTOL	MISOPROST*-25mcg TABLET	Tablet	200-800 mcg/day in 2-4 divided doses	18.30	Diarrhoea, abdominal pain, dyspepsia, constipation, flatulence, nausea and vomiting; abnormal vaginal bleeding, cramps, increased uterine contractility, headache.
11	OXYTOCIN	OXYTOCIN-1ml INJECTION	Injection	10-40 u by infusion at a rate to control uterine atony	17.79	Transient hypotension, reflex tachycardia; nasal irritation, rhinorrhoea, lachrymation
12	METHYLPREDNISOLONE ACETATE	DEPOPRED-2ml INJECTION	Injection	The range of initial doses is 0.11 to 1.6 mg/kg/day	67.12	Aggression or anger Decreased urine output Dizziness Headache

MUSCLE RELAXANTS, ANAESTHETIC & EMERGENCY

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	PROPOFOL	TROYPOFOL 10ML INJECTION	Injection	Adult: 1.5 - 2.5 mg/kg IV upto 15 mg/kg/hr Paed: 3 - 3.5 mg/kg upto 12.5 mg/kg/hr	98.00	Nausea, vomiting, hiccough, ophthalmoplegia, anaphylaxis
2	VECURONIUM BROMIDE	KROVEC-4mg-INJECTION	Injection	Adult: 80 - 100 mcg/kg maintenance: 20 -30 mcg/kg Paed: < 4mth : 10-20mcg/kg > 4 mth: as in adults	135.00	Muscle weakness, paralysis, muscleatrophy (after long term use), hypersensitivity reactions e.g urticaria and erythema.
3	ROPIVACINE	ROPEE-7.5mg-20ml-INJ	Injection	75-200 mg (15-30 mL of 0.5%-1% solution)	173.90	Nausea, vomiting; headache, back pain; fever; itching; numbness or tingly feeling
4	THIOLCHOLCHICOSIDE + ACECLOFENAC	SPERONAC-MR-TABLET	Tablet	Muscle spasms Adult: Initially, 16 mg daily. Intramuscular Muscle spasms. Adult: Up to 8 mg daily.		Photosensitivity reactions.
5	BACLOFEN EXTENDED RELEASE	LIOFEN-XL-20mg TABLET	Tablet	5 mg 3 times/day for 3 days, increased to 10 mg 3 times /day for 3 days	25.00	Sedation, drowsiness, ataxia, dizziness, headache, confusion, hallucinations, skin reactions, GI symptoms, enuresis.
		LIOFEN-XL-30mg TABLET(Baclofen)				
6	IBUPROFEN, PARACETAMOL &CHLORZOXAZONE	FLEXON-MR-TABLET	Tablet	The recommended initial dose is 500 mg 3-4 timesday. May reduce to 250 mg 3-4 timesday subsequently. Max: 750 mg 3-4 timesday	1.84	Dry mouth, dizziness, drowsiness, lightheadedness, blurred vision, nausea, vomiting, and constipation.
7	DROTAVERINE HYDROCHLORIDE & PARACETAMOL	DROTIN PLUS-TABLET	Tablet	40-80 mg 3 timesday.	12.61	Difficulty in breathing, Allergic skin reaction, Swelling of face, lips, eyelids, tongue, hands and feet, Fall in blood pressure.

8	FENTANYL CITRATE	Verfen	Injection	Initial: 50-200 mcg followed by supplements of 50 mcg. Max: 200 mcg. Admin lower infusion rates of 0.05-0.08 mcg /kg/min. Patients w/ assisted ventilation: Initial: 300-3,500 mcg (up to 50 mcg/kg) followed by supplements of 100-200 mcg depending on response. Loading dose (alternatively via bolus): Approx 1 mcg/kg/min given for the 1st 10 min followed by infusion of approx 100 ng/kg/min	46.00	
9	BUPRENORPHINE	MUTRANS-10mcg-PATCHES (BUPRENORPHINE)	Patches	IM Anaesth premed 300 mcg. IV Perioperative analgesia 300-450 mcg. IM/IV Moderate to severe pain 300-600 mcg 6-8 hrly.	995.00	Nausea, vomiting, dizziness, sweating and drowsiness.

NSAIDS & GOUT

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ACECLOFENAC, PARACETAMOL WITH SERRATIOPEPTIDASE	PAIN OFF SP-TABLET	Tablet	The recommended dose is one tablet to be taken twice daily	9.00	Dizziness •nausea (feeling sick) •diarrhoea •increased liver enzymes in the blood
2	PARACETAMOL + ACECLOFENAC	PARAFENAC Z-TABLET(ACECLOFENAC&PARACETAMOL)	Tablets	1 tablet should be taken 1 - 2 times daily	4.50	Nausea, Vomiting, Perforation, GI ulceration, Bleeding, Bronchospasm, Coagulopathy

3	RIZATRIPTAN BENZOATE ORALLY DISINTEGRATING	RIZANET-10MG STRIP	Tablet	Rizatriptan dosage is 5 mg single dose in patients weighing less than 40 Kg and 10 mg single dose in patients weighing 40 kg or more	53.00	Chest pain Drowsiness Dizziness Shortness of breath Feeling of warmth or heat Irregular heartbeat Itching Numbness Dry mouth Nausea and Vomiting Unusual tiredness and weakness
		RIZANET-5MG STRIP			36.00	
4	TRANEXAMIC ACID-500MG+MEFENAMIC ACID-250MG	TEXAKIND-MF TABLET	Tablet	The recommended standard dose is 2 to 3 tablets taken 2 to 3 times daily.	17.30	Nausea, vomiting, diarrhoea
5	DICLOFENAC SODIUM	VOLITRA ENZO-TABLET	Tablet	50 mg orally 2 or 3 times a day or 75 mg orally 2 times a day	15.50	It may cause dizziness and blurred vision
6	ALCOHOL 10 %V/V+DICLOFENAC 4 %W/V	VOLITRA APS SPRAY-30ml	Spray	100-200 mg/day in divided doses	235.00	Itching, Headache, Drowsiness
7	DICLOFENAC (50 MG) + TOLPERISONE (150 MG)	TOLPERITAS-D-TABLET	Tablet	100-200 mg/day in divided doses.		Abdominal pain, Diarrhea, Nausea and Vomiting, Skin Rash, Dry mouth

SEDATIVES AND HYPNOTICS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	CLONAZEPAM	ZAPIZ-0.25mg TABLET	Tablet	Adult: 1.5 mg/day Paed: 0.01 - 0.03 mg/kg/day		Anorexia, loss of memory & coordination, tachycardia, hypotension
2	ZOLPIDEM	SOBRIUM-5mg TABLET	Tablet	Women: 5 mg orally once a day immediately prior to bedtime -Men: 5 to 10 mg orally once a day immediately prior to bedtime Maintenance dose: 5 to 10 mg orally once a day immediately prior to bedtime Maximum dose: 10 mg/day	5.42	Unsteadiness Confusion Dizziness Fast heartbeat Drowsiness Blurred vision Dry mouth Headache Loss of appetite Nausea or Vomiting Xia, loss of memory & coordination, tachycardia, hypotension

VACCINE

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	BACTERIUM SALMONELLA TYPHI	BIOTYPH-2.5ml-INJECTION	Injection	0.5 ML	650.00	Headache, fever, rash, itching, abdominal pain, nausea, diarrhea, vomiting, anaphylactic reaction, nerve damage, muscle pain, hives (area of a skin being raised indicating an allergic reaction), sepsis and weakness
2	DIPHTHERIA, TETANUS, PERTUSSIS (ACELLULAR, COMPONENT), HEPATITIS B (RDNA), POLIOMYELITIS (INACTIVATED) AND HAEMOPHILUS INFLUENZAE TYPE B CONJUGATE VACCINE (ADSORBED)	HEXAXIM-INJECTION	Injection	0.5 ML	3900.00	Injection-site pain, injection-site erythema, injection-site swelling Irritability Pyrexia (body temperature $\geq 38.0^{\circ}\text{C}$)
3	VARICELLA-ZOSTER	NEXIPOX-INJECTION	Injection	Each 0.5ml vial/dose	1779.00	Redness, itching, tenderness, swelling, bruising, or a lump where the shot was given; low fever; or. mild skin rash that looks like chickenpox (can occur up to 1 month after vaccination).

MISCELLANEOUS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	METHOXSALEN TOPICAL SOULUTION	OCTAMOP-30ml LOTION	Lotion	It is applied on the skin along with ultraviolet-A irradiation	118.00	Chewing sugar-free gum may help. Sunburn. Skin irritation. Itching.

2	TACROLIMUS	TACROVERA-0.1%-15ml LOTION	Lotion	Apply a thin layer over the affected skin twice daily.	530.00	High blood pressure. Kidney function that gets worse. Chance of getting an infection. Wash hands often. Stay away from people with infections, colds, or flu. Feeling dizzy. Rise slowly over a few minutes when sitting or lying down. Be careful climbing. Headache. Upset stomach
3	CALCIPOTRIOL	PASITREX-20gm OINMENT	Ointment	Apply 1-2 timesday	330.00	Skin Irritation
4	FLUOCINOLONE ACETONIDE+HYDROQUINONE+TRETINOIN	TRILUMA-15gm CREAM	Cream	Apply 3-4 timesday	495.00	Skin irritation.
5	FERROUS ASCORBATE & FOLIC ACID	OROFER-XT-TABLET	Tablet	The most recommended dosage in adults is one-two tablets/ tablespoon right after a meal.	13.00	Vomiting, Nausea, Altered sleep patterns, Constipation, Flatulence, Anorexia
6	DICALCIUM MALATE, GLYCINE, VITAMIN D3	DICAL-D TABLET	Tablet	Once daily	10.50	Rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat
7	ALPHA LIPOIC ACID, BENFOTIAMINE,CHROMIUM PICOLINATE,FOLIC ACID,INOSITOL,MECOBALAMIN	BENFOMET FORTE TABLET	Tablet	100 mg 4 timesday.	22.50	Allergic reactions, Heart disease, Low blood sugar levels, Thyroid disorder, History of stomach ulcer, Liver disease, Hypersensitivity, Kidney disease, Asthma or any breathing disorder
8	NATIVE TYPE 11 COLLAGEN CAPSULES	GERIJOINT-40mg-CAPSULE	Capsule	2.5 mg daily for up to 24 weeks	41.00	Upset stomach or throwing up. Constipation. Change in color of stool to green. Diarrhea. Belly pain.
9	VITAMIN D3 ORAL SOLUTION	ARACHITOL-NANO-5ml ORAL SOLUTION	Oral Solution	Adults, elderly and children over 12 years of age – Swallow 2 twice daily, ideally two in the morning and two in the evening. Children under 12 – Do not give Vitamin D3 to children under 12 years	65.12	Constipation, wind, feeling sick, stomach ache, diarrhoea •Skin rash •Hypercalcaemia

10	CISSUS QUADRANGULARIS, MORINGA OLEIFERA AND VITAMIN D3	CISBIND-TABLET	Tablet	once a day in the morning with or without food	15.00	Dry mouth, muscle pain, increased thirst, loss of appetite, vomiting, metallic taste, upset stomach, constipation, difficult urination.
11	OMEGA- 3 FATTY ACID, GREEN TEA EXTRACT GINKGO BILOBA, GINSENG, GRAPE SEED EXTRACT, VITAMINS, MINERALS AND TRACE ELEMENTS	FITMUNE-TABLET	Tablet	once a day in the morning	16.00	Drowsiness, dizziness, constipation, stomach upset, blurred vision, or dry mouth/nose/throat
12	CALCIUM AND VITAMIN D3	SHELCAL-HD-TABLET	Tablet	once a day in the morning	6.27	Nausea, vomiting, constipation; increased thirst or urination; muscle weakness, bone pain
13	GINKGO TABLETS	BILOVAS-TABLET(GINKGO)	Tablet	once a day in the morning	19.78	Allergic skin reactions. Bleeding disorders. Constipation. Contact dermatitis
14	UNDENATURATED TYPE-II COLLAGEN, CALCIUM, CHOLECALCIFEROL, VITAMIN C	OSTOSHINE-TABLET	Tablet	once a day in the morning	70.00	Rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat

15	WATER, CETEARYL ALCOHOL, ETHANOLAMINE, STEARTRIMONIUM CHLORIDE, STEARAMIDE DEA OTHER INGREDIENTS : CETEARETH-20, TOLUENE-2, 5 DIAMINE SULFATE, STEARETH-21, STEARAMIDOPROPYL DIMETHYLAMINE, DIMETHICONE, AMODIMETHICONE, BOSEWELLIA SERRATA EXTRACT, MINERAL OIL, RESORCINOL, OLETH-2, LANOLIN, PORTULACA OLERACEA EXTRACT, BEHENTRIMONIUM CHLORIDE, GLYCERYL STEARATE SE, SORBITAN SESQUIOLEATE, SODIUM SULFITE, M-AMINOPHENOL, CYSTEIN HCL, ALLANTOIN, O-AMINOPHENOL, TETRASODIUM EDTA, POLYQUATERNIUM-10, PANTHENOL, ASCORBIC ACID, FRAGRANCE, OLEA EUROPEA (OLIVE) FRUIT OIL	CUTICOLOR HAIR COLOUR	Hair colour	Once in 15 days	1433.50	Causing allergy
16	EPALRESTAT, METHYLCOBALAMIN	EPALRICA-M-TABLET	Tablet	3 times/day		Constipation, diarrhea, or upset stomach
17	CALCITRIOL +CALCIUM CITRATE	OSTEOFIT-C-TABLET		Initial: 0.25 mcg/day		Constipation, wind, feeling sick, stomach ache, diarrhoea.
18	CALCIUM FOLINATE	NEOVORIN-50mg/5ml-INJECTION (CALCIUM FOLINATE)	Injection	200 mg/ml by intravenous infusion over two hours, followed by bolus 400 mg/ml	81.00	Constipation, oral inflammation or cold sores, fatigue, hair loss, or loss of appetite

19	L-METHYLFOLATE, METHYLCOBALAMIN, PYRIDOXAL 5-PHOSPHATE	FOLVITE-ACTIVE-CAPSULE	Capsule	one tablet daily	12.10	Rash, itching/swelling (especially of the face/tongue/throat), dizziness, or trouble breathing
20	LACTULOSE SOLUTION	LOOZE-200ml LIQUID	Liquid	Adult: PO Constipation Initial: 10-20 g (15-30 mL)/day Max: 45 mL (or 40g of the reconstituted oral formulation)/day.	216.30	Diarrhoea, Nausea, Vomiting
21	ALUMINIUM HYDROXIDE 300 MG+DIMETHICONE 40 MG+MAGNESIUM HYDROXIDE 250 MG+Sorbitol 1000 MG /5ML	DIOVOL-170ml SYRUP	Syrup	Take 1 to 3 hours after meals as an antacid	95.00	Bad taste in your mouth. Hard stools
22	LACTITOL	LACTIHEP PLUS-250ml SYRUP	Syrup	2 tablespoon 8-15ml during night	211.00	Dehydration
23	ALPHA KETOANALOUGE	KETOALPHA-TABLET	Tablet	4 to 8 tablets three times daily during meals	34.00	Hypercalcemia
24	PRUCALOPRIDE TABLETS	PRUEASE-1mg TABLET	Tablet	Take one tablet daily.	16.35	Headache Nausea
		PRUEASE-2 mg TABLET	Tablet	2 milligrams per day.	22.90	Diarrhea Abdominal pain
25	OMEGA- 3 FATTY ACID, GREEN TEA EXTRACT GINKGO BILOBA, GINSENG, GRAPE SEED EXTRACT, VITAMINS, MINERALS AND TRACE ELEMENTS TABLET	FITMUNE-TABLET	Tablet	three times a day	16.00	Drowsiness, dizziness, constipation, stomach upset, blurred vision, or dry mouth/nose/throat may occur
26	COENZYME Q10, DEHYDROEPIANDROSTERONE (MICRONIZED) & MELATONIN	OVARES PLUS-TABLET		500 mcg of selenium daily.	47.00	Drowsiness, change in sleep pattern and headache
27	BIOTIN-10MG+CALCIUM PANTOTHENATE-100MG+COPPER-3MG+ZINC-22.5MG+ACETYLCYSTEINE-50MG+SELENIUM-65MCG	BTN-ULTRA TABLET	Tablet	4 to 7 milligrams (mg) per day		Constipation, diarrhea, or upset stomach
28	CLOBETASOL-0.05% W/W+SALICYLIC ACID-3% W/W	CLOP-S-20gm LOTION	Lotion	You usually put Clobetasol propionate on your scalp in the morning and at night	191.40	Skin peeling Skin thinning Application site reactions (burning, irritation, itching and redness)

29	MOMETASONE FUROATE 0.1 %W/V	HH SONE-10ml LOTION	Lotion	Apply affected area one or two times a day	158.00	burning –stinging –itching –tingling
30	LIQUID PARAFFIN-10.2% W/W+WHITE SOFT PARAFFIN-13.2% W/W	COTARYL-50gm CREAM	Cream	Apply to affected skin two or three times a day	76.00	Rash Itching Swelling at applied sites
31	BENZOYL PEROXIDE	BENZAC-AC 5%-20gm GEL	Gel	Apply to affected skin two or three times a day	166.90	Skin reactions such as peeling, itching, irritation, and reddened skin may occur, especially at the start of treatment
32	SAFFLOWER OIL 362.2 MG, FISH OIL 87.7 MG, BORAGE OIL 51.0 MG, ASCORBIC ACID 50.0 MG, ZINC SULPHATE 2.0 MG, VITAMIN E 10.0 IU (AS DL-ALPHA TOCOPHERYL ACETATE). OMEGA 3: EICOSAPENTAENOIC ACID (EPA) 15.8 MG, DOCOSAHEXAENOIC ACID (DHA) 10.5 MG. OMEGA 6: GAMMA LINOLENIC ACID (GLA) 10.2 MG, LINOLENIC ACID (LA) 280.8 MG.	EICOSA-CAPSULE	Capsule	One capsule daily	11.00	Digestive tract, liver or kidneys.
33	HALOBETASOL PROPIONATE	HALONOVA-15gm CREAM	Cream	Topical- Apply a thin layer to the affected skin once or twice daily	129.80	Dry skin, redness, skin scar, white patches, vesicles and rash
34	BETACAROTENE- 10MG+ZINC SULFATE- 27.4MG+SELENIUM DIOXIDE- 0.07MG+COPPER- 1MG+MANGANESE-2MG	ANTOXID-CAPSULE	Capsule	1 hour before or 2 hours after meals. May be taken with food if it upsets your stomach		Nausea, stomach upset, heartburn
35	TENDOCARE TABLET CONTAINS COLLAGEN PEPTIDE(40MG), CHONDROITIN SULFATE (200MG), SODIUM HYALURONATE (30MG) AND VITAMIN C (35MG)	TENDOCARE-TABLET	Tablet	1 tablet once or twice per day	20.33	Nausea or vomiting Headache Bloating Nausea Gastrointestinal symptoms Wheezing

36	COLLAGEN PEPTIDE-TYPE II	REJOINT-UC CAPSULE	Capsule	1 capsule a day	61.00	Nausea, Heartburn, Diarrhea, constipation, Drowsiness Headache, Allergic Skin Reaction
37	TOCOTRIENOL	TOCOFIL-CAPSULE	Capsule	1 capsule a day		Feeling tired or weak (fatigue), stomach pain, loose stools (diarrhea), headache, nausea, or blurred vision
38	SOY PROTEIN	SOYGROWTH-200gm POWDER	POWDER	1 table spoon with water	465.00	Constipation, nausea Bloating, rash, itching Abnormal tissue growth in uterus
39	MECOBALAMIN, ZINC SULPHATE, VITAMIN B6, FOLIC ACID, NIACINAMIDE, CHROMIUM PICOLINATE	NUROSTAR-TABLET	Tablet	1 capsule a day	4.62	Skin irritation, Headache, Dizziness Nausea, Mood changes
40	GLYCERIN HYDROGENATED POLYISOBUTENE CETEARYL ALCOHOL CETEARETH-20 MACADAMIA NUT OIL DIMETHICONE TOCOPHERYL ACETATE STEAROXYTRIMETHYLSILANE (AND) STEARYL ALCOHOL PANTHENOL FARNESOL BENZYL ALCOHOL PHENOXYETHANOL	CETAPHIL DAM-100gm LOTION	Lotion	Recommended for gentle daily cleansing of damaged skin associated with common skin conditions and following dermatological treatments	373.00	Allergic reaction: hives; difficult breathing; swelling of your face, lips, tongue, or throat.
41	ALMOND, ESSENTIAL OIL AND VITAMIN E	KERA-XL HAIR GROWTH-60ml	Hair Growth Serum		1395.00	Abdominal cramps. Photosensitivity. Rash. Burning. Excess of creatinine in urine
42	FOLIC ACID 5 MG+METHYLCOBALAMIN 1500 MCG+PYRIDOXINE 20 MG	HOMISURE-TABLET	Tablet	50mg three times a day		Wheezing; chest tightness; fever; itching; bad cough; blue or gray skin color; seizures; or swelling of face, lips, tongue, or throat.

43	PHENOXYETHANOL, HYDROXYPROPYL OXIDIZED STARCH PG- TRIMONIUM CHLORIDE, SODIUM BENZOATE, CITRIC ACID MONOHYDRATE, SODIUM CITRATE, DECYL GLUCOSIDE, DISODIUM LAURETH SULFOSUCCINATE, SODIUM LAURYL SULFOACETATE	FOLLIGLO-60gm SHAMP00	Shampoo	Wet hair and apply shampoo on scalp evenly - Keep shampoo on the scalp for 5 minutes before rinsing	185.00	Mild skin itching or irritation, dry skin, abnormal hair texture, scalp pustules,
44	MINOXIDIL	ZENOXIDIL ROLL ON-10ml	Roll on	once in a day	110.00	Feeling dizzy. Rise slowly over a few minutes when sitting or lying down. Be careful climbing. Hair growth
45	AZATHIOPRINE	AZR-50mg TABLET	Tablet	3 to 5 mg/kg orally once a day	10.43	Upset stomach
46	EPALRESTAT- 150MG+METHYLCOBALAMI N-1500MCG	EPALRICA-M-TABLET (EPALRESTAT+MC)	Tablet	50mg 3 times/day,	20.93	Constipation, diarrhea, or upset stomach may occur
47	MYCOPHENOLATE MOFETIL	MYCHOPEN-500mg TABLET(mycophenolate mofetil)	Tablet	Twice daily on an empty stomach, 1 hour before or 2 hours after meals.	78.26	Constipation,nausea,headace,diarrhea, vomiting,stomach upset,gas,tremor
48	ANHYDROUS DEXTROSE 15 GM+CALCIUM CHLORIDE 0.257 GM+MAGNESIUM CHLORIDE 0.0508 GM+SODIUM CHLORIDE 5.38 GM+SODIUM LACTATE 4.48 GM /1000ML	PERITONEAL DIALYSIS SOLUTION-2000ml	SOLUTION	Approximately two liters of dialysis solution are instilled into the peritoneal cavity of adults	235.00	Abdominal pain, bleeding, peritonitis, subcutaneous infection