

Sri  
**MANAKULA VINAYAGAR**




Medical College and Hospital

Kalitheerthalkuppam, Madagadipet, Puducherry – 605107.

Department of Pharmacy

# DRUG FORMULARY

Edition - III



## PREFACE

---

This Drug Formulary of Sri Manakula Vinayagar Medical College and Hospital is a concise and handy book designed for quick reference of current essential drugs in medical practice. As it is not possible to include all drugs, commonly used drugs in our hospital with their recommended dosages, both in adults and children, adverse drug reaction and cost have been included. However, as the medical knowledge is vast and continuously being upgraded, changes may have occurred and may not be reflected in the incorporated information. Clinical judgment and experience should be used while prescribing the drugs.

**Note:** Dose of drug varies with indications and severity of disease, the given doses is just for reference, and is for the most common indication. Where feasible, the dose ranges from lowest to maximal permissible doses have been given. Paediatric doses are for children more than 1 month of age i.e. neonatal doses may vary. The price given is for an estimation of the cost of therapy and not the exact selling price, which may vary.

“Let food be thy medicine and medicine be thy food.”

**-Hippocrates**

## CONTENTS

SL. NO.	SECTION	PAGE NO.
1	ANTI-ASTHMA, DECONGESTANTS , OTHER RS	1
2	ANTI-BACTERIALS	4
3	ANTI-CANCER	11
4	ANTI-COAGULANTS AND THROMBOLYTIC	17
5	ANTI-DEPRESSANTS	20
6	ANTI-DIABETIC	21
7	ANTI-DIURETICS AND ITS ANALOGUES	128
8	ANTI-EMETIC	25
9	ANTI-EPILEPTICS	26
10	ANTI-FUNGALS	29
11	ANTI-HISTAMINE	31
12	ANTI-HYPERTENSIVE & HYPOTENSIVE	32
13	ANTI-MALARIAL	37
14	ANTI-PARASITIC	37
15	ANTI-PARKINSONISM	38
16	ANTI-PSYCHOTICS	38
17	ANTI-TUSSIVES & EXPECTORANTS	40
18	ANTI-ULCER	42
19	ANTI-VIRAL	45
20	CNS - OTHERS	46
21	CVS - OTHERS	49
22	DERMATOLOGICAL (EXTERNAL PREPARATION)	99
23	EYE,EAR,NOSE, MOUTH & THROAT	95
24	HORMONES & THEIR ANTAGONIST	53
25	HYPOLIPIDEMIC DRUGS	60
26	IMMUNOMODULATORS	62

27	MULTIVITAMINS, MINERALS AND IRON	111
28	MUSCLE RELAXANTS, ANAESTHETIC & EMERGENCY	64
29	NSAIDS & GOUT	69
30	SEDATIVES AND HYPNOTICS	73
31	VACCINE	75
32	MISCELLANEOUS	76

**“Writing prescriptions is easy, understanding people is  
hard”**

**- Franz kafker, 1883-1924**

## ANTI-ASTHMA, DECONGESTANTS , OTHER RS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ACEBROPHYLLIN	AB-PHYLLINE - 100 MG	Tablets, Capsules, Syrup	100 mg twice daily Children up to 2 years, it`s half a teaspoon, twice daily. Age 2 to 5 is half a teaspoon, three times a day	81.50	Restlessness, Tremor, Palpitation, Tachypnoea,
		AB-PHYLLINE SR - 200 MG			141.00	
2	ACETYLCYSTEINE	MUCOMIX - 2ml	Respules, injection,	PO As a mucolytic As loz, granules or effervescent tab: 600 mg daily as a single or in 3 divided doses. Paracetamol poisoning As 5% soln: Initial:140 mg/kg followed by 70mg/kg 4 hrly for additional 17	49.95	Bronchospasm, angioedema, rash, pruritus, hypotension, HTN, flushing, nausea, vomiting.
		MUCOMIX	Tablet, Capsule		320.00	
		MUCINAC - 600mg			179.00	
3	ACEBROPHYLLIN & N-ACETYLCYSTEINE	AB-PHYLLINE-N-TABLET	Tablet	Acetabrophylline 100 mg+N-Acetylcysteine 600 mg OD	149.00	Restlessness, Tremor, Palpitation, Tachypnoea, Flushing, Hypotension .
4	AMBRISENTAN	PULMONEXT-5mg	Tablet	Once daily	1450.00	Nasal congestion, sinusitis, and flushing.
5	AMINOPHYLLINE	AMINOPHYLLINE. I.P. - 10 ML	Injection	Adult: Loading dose IV: 6.3 mg/kg over 30 min Maintenance dose : 0.3 - 0.8 mg/kg/hr Paed dose: Maintenance dose IV 0.8 mg/kg/hr.	20.00	GI upset, Rapid IV causes precordial pain, syncope & even sudden death.
7	AZELASTINE HYDROCHLORIDE & FLUTICASONE FUROATE	AZEFLO-FT NASAL SPRAY	Nasal Spray	Nasal Rhinitis 140 mcg into each nostril twice daily.	395.00	Irritation, stinging & itching of the nasal mucosa. Sneezing, nosebleeds, headache; nausea, taste disturbances, somnolence,dry mouth .
8	AZELASTINE HYDROCHLORIDE & FLUTICASONE PROPIONATE	DUONASE NASAL SPRAY	Nasal spray	Nasal Rhinitis 140 mcg into each nostril twice daily.	264.00	Irritation, stinging & itching of the nasal mucosa. Sneezing, nosebleeds, headache; nausea, taste disturbances, somnolence, dry mouth.
10	BECLOMETASONE & FORMOTEROL	FULL FORM-200 R/C	Rotacaps	2 spray 2 times a day.	157.30	Back pain,Pneumonia Bronchitis (inflammation of the airways),Sinus inflammation,Cough Oropharyngeal pain,Joint pain.
12	BILASTINE	BILAGRA-20mg	Tablet	Once daily.	120.00	Headache and drowsiness.
13	BILASTINE & MONTELUKAST	MONTEK-BL	Tablet	OD at bed time	175.00	Cough Upper respiratory infection Vomiting Headache Diarrhea Sinus infection Abdominal pain.
14	BUDESONIDE	BUDECORT - 200	Inhaler,	Budesonide: 100/200 mcg	396.00	Dryness, Irritation, Smell & taste disturbance.
		BUDECORT - 100 mg	Rotacaps,Respules		67.00	
		BUDECORT - 200 mg			86.00	
		BUDECORT - 0.5 mg			30.00	

15	DOXOFYLLINE	DOXOLIN - 400 mg	Tablet	Adult: Up to 1200 mg daily.	54.30	Restlessness, tremor, palpitation, tachypnoea, flushing, hypotension
		DOXOLIN-200mg		The recommended dose is up to 1,200 mg/day.	66.00	
16	EPHEDRINE	EFIPRES 1ml	Injection	As ephedrine sulfate: 12.5-25 mg.	23.40	Tachycardia, anxiety, nausea, restlessness, insomnia, tremor, dry mouth, impaired circulation to the HTN, headache, cardiac arrhythmia.
17	ETOPHYLLINE & THEOPHYLLINE	DERIPHYLLIN RETARD - 150 mg	Tablet, Injection	Eto: 115 mg	5.80	Restlessness, tremor, palpitation, tachypnoea, flushing, hypotension.
		DERIPHYLLIN - 100 mg		Theo: 35 mg	3.55	
		DERIPHYLLIN			4.35	
18	FLUTICASONE FUROATE & VILANTEROL	BESVIL-100 OCTACAPS	Octacaps	Once daily	445.00	Headache, Runny nose, Sore Throat (Pharyngitis), Joint pain, Nervousness.
19	FLUTICASONE PROPIONATE	FLUTICONE - 12 ml	Nasal spray, Inhaler, Capsule	100 mcg bd	251.00	Topical: Pruritus, hypertrichosis, dryness, numbness of fingers, burning, eruptions, hypopigmentation.
		FLUTICONE-OX			330.00	
		FLUTIFLO			189.00	
		NEZAFLO		The usual starting dose is 2 sprays into each nostril once a day, preferably in the morning.	262.00	
20	FLUTICASONE FUROATE	ENHHALE	Inhaler, Rotacaps, Respules	once daily	275.00	allergic contact dermatitis, secondary infection, skin atrophy
		FURAMIST		290.00		
		FLUTIFLO-FT		263.50		
21	FLUTICASONE FUROATE & OXYMETAZOLINE	FLUTICONE-OX	Nasal Spray	1 spray in each nostril 2 times a day.	300.00	Headache, Dry nose, Fever
22	FLUTICASONE AZELASTINE	FURAMIST-AZ	Nasal spray, Respules		326.50	
23	FLUTICASONE FORMOTEROL	MAXIFLO - 250	Inhaler, Rotacaps		244.00	
		MAXIFLO - 100		152.00		
		MAXIFLO - 250		490.00		
24	FORMOTEROL FUMARATE & BUDESONIDE	FORACORT 200	Inhaler, Rotacaps, Respules.	Formoterol: 6mcg	175.00	Nasal irritation, sneezing, itching of throat & dryness.
		FORACORT - 0.5 mg-2ml		Budesonide: 100/200 mcg	39.00	
		FORACORT - 200 mg			323.00	
		FORACORT - 400 mg			426.00	
		FORACORT 1MG-2 ml			44.00	
		FORMONIDE - 200 mg			333.00	

		FORMONIDE - 100 mcg			161.00	
		FORMOFLO -100			143.50	
		FORMONIDE - 200 mg			333.00	
		BUDAMATE - 200			322.50	
		BUDAMATE - 200			159.00	
		BUDAMATE - 400			408.50	
		BUDAMATE -400			205.50	
		FORACORT-100mcg			235.99	
		FORMOFLO-250mg			598.65	
		FORMONIDE-0.5mg/2ml			43.90	
		FOMTIDE-200mg			287.43	
25	FORMOTEROL FUMARATE & GLYCOPYRRONIUM	FORGLYN	Inhaler	Two times a day.	544.50	Headache, Dizziness, Dry mouth,Nausea, Painful and frequent urination, Chest pain, Muscle cramps, Anxiety.
27	FORMOTEROL FUMARATE, BUDESONIDE & IPRATROPIUM BROMIDE.	TRIMIUM TRANSHALER	Inhaler	Formoterol: 6mcg Budesonide: 100/200 mcg.	751.00	Nasal irritation, sneezing, itching of throat& dryness.
28	GLYCOPYRRONIUM BROMIDE / GLYCOPYRROLATE(9.0 MCG) + FORMOTEROL FUMARATE(4.8 MCG) + BUDESONIDE(160.0 MCG)	FORGLYN PLUS	Inhaler	Twice a day	823.9	Indigestion. High blood pressure. Nasopharyngitis (inflammation of the throat and nasal passages) Chest pain. Fever.Urinary retention.Sore throat.
29	GLYCOPYRRONIUM BROMIDE	NEBZMART-G-2ml	Capsules, Respules	Inhalation is one capsule once daily using the inhaler.	39.40	Constipation, Dry mouth.
35	IPRATROPIUM BROMIDE	IPRAVENT (RESP)SOLUTION 15M	Inhaler, Respules	Adult: Nebulized 500 mcg 3-4 times/day Paed:Nebulized 125-250mcg 3-4 times/day.	46.00	Drymouth, Urinary retention, Broncho spasm
		IPRAVENT- 2 ML			14.00	
		DUOLIN INHALER 200 MD			233.00	
36	IPRATROPIUM BROMIDE & LEVOSALBUTAMOL	DUOLIN RESPULES-2.5ml	Respules	One respule 3-4 times/day by nebulization.	10.59	Duolin can include fine tremor of skeletal muscles and nervousness and less frequently, tachycardia, dizziness, palpitations or headache, especially in hypersensitive patients.
38	LEVOCETRIZINE	LEVOCKAST-Z	Tablet	5 mg orally once a day in the evening.	16.50	Dizziness, Fatigue.
		LEVOCETNE -Z			8.90	
		XYZAL-5mg			9.07	
40	LEVOSALBUTAMOL	LEVOLIN - 0.31 mg - 2.5 ml	Tablet, Inhaler,	Adult: 0.63 mg TID (Nebulizer) 2 mg (Tablet)	3.37	Headache, Tremor, Tachycardia.
		LEVOLIN - 2 mg	Respules, Rotacaps,		7.78	
		LEVOLIN INHALER	Syrup		120.00	
		LEVOLIN SYRUP - 100 ml			16.55	
		DUOLIN ROTOCAPS			63.00	

41	MONTELUKAST SODIUM	MONTAIR - 10 mg	Tablet	10 mg/day	137.00	Churg-Strauss syndrome; headache, dizziness, asthenia /fatigue, dream abnormalities, hallucinations, agitation.
		MONTAIR - 4mg		Once daily.	122.00	Headache, Stomach pain, Body aches or Pain, Pus in urine, Agitation and anxiety, Skin rash.
42	MONTELUKAST SODIUM & LEVOCETRIZINE	MONTEK-LC	Tablet, Syrup		114.00	vomiting, dryness in the mouth
		TELEKAST-L-KID			69.00	
44	MONTELUKAST SODIUM & FEXOFENADINE	MONTI - FX	Tablet	Adult : 10mg + 120mg	118.10	Dizziness, Fatigue.
		TELEKAST-F			121.00	
45	MONTELUKAST SODIUM & ACEBROPHYLLIN	TELEKAST-A	Tablet	Adult: 10 mg + 100mg	126.00	Dizziness, Fatigue.
46	ORCIPRENALINE SULFATE	ORCIBEST - 10 mg	Tablet	20 mg 3-4 times/day	62.00	Tachycardia, nervousness, increased serum glucose, increased potassium levels, tremor, palpitation.
47	PIRFENIDONE	PIRFENEX - 200 mg	Tablet	267 - 534 TDS	185.00	Increased ALT, increased AST, increased gamma glutamyl transferase, Photosensitivity, Rash, abdominal pain/ distension/ discomfort, headache,anorexia.
51	SALBUTAMOL	ASTHALIN INHALER	Tablet, syrup, Respules, Rotacaps, Inhaler	Oral: 2-4 mg 100- 200 mcg / inhalation	127.13	Palpitation, tremor, restlessness.
		ASTHALIN - 2.5 ml RESPULES			4.00	
		ASTHALIN(RESP)SOLUTION-15m			15.20	
		ASTHALIN SYRUP			16.33	
		ASTHALIN - 2 mg			4.80	
		ASTHALIN - 4 mg			6.00	
52	SALMETEROL & FLUTICASONE PROPIONATE	SEROFLO - 125 mg INHALER	Inhaler, Rotacaps	125 - 200 mcg	346.00	Mouth & throat candidiasis throat irritation
		SEROFLO - 250 mg INHALER			485.00	
		SEROFLO - 250 ROTOCAPS			253.00	
		SEROFLO 100 ROTACAPS			130.50	

### ANTI-BACTERIALS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMIKACIN	MIKASTAR - 250mg	Injection	250, 500 mg	31.00	Ototoxicity,. Nephrotoxicity,



2	AMOXICILLIN	NOVAMOX - 250mg	Capsules, Syrup, Tablet	Adult: 250, 500 mg	47.86	Diarrhea, Alteration of bacterial flora, rashes,
		NOVAMOX 250 REDIUSE 30ml		Paediatric dose:	48.00	
		BLUMOX-500mg			62.35	
3	AMOXYCILIN & POTASSIUM CLAVULANATE	MOXCLAV - 625mg	Injection, tablet, syrup,	amoxycillin	250.55	Diarrhea, Alteration of bacterial flora, rashes, HS reactions.
		NOVAMOX-CV - 30ml		Adult: 250, 500 mg	64.91	
		NOVAMOX-CV 375mg		Paediatric dose:	140.00	
		CLAMP - 300mg	Tablet	20- 40 mg/kg/day	75.50	
		AUGMENTIN - I.V - 600mg		Clavulanic acid:125mg	99.41	
		AUGMENTIN - 30ml			64.88	
		AUGMENTIN 625mg			259.73	
		SENSICLAV-375mg		Amoxycillin Clavulanic Acid	18.83	Diarrhea, Alteration of
		ADVENT FORTE 457mg-30ml	Syrup	Two times a day.	176.13	Nausea, Vomiting, Abdominal pain, Diarrhea, Allergy, Skin rash.
4	AMOXYCILIN + CLAVUNIC ACID	MOXYNIC-1.2mg	Injection, tablet, syrup,		131.96	Rash, Vomiting, Allergic reaction, Nausea, Diarrhoea
		ADVENT FORTE 457mg-30ml			152.50	
		AUGMENTIN-375mg		Adult (inj): Based on amoxicillin dose 500 mg tds by slow inj over 3-4 min Children (inj): Based on amoxicillin dose <3 months: 25mg/kg bd; >3 months: 25mg/kg tds	229.00	
		CLAVAM-1.2mg			127.62	
		CLAVAM-600mg		600 mg IV 24 hourly.	101.02	
6	AMOXICILLIN-250MG + CLAVULANIC ACID-50MG	AUGMENTIN-I.V-300mg	Injection	Once daily	68.00	Pain and swelling at the injection site Skin rashes Nausea Vomiting Bloating Gas.
7	AMOXICILLIN AND CLAVULANIC ACID	SENSICLAV-I.V-300mg	Injection	Once daily	83.40	Diarrhoea Anaphylactic reaction Steven-johnsons syndrome Angioedema Erythema Hypersensitivity Convulsion.
8	AMPICILLIN	ROSCILLIN-500mg	Capsules, Injection	Adult dose : 500 mg 100 - 400mg/kg/ day in 4 divided doses	43.96	Diarrhea, Alteration of

10	AMPICILLIN + CLOXACILLIN	AMPOXIN - 500mg	Capsules, Injection	Paediatric dose: upto 2 years : 62.5mg amp & 62.5 mg clox 4-6 hrly  2-10 years :  125mg amp + 125mg  clox 4-6 hrly	15.90	Diarrhea, Alteration of bacterial flora, rashes, HS reactions.
11	AZITHROMYCIN	AZIBEST- 500mg	Tablets, Syrup, Injection	Adult dose: 500 mg	65.00	Mild GI upset, Headache & dizziness     Genital itching, white patches in mouth.
		AZEE - 500mg		OD on day 1	203.34	
		AZITHRAL 15ml		then 250 mg OD	26.85	
		AZITHRAL 200mg		Paediatric dose:	63.40	
		AZIWOK 250mg		10mg/kg/day	71.38	
		AZIWOK 500mg		on day 1 then 5mg/kg/day	118.91	
		AZITHRAL-500mg		Once daily	119.50	
12	BENZATHINE PENICILLIN	PENIDURE-LA-12 LAKHS	Injection	Adult: 400 mg.	13.05	Diarrhea, alteration of bacterial flora, rashes, HS reactions.
14	CEFOPERAZONE & SULBACTUM INJ	LARICEF - 1.5gm	Injection	Adult Dose: 1 to 4 g	29.29	GI upset, Skin rash, Perineal pruritis, Depressed
		CEFGLOBE-S-FORTE 1.5gm		bd/ tid upto 12g/ day	204.50	
17	CEFIXIME	CEFTAS CV - 200mg	Tablet, Syrup	Adult: 200 mg BD	170.00	GI upset, Skin rash, Perineal pruritis, Depressed Haematopoiesis, HS reactions.
		MAHACEF - 200mg		Paed: 8mg/kg in 2 divided doses.	37.40	
		OMNIX-50mg - 30ml			71.00	
		OMNIX-100mg - 30ml			50.29	
		CEFTAS - 100mg			50.00	
		CEFTAS - 200mg			74.80	
18	CEFODROXYL	DROXYL - 500mg	Tablet	Adult: 500 - 1000 mg OD/BD Paed: 30mg/kg/day in 2 divided doses.	23.57	GI upset, Skin rash, Perineal. Pruritis, Depressed Haematopoiesis, HS reactions.
		DROXYL-DT-250mg			45.40	
19	CEFOTAXIME SODIUM	TAX-O-BID - 1gm	Injection	Adult: IM/IV 1-2g BD - QID	16.54	Inflammation at injection site, GI upset, Skin rash, Perineal pruritis, Depressed Haematopoiesis, HS reactions.
		TAXIM - 250mg		Paed: IM/IV: 100-150 mg/kg/ day in 2-4 divided doses.	21.83	
		TAXIM - 500mg			41.00	
20	CEFPODOXIME PROXETIL	CEPODEM-100mg/5ml - 30ml	Tablet, Suspension	Adult: 100 -200 mg BD Paed: 5 mg/kg BD.	139.00	GI upset, Skin rash, Perineal, pruritis, Depressed Haematopoiesis, HS reactions.
		CEPODEM -200mg			139.00	

21	CEFPODOXIME + POTASSIUM CLAVULANATE	KEFPOD CV - 200mg	Tablet, Syrup		24.40	Thrombocytopenia.
		CEPODEM -XP--325mg		Two times daily.	100.00	Eosinophilia, Leukopenia.
22	CEFPODOXIME + OFLOXACIN	CEPODEM-O	Tablet	Three times a day		Upset stomach or throwing up. Many small meals, good mouth care, sucking hard, sugar-free candy, or chewing sugar-free gum may help. Loose stools (diarrhea).
23	CEFAZOLIN	NOSTOF-500mg	Injection	Every 6 to 8 hours	25.00	Allergic reactions, Confusion, Diarrhea, Drowsiness, Eosinophilia.
24	CEFTAZIDIME	STEF-1gm	Injection	Adult :IM/IV 1-2 g OD Paed: IM/IV 50-100 mg /kg/day. Paed: IM/IV 50-100 mg /kg/day.	229.27	GI upset, Skin rash, Perineal pruritis, Depressed haematopoiesis, HS reactions
25	CEFTRIAXONE	XONE - 1gm	Injection	Adult :IM/IV 1-2 g OD; Paed: IM/IV 50-100	55.50	GI upset, Skin rash, Perineal. Pruritis, Depressed Haematopoiesis, HS reactions.
		XONE - 500mg.			139.00	
26	CEFTRIAXONE + SULBACTUM	XONE SB-1.5gm	Injection		150.00	Thrombocytopenia,Eosinophilia, Leukopenia Diarrhea, Increased liver enzymes, Allergic reaction, Rash.
27	CEFUROXIME AXETIL	CEFTUM - 250mg	Tablet	Adult dose:125-500mg.	345.00	GI upset, Skin rash, Perineal,pruritis, Depressed, haematopoiesis, HS reactions.
		CEFTUM - 500mg			129.00	
28	CEFUROXIME SODIUM	CEROXIM-500mg	Tablet	The recommended dose range is 125mg to 3gms day.	35.60	Diarrhea, Chills, Headache, Itching of vagina or genital area
		CEROXIM-250mg			19.80	
		BIPACEF-1.5gm	Injection	Six to eight hours usually for 5 to 14 days.	380.00	Diarrhea, nausea, vomiting.
		CT RAX-500mg	Tablet	Two times a day.	499.90	Rash, itching, hives, difficulty breathing or swallowing, wheezing. swelling of the face, throat, tongue, lips, or eyes.
29	CEFTAZIDIME & TAZOBACTAM	BIOTUM-TZ	Injection	125mg to 3gmsday 750 mg IM 8 hourly for up to 24-48 hours.	539.00	Diarrhea, Rash, Increased white blood cell count (eosinophils), Decreased white blood cell count, Low blood platelets, Increased liver enzymes.
30	CEPHALEXIN	SPORIDEX - 125mg- 30ml	Capsule,Suspension	Adult: 250 -500mg,QID Paed: 25 - 100 mg/kg/day in 3 divided doses.	134.79	GI upset, Skin rash, Perineal,pruritis, Depressed haematopoiesis, HS reactions,Thrombocytopenia,Eosinophilia, Leukopenia.
		SPORIDEX - 500mg			60.71	
32	CIPROFLOXACIN	CIPLOX-TZ	Tablet, Injection	Adult dose:250-500mg B.I.D	106.54	Bone marrow depression, HS reaction, Grey baby syndrome, Superinfections.
		CIFRAN - 500mg			42.03	
		ZOXAN - 500mg			40.50	
		CIPROFLOXACIN I.V - 100ml			17.85	
		CIFRAN-250mg TABLET			21.16	

35	CLARITHROMYCIN	CLARIBID - 500mg	Tablet, Injection	Adult dose: 250 - 500mg B.I.D, Paed dose: 7.5 - 15 mg/kg every 12 hrs	482.90	GI upset, Taste perversion, insomnia and confusion.
36	CLINDAMYCIN	CLINDATEC 300mg - 2ml	Injection, gel, capsule	Adult dose: 600 -1800 mg per day in four divided doses i.m /i.v: 300 - 2400 mg/day in 2-4 divided doses Adult dose: 600 -1800 mg per day in four divided doses i.m /i.v: 300 - 2400 mg/day in 2-4 divided doses child <1 mth: 15-20mg/kg /day in 3-4 divided doses child>1mth: 20-40mg/kg/day in 3-4 divided doses. child>1mth: 20-40mg/kg/day in 3-4 divided doses child <1 mth: 15-20mg/kg /day in 3-4 divided doses	92.95	GI upset, pseudomembranous. colitis, angioneurotic edema. hypotension & cardiac arrest. after rapid i.v.
		DALACIN - C - 300mg			394.00	
		CLAMIN-300mg/2ml			108.00	
		FACECLIN - 20gm			126.00	
39	COLISTIMETHATE SODIUM	COLY-MONAS-1000000IU INJECTION(COLISTIMETHATE SODIUM)	Injection	Once Daily for 2 to 5 days	1310.00	Upset stomach, Rash, Paresthesia (tingling or pricking sensation), Dizziness, Slurred speech, Vertigo, Fever.
40	DOXYCYCLINE	DOXY-1 LDR	Capsule	200 mg on day 1 as a single or in divided doses, followed by 100 mg/day.	65.25	Hypotension, Pericarditis. Angioneurotic oedema, Dyspnoea, Serum sickness,Peripheral oedema, Tachycardia, Urticaria, Haemolytic anaemia, Thrombocytopenia, Neutropenia.
		DOXYCYCLINE - 100mg	Capsule	Adult dose: 100 mg OD children > 8 yrs : 2-5mg/kg/day in 1 to 2 divided doses.	35.00	Anorexia, GI upset, photosensitivity, exfoliative dermatitis, superinfections, liver damage.
41	ERYTHROMYCIN	ERYTHROCIN - 500mg	Tablet	Adult dose: 250 to 500 mg QID max 4g/day. Paed dose: 20 - 40 mg/kg/day in four divided doses.	56.68	Mild GI upset, headache & dizziness.
42	FAROPENEM SODIUM	DUONEM ER - 300mg	Tablet, Capsule	150 – 200 mg, thrice daily.	473.50	Diarrhea, nausea, vomiting,
		FAROBACT - 200mg			332.50	GI haemorrhage, fever,chills, tachycardia, hypertension,hypotension, MI, Dyspepsia,Hypochromic anaemia.
43	FOSFOMYCIN TROMETHAMINE	MONUROL-SACHET	Sachet	1 packet (sachet) as a single dose. Mix this drug with water	380.00	Nausea, diarrhea, headache, vaginal discomfort, or dizziness may occur.
48	GENTAMICIN	GENTICYN 80mg-2ml	Injection	Adult: 1-1.7 mg/kg every 8 hrs ; Paed 2-2.5 mg/kg/day in 3 divided doses.	4.87	Ototoxicity, nephrotoxicity & neuromuscular blockade.
49	IMIPENEM + CILASTATIN	PRIMAPEN - 500mg	Injection	Adult: 250-500 mgIV infusion 6 hrly Paed: 25 mg/kg IV infusio 8 hrly.	866.00	Nausea, vomiting, diarrhea, fever, hypotension, pruritis, utricaria, rashes.

51	LEVOFLOXACIN	LEON - 750mg	Tablet, injection	Adult dose : 500 mg OD	76.00	Dizzines, drowsiness, disorientation, hot & cold flushes, slurring of speech
		LEVOFLOX - 100ml			118.00	
		LEVOFLOX - 500mg			103.00	
		LOXOF-500mg TABLET	Tablet	Once daily with or without food.	8.47	Diarrhea, Abdominal pain, Confusion,Fever, Redness of skin, Change in taste, Constipation, Dizziness,Headache,Nausea or vomiting.
52	LINEZOLID	LIZOFORCE - 30ML	Tablet, Syrup,	Adult: PO 400 mg 12 hrly	108.77	Reversible myelosuppression,convulsions and seizures, rashes,dizziness, nausea and vomiting, (e.g. leucopenia, thrombocytopenia, anaemia,pancytopenia); headache, fever,metallic taste, insomnia, constipation, diarrhoea, abdominal pain, skin disorder,cough, pharyngitis.
		LINID-TABLET(LINEZOLID)	Injection	PO/IV600 mg 12 hrly .	347.00	
53	MEROPENEM	EUPEN - 1GM	Injection	Adult: 1-3g TID Paed: 60-120 mg/kg/day in 3 doses	1061.00	Diarrhea, nausea, vomitting, GI haemorrhage, fever, chills, tachycardia, hypertension, hypotension, MI, Dyspepsia, Hypochromic anaemia.
		EUPEN-500gm			512.00	
61	NITROFURANTOIN	NIFTAS-50mg	Tablet	50-100 mg 4 times/day.	70.00	Nausea, vomiting, anorexia, abdominal pain, diarrhoea;headache, drowsiness, vertigo, dizziness, nystagmus, benign intracranial hypertension; rash, urticaria.
		URIFAST-100MG	Tablet	50-100 mg 4 times/day	123.90	
63	NORFLOXACIN	NORFLOX - 400MG	Tablet	Adult: 400 mg BD	48.50	QT prolongation, phototoxicity, GI upset
		NORFLOX-TZ			66.80	
64	OFLOXACIN	OFLER - 200 MG	Tablet, Injection,	Adult: 200-800 mg OD.	52.78	Abd pain, Diarrhoea, Nausea, Vomiting
		OFLOX - 100 ML	Syrup		94.00	
		OFLOX - 200 MG			56.69	
65	OFLOXACIN + TINIDAZOL	OFLOX - TZ	Tablet, Injection,	Adult: 200-800 mg OD.	96.50	
70	PENCILLIN G POTASSIUM	PENTIDS - 400 MG	Tablet	Adult: 400 mg	6.24	Diarrhea, alteration of bacterial flora, rashes, HS reactions.
71	PIPERACILLIN & TAZOBACTAM	MICROTAZ-2.25gm	Injection	3.375 g IV every 4 hours or 4.5 g IV every 6 hours	191.32	Nausea, constipation, diarrhoea, headache, and trouble sleeping.
		PIROTAZ-4.5gm	Injection		546.24	
72	PIPERACILLIN + TAZOBACTAM	PIPZO - 2.25MG	Injection	Adult: 2.25- 4.5 g IV 6 hrly	154.50	GI disturbance, hypokalemia, hyponatremia, platelet dysfunction.
		TAZAR-1.125GM			98.50	
		ZOPERCIN - 4.5GM			375.00	
		PIPTUM- 4.5gm- INJECTION(PIPERACILLIN&TAZ OBACTAM	Injection	Adult: 2.25- 4.5 g IV 6 hrly	398.00	

75	RIFAXIMIN	RIFAGUT - 400MG	Tablet	200 mg orally 3 times a day.	239.00	GI upset, periphaeral edema, Pyrexia, Asthenia.
76	ROXITHROMYCIN	ROXID - 150 MG	Tablet	Adult: 150-300 mg BD Paed: 2.5 - 5 mg/kg BD	102.32	Mild GI upset, headache & dizziness.
79	SULPHAMETHOXAZOLE AND TRIMETHOPRIM	SEPTRAN-DS	Tablet	Once daily	25.20	High potassium levels in blood causing abnormal heartbeats Fungal infection (thrush or candidiasis) of mouth or vagina Headache, nausea, diarrhoea Skin rashes.
80	SULFAMETHOXAZOLE	COTRIM-5ml INFUSION	Injection	120mg/kg/day divided into a 6 to 8 hourly dosing regimen for 14 days.	64.50	Nausea,vomiting,loss of appetite,diarrhea,joint or muscle pain.
85	VANCOMYCIN HCL	VANKING - 1 GM	Injection	Adult: 1g IV 12 hrly	591.00	Allergic reaction, Rash, Itching, Decrease in blood pressure, Breathlessness, Flushing.
		VANKING - 500MG			379.00	
86	METRONIDAZOLE	FLAGYL - 100ml	Tablet, Suspension	Adult: TAB 250 mg - 750mg INJ: 15mg/kg IV.	12.83	GI upset, metallic taste, seizures, Peripheral neuropathy.
		FLAGYL - 400mg			12.29	
		METRONIDAZOLE - 400			12.29	
		FLAGYL-60ml			17.01	
		METROGYL GEL-30gm			63.10	
87	METRONIDAZOLE + NORFLOXACIN	NOR- METROGYL-30ml	Suspension	Paed : 100mg +100 mg.	12.54	Dizziness, headache, stomach upset, nausea, vomiting, loss of appetite, diarrhea, constipation, or metallic taste in your mouth.
88	ORINDAZOLE	ORDENT	Tablet	Adult: 500 mg - 1.5g Child 10-40mg/kg OD.	97.00	GI upset, vertigo, headache.
89	SATRANIDAZOLE	SATROGYL - 300mg	Tablet	Adult: 300 - 600 mg OD.	42.25	Headache, dry mouth, weakness and dizziness.
90	SATRANIDAZOLE + OFLOXACIN	SATROGYL - O	Tablet	Once Daily	106.00	Nausea.Vomiting. Stomach pain. Loss of appetite. Dizziness.
91	SECNIDAZOLE	SECNIL FORTE - 1gm	Tablet	Adult: 1.5-2g OD	100.00	Nausea, gastralgia, change of taste, metallic taste, stomatitis, urticaria, rashes, leucopenia. Rarely vertigo, moderate neurological, digestive disturbances.

92	TINIDAZOLE	FASIGYN - 500mg	Tablet	Adult: 2g OD	28.00	Transient leucopenia, ataxia, convulsions, peripheral neuropathy, sensory disturbances, vertigo, metallic taste, flushing. Potentially Fatal: Hypersensitivity.
93	BETAMETHASONE + NEOMYCIN	BETNOVATE-N-20gm	Cream	Two or three times a day.	330.30	Skin itchiness, drying or cracking of skin, acne, changes in skin colour, persistent infections.
94	BETNOVATE VALERATE CREAM I,P	BETNOVATE - 20gm	Cream	apply to affected area 2-3 times per day	20.00	Itching, redness.

### ANTI-CANCER

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ATEZOLIZUMAB	TECENTRIQ-1200mg	Injection	Every Three Weeks.	396725.00	Fatigue, Weakness, Decreased appetite, Nausea, Cough, Breathlessness, Constipation.
2	BEVACIZUMAB	BEVACIREL-400mg	Injection	Once every 2 or 3 weeks.	47016.17	Dizziness, loss of appetite heartburn, change in ability to taste food, diarrhea, weight loss
		BEVACIREL-100mg	Injection	Once every 2 or 3 weeks.	14951.84	sores on the skin or in the mouth voice changes.
3	BEVACIZUMAB	BEVATAS-100mg	Injection	Once in every 2-3 weeks	11336.00	Rectal bleeding, Taste change, Headache, Nosebleeds, Back pain, Dry skin High blood pressure, Protein in urine Inflammation of the nose.
4	BENDAMUSTINE	BENDIT-100mg	Injection	Once daily.	3643.35	Anemia (low number of red blood cells), Decreased white blood cell count, Infection, Fatigue, Nausea, Fever, Vomiting, Low blood platelets, Mucosal inflammation.
5	BICALUTAMIDE	TABI-50mg	Tablet	Once daily.	1291.20	Anemia (low number of red blood cells), Rash, Abdominal pain, Breast enlargement in male, Breast tenderness, Constipation, Decreased appetite, Decreased libido, Depression, Dizziness.
6	BORTEZOMIB	BORTENAT-2mg	Injection	Twice weekly for two weeks.	2787.92	Chest pain, chills, confusion, cough.
		BORTECAD-2mg	Injection	1.3 mg/m2 as a bolus IV injection twice weekly	2605.85	General weakness, tiredness, nausea, vomiting, diarrhea, constipation, loss of appetite.

7	CARBOPLATIN	CABOO-150mg/15ml	Injection	Adult(Previously untreated patient):400mg/m2 as single short term IV infusion over 15-60min. If previously treated reduce initial dosage by 20-25%(300-320mg/m2)	809.90	Pain at the injection sitesevere, Low WBC and Platelet countsevere, Fever and chillssevere.
		CABOO-450mg	Injection		2222.00	
8	CISPLATIN	CISMAX-10mg/10ml	Injection	Administered by slow iv infusion 50-100mg/m2 BSA every 3-4 weeks	70.00	Acute renal failure, Tinnitus, Hearing loss, Leukopenia, Fever, Infection, Nausea, Vomiting, Hypocalcemia.
		CISMAX-50mg/50ml	Injection		350.00	
		CELPLAT-10mg	Injection	Once a week.	76.00	Black, tarry stools, blood in urine or stools, burning, numbness, tingling, or painful sensations, change in frequency of urination or amount of urine, cough or hoarseness, difficulty
		CELPLAT-50mg	Injection		428.00	
9	CITICOLINE (250MG)	CITIZAC-2ml	Injection	Adult: up to 1 g IM/IV daily	99.00	Decreased blood pressure, Stomach pain, Diarrhoea, Injection site reactions (pain, swelling, redness), Irregular heart rate.
10	DACARBAZINE	ZICARB-200 mg	Injection	150 mg/m <sup>2</sup> IV a Day for 5 days, OR	215.00	Nausea, Vomiting, Injection site pain, Leukopenia, Thrombocytopenia.
		ZICARB-500 mg	Injection	375 mg/m <sup>2</sup> IV on Day 1; repeat every 15 Days	470.45	
11	DACTINOMYCIN	DACILON-0.5MG	Injection	10 to 15 mcg/kg daily for 5 days, with repeat courses every 2 to 4 weeks.	369.00	Black, tarry stools. blood in the urine or stools. Cough,diarrhea (continuing) difficulty with swallowing, hoarseness.
12	DENOSUMAB	DENOCI-120mg	Injection	120 mg, for every 4 weeks.	28500.00	Fatigue/weakness, decreased level of phosphate and calcium in blood, nausea, diarrhea, headache and cough.
13	DOCETAXEL	DOCAX -120mg/3ml	Injection	Adult: 60-100mg/m2 by infusion over 1 hr every 3 wk	2756.00	Nerve damage, Fluid retention, Muscle pain, Vomiting, Weakness, Nausea, Infection, Difficulty in swallowing, Loss of appetite, Hair loss, Diarrhoea, Skin reaction, Breathing difficulty.



14	DOXORUBICIN	METODOX-10mg	Injection	Adult: 60-75 mg/m2 BSA once every 3 week	90.20	Vomiting, Allergic reaction, Nausea, Mouth sore, Hair loss, Stomach pain, Diarrhoea, Decreased appetite, Eye disorder, Increased thirst, Decreased white blood cell count (neutrophils), Heart beat irregular, Nail disorder.
		METODOX-50 -50mg	Injection	AIDS-related Kaposi's sarcoma As pegylated liposome: 20 mg/m2 once 2-3 wkly. Ovarian carcinoma As pegylated liposome: 50 mg/m2 once 4 wkly. Metastatic breast carcinoma w/ cyclophosphamide: 60-75 mg/m2 once 3 wkly.	304.00	Leucopenia, thrombocytopenia, nausea, vomiting, diarrhoea. Rarely facial flushing, rash, alopecia. Blurred vision, headache, seizures, paraesthesia, confusion, malaise, lethargy, skin pigmentation.
15	DOXORUBICIN LIPOSOME	LIPPOD-50MG	Injection	60-75 mg once in 21 days or 30 mg daily for 3 days every four weeks.	9460.00	Hair loss, Bone marrow suppression Vomiting, Rash, Inflammation of the mouth.
16	DONEPEZIL + MEMANTINE	DONAMEM-10mg	Tablet	10 mg tablet taken once a day with or without food, in the evening just before bedtime.	183.00	Nausea, Vomiting, Loss of appetite, Diarrhoea, Headache, Dizziness, Bleeding under the skin.
		DONAMEM-5mg	Tablet		125.00	
17	EPIRUBICIN	EPINEON 10mg	Injection	Dose:60-90mg/m2 iv over 5 min,repeated at 3 weeks,maximum dose<900mg/m2.	244.00	Rash, Vomiting, Weakness, Conjunctivitis, Nausea, Mouth sore, Hair loss, Absence of menstrual periods, Itching, Diarrhoea, Hot flashes.
		EPINEON 50 mg	Injection		982.00	
18	ETOPOSIDE	ETOVEL-100 mg	Injection	Dose:50-100mg/m2/day iv for 5 days	195.00	
		ETOVEL-50 mg	Capsule	100-200mg/day oral	560.00	
19	ERLOTINIB	ERLIB-150mg	Tablet	once daily	3380.10	Diarrhea, dry skin, muscle/joint pain, mouth sores, unusual eyelash growth, or eye dryness/redness/irritation may occur.
		ERLOCIP-150mg	Tablet	A 100 mg single IV dose.	8226.90	Diarrhea, loss of appetite, nausea, vomiting, heartburn, gas, constipation, mouth sores, weight loss, extreme tiredness.
20	ETOPOSIDE	ETOPLAST-100mg*/5ml	Injection	Testicular cancer- Adult: IV- The recommended dose is 50-100 mgm 2day from days 1-5. Small cell lung cancer- The recommended dose is 35 mgm 2day for 4 days to 50 mgm 2day for 5 days. May repeat course 3-4 weekly.	131.00	Stay away from people with infections, colds, or flu. Anemia, low white blood cell count, and low platelet count. Feeling tired or weak. Upset stomach or throwing up. Many small meals, good mouth care, sucking hard, sugar-free candy, or chewing sugar-free gum may help. Loose stools (diarrhea). Irritation or swelling where the shot was given. Hair loss.
		ETOPLAST-50mg	Capsule		585.00	

		POSID-50mg	Capsule	50-100 mg/m <sup>2</sup> /day from days 1-5, May repeat course 3-4 weekly.	452.57	Nausea, vomiting, loss of appetite, tongue inflammation, constipation, abdominal pain, diarrhea and taste alteration.
21	FILGRASTIM	FILNEON-300mg/0.5ml	Injection	Adult:5-10mcg/kg daily as a single daily SC inj, as a continuous IV or SC infusion, or as a daily IV infusion over 15-30 mins.	1186.65	Bone pain, Weakness, Headache, Rash, Nausea, Increased lactate dehydrogenase level in blood, Vomiting.
22	FLUOROURACIL	CHEMOFLURA - 500mg-10ml	Injection	12 mg/kg/day (max: 0.8-1g/day) for 3-4 days.	21.60	Leucopenia, thrombocytopenia, stomatitis, GI ulceration, bleeding and diarrhoea, haemorrhage from any site (stop treatment).
		FLURACIL - 250mg			11.61	
		CHEMOFLURA-250mg	Injection	Adult: 12mg/kg/day(max 0.8-1g/day) for 3-4 days, maintenance dose:5-15mg/kg/wk(max 1g/wk. As an infusion,15mg/kg/day (max.1g daily) in 500ml NS or 5% glucose over 4 hrs.	10.45	Nausea, vomiting, loss of appetite, stomach ache, diarrhea, or darkening of the skin/nails.
		5FLUCEL-250mg	Injection	Once a week.	12.20	Bruising and bleeding, Anaemia (low number of red blood cells), Feeling sick, Diarrhoea, Sore mouth and throat.
23	FULVESTRANT	FULVENAT-250mg	Injection	IM on days 1, 15, 29, then once monthly.	12229.66	Nausea, vomiting, constipation, diarrhea, stomach pain, loss of appetite, sore throat, mouth sores, weakness, hot flashes or flushing.
24	GEFITINIB	GEFTIGET-250mg	Tablet	Adult: 250mg once daily	4355.00	Nausea, Rash, Vomiting, Weakness, Loss of appetite, Dry skin, Diarrhoea, Stomatitis.
		GEFTINAT-250mg	Tablet		5900.00	
25	GEMCITABINE	GEMCHEM-1000mg	Injection	Adult: 1,000 mg/m <sup>2</sup> once weekly via infusion over 30 mins for 3-7 week and then followed by 1 week rest.This cycle is again repeated.	1300.75	Nausea, Vomiting, Loss of appetite, Hair loss, Low blood platelets, Allergic reaction, Breathlessness, Decreased white blood cell count, Increased liver enzymes, Anemia .
		GEMCHEM-200mg	Injection		402.00	
		GEMGET-1000mg	Injection		5165.00	
		GEMGET-200mg	Injection		1170.00	
26	IFOSFAMIDE & MESNA	HOLOXAN 1gm + UROMITEXAN	Injection	1.2 g/m <sup>2</sup> per day for 5 consecutive days. Treatment is repeated every 3 weeks.	463.22	Hair loss, Nausea, Vomiting, Anemia, Blood in urine, Decreased white blood cell count, Increased risk of infection, Neurotoxicity.
		HOLOXAN 2gm + UROMITEXAN	Injection		1157.96	
27	IMATINIB	CHEMOTINIB-400mg TABLET	Tablet	400 mg a Day		Neutropenia, Nausea, Muscle cramps Musculoskeletal pain, Thrombocytopenia, Rash.
28	LEUPROLIDE	LUPRODEX -(3.75)	Injection	3.75 mg alone or in combination with norethindrone acetate is six months.	2719.00	Decreased libido, Testicular atrophy, Increased sweating, Fatigue, Muscle weakness, Erectile dysfunction, Bone pain, Injection site reaction, Hot flashes.
		LUPRODEX-3m(11.25)	Injection	Adult: 1mg as single daily dose by SC inj.	11025.00	Decreased libido, Testicular atrophy, Increased sweating, Fatigue, Muscle weakness, Erectile dysfunction, Bone pain, Hot flashes.

29	LOMUSTINE	BELUSTINE-40mg	Capsule	single dose(40mg) every 6 to 8 weeks.	1125.00	headache, stomach pain, nausea, diarrhea, vomiting, gas.
		LOMOOTHER-40mg TABLET(LOMUSTINE)	Tablet	Single dose every six to eight weeks.	599.76	Nausea, vomiting, loss of appetite, diarrhea.
30	LEUCOVORIN	NEOVORIN-15mg-TABLET	Tablet	15 mg (10 mg/m2) every 6 hours.	177.00	Skin rash, hives, or itching wheezing.
31	NIMOTUZUMAB	BIOMAB EGFR-(4x50mg)	Injection	200 mg as a continuous intravenous infusion weekly with combination of radiation therapy for 6 weeks followed by 200 mg every 15 days.	59730.60	Gastrointestinal: Nausea, vomiting • Skin: Skin rash • Cardiovascular: Changes in blood pressure • Central nervous system: Drowsiness, disorientation • Others: Fever, chills, rigors, weakness, allergic reaction, blood in the urine, increase in liver enzymes.
		OPDYTA-100mg/10ml	Injection	3mg/kg to be injected as an intravenous infusion over one hour. It is administered once every two weeks.	99500.00	Chest pain, severe cough, breathing difficulties, abdominal pain, diarrhea, jaundice, ankle swelling, hormonal problems, muscle weakness.
		OPDYTA-40mg/4ml	Injection		39800.00	
32	MESNA	CANCENA-200mg/2ml	Injection	Adult and children>4 month:240mg/m2(if receiving 1.2g/m2 ifosfamide dose) or 20% of cyclophosphamide dose administered either by short(15-30 mins) or continuous (24 hrs) infusion	26.85	Nausea, Headache, Lightheadedness, Lethargy, Abdominal pain, Flushing, Diarrhoea, Rash, Fever, Flu-like symptoms.
33	OXALIPLATIN	OXYREA-100mg	Injection	Adult:85mg/m2 every 2	1376.00	Decrease in platelet counts, nausea, vomiting and diarrhea,constipation and abdominal pain.
		OXYREA-50mg	Injection		751.00	
		OXITAN-100mg	Injection	2 to 6 hours	2000.00	Nausea, vomiting, mouth ulcers, loss of appetite, indigestion, numbness, nerve pain, tiredness.
34	PACLITAXEL	TAXELEON-100mg-16.5ml/7ml	Injection	Adult:135mg/m2 infused over 24 hr and repeat at 3 weeks intervals.	618.00	Nausea, Vomiting, Weakness, Rash, Low blood platelets, Upper respiratory tract infection, Urinary tract infection, Bleeding, Anemia.
		TAXELEON-260mg/43.34ml	Injection		1536.00	
35	PACLITAXEL NANOPARTICLE	PACLITAX NAB-100mg	Injection	20 mL of 0.9% Sodium Chloride Injection.	8232.68	Low red blood cell count (anemia), Feeling weak or tired, Hair loss, Numbness, tingling or burning in hands or feet (neuropathy), Joint and muscle pain.
36	PAZOPANIB	PAZOICI-400MG	Tablet	Once daily.	9995.00	Headache, Nausea, Vomiting, Musculoskeletal (bone, muscle or joint) pain, Fatigue, Change in color of hair, Diarrhea, High blood pressure, Decreased appetite, Weight loss, Taste change, Abnormal skin pigmentation.

37	PEMETREXED	PEMETERO-100mg	Injection	45 mL/min or greater is 500 mg/m <sup>2</sup> as an intravenous infusion over 10 minutes	1014.00	Nausea, Vomiting, Sore throat, Loss of appetite, Stomatitis (Inflammation of the mouth), Hair loss.
38	PEGFILGRASTIM	PEGNEON-6mg/0.6ml	Injection	Adult: 6 mg subcutaneously once a week for 2 doses ,  Child: Less than 10 kg: 0.1 mg/kg 10 to 20 kg: 1.5 mg 21 to 30 kg: 2.5 mg 31 to 44 kg: 4 mg <del>45 kg or greater: 6 mg</del>	7200.00	sudden or severe pain in your left upper stomach spreading up to your shoulder;
		PEG RELIGRAST-6mg/0.6ml	Injection	At once.	3855.38	Nausea, Headaches, Bone pain, Pain in extremity, Pain at the site of injection.
39	PERTUZUMAB	PERJETA-420MG	Injection	The initial recommended dose is 840 mg administered as a 60-minute intravenous infusion, followed every 3 weeks thereafter by 420 mg administered as a 30 to 60 minute intravenous infusion..	212000.00	Nausea, diarrhea, temporary hair loss, fever, low blood cell counts, tiredness, rash, numbness, tingling.
40	PEMETREXED	PEMETERO-500mg	Injection	Intravenously on Day 1 of each 21-day cycle.	2673.00	Nausea, Vomiting, Sore throat, Loss of appetite, Stomatitis (Inflammation of the mouth), Hair loss, Low blood platelets, Rash, Fatigue.
41	IRINOTECAN	INTENSIC-100mg/5ml	Injection	Adult: 125 mg/m <sup>2</sup> infused IV over 90 mins once weekly for 4-6 weeks.	781.00	Fatigue, Nausea, Vomiting, Weakness, Hair loss, Fever, Anemia, Diarrhoea, Decreased white blood cell count, Decreased appetite, Mucosal inflammation.
		INTENSIC-40mg/2ml	Injection		442.00	
42	IFOSFAMIDE	IPOGET-1 gm	Injection	Adult: 1.2g/m <sup>2</sup> /day for 5 days (slow iv infusion for 30 mins)repeat every 3 week	493.00	Nausea, Vomiting, Anemia (low number of red blood cells), Hair loss, Decreased white blood cell count, Infection, Blood in urine, Cns toxicity.
		IFONEON-1000mg	Injection	1.2 g/m <sup>2</sup> per day for 5 consecutive days.	466.50	Nausea, Vomiting, Anemia (low number of red blood cells), Hair loss, Decreased white blood cell count, Infection, Blood in urine, CNS toxicity.
43	TAMOXIFEN	ENTAX-20mg	Tablet	20 mg given in 1-2 divided doses. Max: 80 mg/day.	28.00	Vaginal discharge, Hot flashes.

44	TEMOZOLAMIDE	TEMOTERO -100mg	Capsule	150mg/m2 once daily for 5 consecutive days, then 23 days off; for Cycles 2 through 6: increase to 200mg/m2 once daily for 5 consecutive days if tolerated, then 23 days off.	7534.90	Nausea, vomiting, diarrhea, constipation loss of appetite, sores in the mouth and throat, headache.
45	TRIPTORELIN ACETATE	DECAPEPTYL-0.1mg	Injection	In group I, the patients received one single injection of 1.87 mg Triptorelin in previous mid-luteal phase. In group II, Decapeptyl 0.1 mg per day started from previous mid-luteal phase.	337.37	Headache, heartburn, constipation.
46	TRASTUZUMAB LYOPHILIZED	TRUMAB-440mg	Injection	Once a week.	20000.00	Diarrhea, redness/irritation at injection (IV) site, muscle/joint/back pain, stomach/abdominal pain.
		TRASTUREL-440mg	Injection		20000.00	
47	VINBLASTINE	CHEMOBLAST-10mg/10ml	Injection	Once weekly. 1st dose: 3.7mg/ml as a single dose, continue to increase dose by increments.	201.50	Nausea. Vomiting. Decreased appetite. Decreased white blood cell count. Diarrhoea.
		UNIBLASTIN-10MG/10ml	Injection	Once a week	223.50	Weakness Stomach Pain Vomiting Breath Shortness Headache Increase in Blood Pressure Weakness.
48	VINORELBINE	NEOBEN-50mg/5ml	Injection	Once a week.	5816.00	Nausea, Vomiting, Weakness, Fatigue, Hair loss, Low blood platelets, Increased liver enzymes, Fever, Anemia (low number of red blood cells), Bleeding, Constipation.
49	ZOLEDRONIC ACID	ZOLDONAT-4mg	Injection	Adult: 4-5mg as a single infusion over at least 15 min. Retreatment after at least 7 days if necessary.	2500.00	Headache, Back pain, Musculoskeletal (bone, muscle or joint) pain, Indigestion, Heartburn, Diarrhoea.

### **ANTI-COAGULANTS AND THROMBOLYTIC**

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	APIXABAN	PIXAFLO-2.5mg	Tablet	Adult: 2.5 mg - 5 mg BID	127.50	Anemia, hemorrhage, hypotension, GI upset, CNS toxicity Skin rash, allergic reactions, fainting, nausea, and anemia.
		PIXAFLO-5mg			157.50	
		ELIQUIS-5mg			725.00	
		ELIQUIS-2.5mg			725.00	

2	CARBETOCIN	CARITEC-1ml	Injection	At once.	440.00	Abdominal pain, dizziness, faintness, feeling of warmth, headache, light-headedness, itching skin, trembling.
3	CILOSTAZOL	PLETOZ- 50mg	Tablet	100 mg bid	77.00	Dizziness, palpitations, diarrhoea, abnormal stools; pain, infection; peripheral oedema.
		STILOZ-50mg			97.50	
4	CLOPIDOGREL	CLOPILET - 75mg	Tablet, Capsule	75 mg OD	71.01	Dyspepsia, abdominal pain, nausea, gastritis, vomiting, flatulence, constipation, gastric and duodenal ulcers.
		CLOPILET-A-150mg			38.11	
		CLOPILET -A-75mg			34.25	
		PLAVIX-75mg			147.44	
5	DABIGATRAN	DABIGO-110mg	Tablet	110 mg BID	148.00	Abdominal pain upper, abdominal pain, abdominal discomfort, and epigastric discomfort.
		PRADAXA-150mg	Capsule	Twice daily.	718.00	
		PRADAXA-110mg			718.00	
6	ENOXAPARIN SODIUM	ENCLEX - 60mg -0.6ml	Injection	20-40 mg OD	552.50	Haemorrhage (including at the inj site), peripheral or unspecified oedema, anaemia, haematuria, ecchymosis, peripheral or unspecified oedema, fever, confusion, nausea, diarrhoea, dyspnoea, inj site pain.
		ENCLEX-40mg		Once daily	400.72	
		LMWX-60mg			638.26	
		LMWX-40mg			427.79	
7	ERYTHROPOITINE	EPOSIS 4000IU	Injection	600 u/kg twice wkly for 3 wk before surgery.	850.00	Hypertension, myalgia, arthralgia, flu-like syndrome, rashes and urticaria.
8	ETHAMSYLATE	ETHAMSYLATE - 2ml	Injection	Adult: 250-500 mg Low birth-weight neonate: 12.5 mg/kg	23.00	Transient hypotension.
9	FONDAPARINUX SODIUM	FONDAZEST-2.5mg/0.5ml	Injection	The usual dose is 2.5 mg once a day, injected at about the same time each day. If there is kidney disease, the dose may be reduced to 1.5 mg once a day.	820.00	Bleeding, Anemia (low number of red blood cells).
		FONDAFLO-2.5mg/0.5ml	Injection		784.50	
10	HEPARIN SODIUM	THROMBOPHOB - 20gm	Injection, Ointment	75-80 U/kg or 5,000 U then 18 U/kg/hr or 1,000-2,000 U/hr.	75.00	Hypersensitivity reactions painful, ischaemic and cyanosed limbs; osteoporosis delayed and cyanosed limbs; osteoporosis delayed transient alopecia, priapism, rebound hyperlipaemia.
		LOFH-25,000 IU			237.30	
		HEPAGLAN-25000IU			220.00	
		HEPARINE-25000IU			145.00	
		CELHEP-25000IU		339.02	Red itchy patches or lumps at the site of injection, unusual bleeding, or skin redness.	
				20,000 to 40,000 units per 24 hours by continuous IV infusion.		

11	NICOUMALONE	ACITROM -1mg	Tablet	4-12 mg on the 1st day and 4-8 mg on the 2nd day.	150.00	Alopecia; fever, nausea, vomiting, diarrhoea;skin rash; cholestatic liver damage Haemorrhage.
		ACITROM - 2mg			235.00	
		ACITROM - 3mg			300.00	
		ACITROM - 4mg			467.20	
12	N-BUTYL CYANOACRYLATE	ENDOCRYL-1ml	Injection	At once.	1310.49	Nausea, Acute Toxicity, Rash, Stomach Upset.
13	PROTAMINE SULPHATE	PROTA - 5ml	Injection	1-1.5 mg per 100 u of heparin	48.88	Hypotension, bradycardia, pulmonary and systemic hypertension, dyspnoea, warm sensation, transitory flushing, nausea and vomiting, lassitude.
14	RIVAROXABAN	RIVAXON-2.5mg	Tablet	Twice a day	70.00	Pain in stomach, Nausea or vomiting, Loose motions or constipation, Dizziness or fainting, Indigestion.
15	STREPTOKINASE	ICKINASE - 1500000 I.U	Injection	1.5 million u	2270.00	Fever, Headache, Nausea and Vomiting, Tiredness and weakness, Shortness of breath, Diarrhea, Reduction in blood cell count, Infections.
16	TENECTIPLASE	TENECTASE-20mg	Injection	Tenecteplase is for intravenous administration only. The recommended total dose should not exceed 50 mg and is based upon patient weight.	28900.00	Bleeding or bruising of any kind, especially around the place of injection collection of blood under skin.
		ELAXIM-40mg		Less than 60 kg: 30 mg IV bolus administered over 5 seconds.	43900.00	Abdominal cramps, cardiac arrhythmias, headache, transient blanching, increased arterial pressure. Potentially Fatal: MI, cardiac failure.
17	TERLIPRESSIN	TERLIEON - 1mg/10ml	Injection	1-2 mg	1467.00	Bleeding, thrombocytopenia, bradycardia, dissection of the coronary artery, dizziness, oedema, pelvic pain.
18	TIROFIBAN	AGGRIBLOC - 5mg -100ml	Injection	0.4 mcg/kg/min for 30 mins, then 0.1 mcg/kg/min for $\geq$ 24-48 hr	5230.37	Tiredness, Injection site reactions (pain, swelling, redness), Nasal congestion, Musculoskeletal (bone, muscle or joint) pain.
19	TRANEXAMIC ACID	TRANEMIC-5ml	Injection	2 or 3 times a day	64.20	Pain in stomach, Nausea or vomiting, Loose motions or constipation, Dizziness or fainting, Indigestion.
20	TRANEXAMIC ACID (100MG)	TRAPIC-5ml	Injection	Adult: 0.5-1g or 10mg/kg tid or 25-50 mg/kg daily by continuous infusion Child: 10mg/kg bid or tid	69.85	Headache. Back pain. Muscle pain. Nose stuffiness.
		TRAPIC	Tablet		187.00	Diarrhoea, nausea, vomiting.

21	TRANEXAMIC ACID + MEFANAMIC ACID	TRAPIC - MF	Tablet		211.00	Nausea, Stomach pain, Diarrhoea, Headache, Nasal congestion, Sinus pain, Sinus inflammation, Rash, Tiredness, Musculoskeletal (bone, muscle or joint) pain, Anemia (low number of red blood cells).
22	TRANEXAMIC ACID (250MG) + ETHAMSYLATE (250MG)	TRAPIC-E	Tablet	1 – 1.5 g three times per day.	188.00	Nausea, Stomach pain, Diarrhoea, Headache, Nasal congestion, Sinus pain, Sinus inflammation, Rash, Tiredness, Musculoskeletal (bone, muscle or joint) pain, Anemia (low number of red blood cells).

### ANTI-DEPRESSANTS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMITRIPTYLINE HCL	TRIPTAZ - 10mg	Tablets	Adult:50-200mg	21.50	GI upset, cns toxicity, heart block, arrhythmias, orthostatic hypotension, palpitation, dry mouth, urinary retention, blurred vision.
		TRYPTOMER - 10mg			43.00	
		TRYPTOMER - 25mg			36.86	
2	CLONAZEPAM + ESCITALOPRAM OXALATE	CITALIN FORTE	Tablets	CLONAZEPAM 0.5mg+ ESCITALOPRAM 10mg combination tablet taken once daily	103.00	Nausea, Vomiting, Diarrhoea, Confusion, Memory impairment, Uncoordinated body movements, Drowsiness, Tiredness, Delayed ejaculation, Anorgasmia (decreased orgasm), Low sexual desire.
		NEXITO FROTE	Tablets	Once daily.	153.00	Delayed ejaculation, Confusion, Vomiting, Memory impairment, Drowsiness, Tiredness, Anorgasmia (decreased orgasm), Low sexual desire, Nausea, Diarrhea.
3	DESVENLAFAXINE EXTENDED RELEASE	D-VENIZ-50mg	Tablet	Once daily	13.00	Headache, tiredness, sleep changes, and brief feelings similar to electric shock.
4	DOSULEPIN	PROTHIADEN-25mg	Tablets	25mg tid, gradually increased	46.00	Cholestatic jaundice, sexual dysfunction, exacerbation of psychotic manifestation, cardiac arrhythmias, hypotension, dry mouth constipation.
		PROTHIADEN-75mg			95.50	
5	DULOXETINE HCL	SYMPTA-20mg	Tablets	30-80mg	57.50	GI upset, cns toxicity, agitation, insomnia, raise in BP.



		DUZELA-30mg	Tablet	40-60 mg/day PO initially (in single daily dose or divided q12hr for 1 week if patient needs to adjust to therapy) For some patients, consider starting at 30 mg day for 1 week, to allow patients to adjust to medication before increasing to 60 mg day Target dosage: 60 mg/day PO; not to exceed 120 mg/day.	174.00	GI upset, cns toxicity, agitation, insomnia, raise in BP.
6	ESCITALOPRAM OXALATE	NEXITO - 10mg	Tablets	5 - 20 mg/day.	76.00	Nausea, diarrhoea, increased sweating, insomnia, impotence, ejaculation disorder, fatigue, somnolence; postural hypotension, sinusitis, taste disturbances. Increased appetite and wt gain.
		NEXITO - 5mg			45.00	
7	ESCITALOPRAM OXALATE + CLONAZEPAM	NEXITO PLUS	Tablets	Once daily.	100.30	Nausea, diarrhoea, increased sweating, insomnia, impotence, ejaculation disorder, fatigue, somnolence; postural hypotension, sinusitis, taste disturbances. Increased appetite and wt gain.
		ESCILATE Z -10mg			85.20	
8	FLUOXETINE	PRODEP - 10mg	Capsules	20-50 mg	36.00	GI upset, cns toxicity, epistaxis, serotonin syndrome .
		PRODEP - 20mg			40.17	
9	IMIPRAMINE HYDROCHLORIDE	ANTIDEP	Tablets	Initial: 75 mg/day, up to 150-200 mg/day if needed.	8.61	Sinus tachycardia, AV/bundle-branch block, postural hypotension, dry mouth, wt loss/gain constipation, urinary hesitancy/retention, impotence.
10	MIRTAZAPINE	MIRTAZ -15mg	Tablets	15-45mg	79.00	GI upset, cns toxicity, agitation, insomnia, raise in BP.
11	SERTRALINE	SERTA-25mg	Tablets	50-200mg	38.00	GI upset, cns toxicity, epistaxis, serotonin syndrome .
		SERTA-50mg			70.00	
12	TOFISOPAM	TOFICALM-50mg	Tablet	50-300 mg per day divided into three doses.	12.80	Nausea stomach discomfort fever dry mouth skin rash sleeplessness headache vomiting drowsiness.
13	VENLAFAXINE ER	VENIZ XR -75mg	Capsules	75 mg/day in 2 or 3 divided doses	68.00	Changes in behaviour, suicidal ideation, agitation, tremor, nervousness, anxiety, insomnia, confusions, abnormal dreams, HTN, nausea, headache, asthenia.
14	VORTIOXETINE	VORTICA-5MG	Tablets	Once daily	90.00	Nausea. vomiting. diarrhea. constipation. gas. dry mouth. dizziness.
		VORTICA-10MG	Tablets	Once daily	160.00	

## ANTI-DIABETIC

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	CANAGLIFLOZIN	SULISENT - 100mg	Tablet	100 mg orally once daily increased to 300 mg once daily	1647.00	Urinary tract infections,increased urination,yeast infections,vaginal itching,thirst,constipation, nausea, fatigue.
2	DAPAGLIFLOZIN	FORXIGA-10mg	Tablet	Once daily.	450.00	Thrush, Back pain, Peeing more than normal, Feeling dizzy, Mild skin rash.
		UDAPA -5mg			126.50	Weakness, feeling tired, Nausea, vomiting, pain in the stomach, Rash, hives, itching of the skin, breathing difficulties, swelling of the face, lips, mouth or the tongue.
3	DAPAGLIFLOZIN, METFORMIN	UDAPA M -10/500mg	Tablet	Two times a day.	114.50	Nausea, Vomiting, Diarrhea, Genital fungal infection, Nasopharyngitis.
4	EMPAGLIFLOZIN	GIBTULIO-10mg	Tablet	Once daily.	513.00	Frequent urination, dizziness, or lightheadedness.
5	GLIBENCLAMIDE	DAONIL - 5mg	Tablet	Adult dose:start at	11.11	Abdominal cramps, diarrhea, joint and muscle pain, leukopenia.
6	GLIBENCLAMIDE + METFORMIN	GLUCORED	Tablet	1.25 - 2.5 mg gradually increase upto 20 mg/day	24.50	Abdominal cramps, diarrhea, joint and muscle pain, leukopenia.
		GLUCORED FORTE			32.50	
7	GLICLAZIDE	GLIZID-80mg	Tablet	Adult dose: 160 mg in two divided doses.	90.75	Abdominal cramps, diarrhea, joint and muscle pain, leukopenia.
		DIAMICRON XR - 60mg			152.28	
		DIAMICRON-MR			63.39	
8	GLICLAZIDE +METFORMIN	RECLIMET	Tablet	Once daily	120.00	Dizziness,Sweating,Palpitations Hunger pangs ,hypoglycaemia,Dry mouth,Dry skin.
9	GLIMEPIRIDE	GP-1mg	Tablet	Adult dose: Initially 1mg once daily followed by maximum upto 6 mg.	38.49	Abdominal cramps, diarrhea, joint and muscle pain, leukopenia.
		GP-2mg			62.77	
		GLIMULIN - 1mg			60.15	
		GLIMULIN - 2mg			107.40	
10	GLIMEPIRIDE + METFORMIN	GEMER - 2mg	Tablet	Once daily.	81.00	Stomach pain, nausea, diarrhoea, vomiting, headache or metallic taste.
		GEMER - 1mg			54.00	
		GLIMESTAR-M1			39.00	
		GLIMESTAR-M2			59.40	
		GLYCOMET GP - 1mg			80.50	
		GLYCOMET-GP - 2mg			118.25	
		ISRYL-M1			22.50	
		ISRYL-M2			36.00	
		GLYCOMET GP - 0.5mg			44.85	

11	GLIMEPIRIDE+ METFORMIN+ VOGLIBOSE	GLYCOMET TRIO-1mg	Tablet	Once daily.	129.24	Diarrhoea, Joint Pain, Skin Rash, Anxiety, Depression, Night Terrors, Seizure, Change In Stools, Unusual Taste In Mouth, Abdomen Discomfort, Cough, Fever.
12	GLIPIZIDE	GLYNASE - 5mg	Tablet	Adult dose: initially the maximum of 15mg	5.07	Abdominal cramps, diarrhea, joint.
13	GLIPIZIDE + METFORMIN	GLYNASE - MF	Tablet		11.53	
14	INSULIN ASPART	NOVORAPID FLEXPEN	Injection	0.2-0.4U/kg/day	612.65	Hypoglycemia, local reactions, allergy, edema.
		FIASP FLEX TOUCH		Once Daily.	756.00	Low blood sugar.
15	INSULIN GLARGINE	LANTUS CARTRIDGE - 3ml	Injection	0.2U/kg or upto 10U once daily.	493.60	Hypoglycemia, local reactions, allergy, edema.
16	INSULIN ASPART PROTAMINE & INSULIN ASPART	NOVOMIX-50 FLEXPEN	Flex Pen	Twice or three times a day.	1089.00	Redness, pain, itching, swelling or hives (itchy, painful eruptions that sting).
17	INSULIN ISOPHANE/NPH (50%) + HUMAN INSULIN/SOLUBLE INSULIN (50%)	HUMAN MIXTARD-50/50	Injection	Type 1 diabetes: Ketones moderate or less: 0.5 U/kg/day SC, Ketones large: 0.7 U/kg/day SC. Type 2 diabetes: Initial dose: 0.5-1 units/kg/day in divided doses Morning: Give 2/3rds of daily insulin SC Ratio of regular insulin to NPH (isophane) insulin 1:2 Evening: Give 1/3 of daily insulin SC.	290.00	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision.
18	BIPHASIC ISOPHANE INSULIN INJECTION	HUMAN MIXTARD 30/70	Injection	Type 1 diabetes: Ketones moderate or less: 0.5 U/kg/day SC, Ketones large: 0.7 U/kg/day SC. Type 2 diabetes: Initial dose: 0.5-1 units/kg/day in divided doses Morning: Give 2/3rds of daily insulin SC Ratio of regular insulin to NPH (isophane) insulin 1:2 Evening: Give 1/3 of daily insulin SC.	148.37	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision.

19	MIXTURE OF INSULIN & PROTAMINE CRYSTALLISED	NOVOMIX FLEXPEN	Injection	0.5-1.0 IU/kg/body weight/day.	611.31	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision.
20	SOLUBLE INSULIN	HUMAN ACTRAPID	Injection	Adult(IM):Initial loading dose of 20 units ,followed by 6 units/hr until blood glucose drops to 10 mmol/l when the dose is given 2 hrly IV: given in conc of 1 unit/ml using an infusion pump: Initially infuse at a rate of 6 units/hr ,double or quadruple the rate if blood glucose conc do not decrease by about 5 mmol/l/hr.	149.40	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision.
21	ISOPHANE INSULIN	HUMAN INSULATARD	Injection	0.3-1.0 IU/kg/body weight/day.Given once or twice a day.	149.40	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision.
22	MIXTURE OF INSULIN & PROTAMINECRYSTALLISED	NOVOMIX FLEXPEN	Injection	0.2 microgram/kg body wt/day. Titrate as reqd.	611.31	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision.
23	INSULIN ASPART + INSULIN DEGLUDEC	RYZODEG PENFILL-3ml	Injection	Adult(IM):Initial loading dose of 20 units ,followed by 6 units/hr until blood glucose drops to 10 mmol/l when the dose is given 2 hrly IV: given in conc of 1 unit/ml using an infusion pump: Initially infuse at a rate of 6 units/hr ,double or quadruple the rate if blood glucose conc do not decrease by about 5 mmol/l/hr.	1468.80	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred vision.
24	LINAGLIPTIN	ONDERO-5mg	Tablet	Once Daily.	490.00	Increased lipase level in the blood, Stuffy or runny nose, Sore throat.
26	METFORMIN	GLYCIPHAGE - 500mg	Tablet	Adult dose: 0.5 - 2.5 g in 2 divided doses.	30.26	Lactic acidosis, vitamin B12 deficiency.
		GLYCOMET - 1gm			59.17	
		GLYCOMET - 850mg			25.25	
		GLYCOMET SR- 500mg			36.12	
27	METFORMIN+TENELIGLIPTIN	DYNAGLIPT-M FORTE	Tablet	Starting dose Once daily depends upon the sugar level increased by twice daily.	105.00	Low blood sugar signs include, anger, shaking, a fast heartbeat, confusion, or sweating. Keep hard candies, glucose tablets, liquid glucose, or juice on hand for low blood sugar.

28	METFORMIN + VILDAGLIPTIN	JALRA-M - 50mg/500mg	Tablet	Two times daily.	216.70	Headache, Low blood sugar levels, (hypoglycaemia), Change in taste, Nausea and vomiting, Diarrhoea, Stomach pain, Loss of appetite.
29	SITAGLIPTIN PHOSPHATE	ISTAVEL - 50mg	Tablet	25 - 100 mg OD.	269.00	Allergic reaction, myalgia, GI discomfort.
30	SITAGLIPTIN & METFORMIN	SITARED-XR	Tablet	Twice a day	145.00	Blistering, peeling, or loosening of the skin. darkened urine.
36	SITAGLIPTIN & METFORMIN	EMSITA-M SR 100/1000	Tablet	Once daily	189.90	Low blood sugar levels, stomach upset, constipation, flatulence, metallic taste in mouth, sore throat, stuffy nose and joint pain.
37	VOGLIBOSE	VOLIBO - 0.2mg	Tablet	Adult dose: 0.2 - 0.3 mg OD.	78.00	Flatulence, abdominal discomfort, loose stool.
		VOLIBO - 0.3mg			109.50	
38	VOGLIBOSE + METFORMIN	VOLIBO - M - 0.2mg	Tablet		84.25	
		VOLIBO - M - 0.3mg			114.00	
38	VILDAGLIPTIN	GALVUS-50mg	Tablet	50mg tablet taken twice daily	328.00	Allergic reaction, myalgia, Gidiscomfort.
		JALRA OD		Once daily.	180.00	Hypoglycemia (low blood glucose level), Tremors, Headache, Dizziness, Fatigue, Nausea.
		VYLDA OD-100mg	Tablet	Twice a day	269.95	Hypoglycemia, tremor, headache, dizziness, fatigue, and nausea.

### ANTI-EMETIC

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	DOMPERIDONE	DOMSTAL - 30ml	Tablets, Syrup	Adult: 10-20 mgevery 4-6 hours.	35.60	Galactorrhea, Gynaecomastia, weakness, headache.
		DOMSTAL			26.04	
2	DOMPERIDONE & NAPROXEN SODIUM TAB	NAXDOM-500mg	Tablet	Usually 2 or 3 times a day.	7.20	Upset stomach, nausea, heartburn, headache, drowsiness, or dizziness may occur.
		NAXDOM-250mg			63.50	
3	LEVOSULPIRIDE	NEOPRIDE	Tablet	25 mg 3 times/day.	53.00	Amenorrhoea, gynaecomastia, galactorrhea, changes in libido.
4	METOCLOPRAMIDE	PERINORM	Tablet,	10-15 mg 4 times/day.	4.67	Extrapyramidal symptoms (usually acute dystonic reactions); parkinsonism; tardive dyskinesia; restlessness, drowsiness, dizziness.
		PERINORM	Injection		12.08	
5	ONDANSETRON	EMESET - 30ml	Syrup,	Adult; IV 0.15 mg/Kg, 8 mg Orally OD, Paed: 4-12 yrs 4 mg OD.	36.14	Headache, GI upset, constipation.
		EMESET - 4mg	Tablets,		55.14	
		NEOMIT - 4mg - 2ml	Injections		16.60	

		VOMIKIND MD-4mg	Tablet	4 mg orally 3 times a day		Headache, Constipation, Fatigue, Painful, Fever, Dry mouth, Anxiet.
		ZOFER-30ml	Syrup	2mg syrup every 12 hours	35.16	Headache, feeling tired or weak, hard stools (constipation).
6	PROCHLOPERAZINE MALEATE	STEMETIL	Injections	Adult: 5-10 mg TID/QID, Injection: 12.5mg IM.	49.40	Drowsiness, dizziness, amenorrhea, blurred vision, skin reactions.
		STEMETIL-MD	Tablet		65.00	
7	PROMETHAZINE	PHENERGAN-100ml	Elixir	Children(2-5yrs):5mg Children(5-10yrs):10mg Children >10 yrs and adults: 25mg To be taken in the night before the journey and then repeat after 6-8 hours as required.	28.5	Liver problems that may cause the eyes or skin to go yellow (jaundice),Muscle stiffness or shaking.
		PHENERGAN-2ml	Injection	Adult: 125-25mg, repeated at intervals of not<4hr.(Max:100mg daily).	8.35	Drowsiness, dizziness, constipation, blurred vision, or dry mouth.

### ANTI-EPILEPTICS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	BRIVARACETAM	BREVIPIIL-50mg	Tablet	Twice daily	148.00	Dizziness, Fatigue, Nausea, Sleepiness, Vomiting, Constipation, Irritation.
		BREVIPIIL-75mg			235.00	
		BRIVGARD-50mg			162.70	Constipation, nausea, vomiting, extreme tiredness or lack of energy.
		BRITZILAM-10mg	Injection	Twice a day	119.9	Anxiety, changes in behavior. chest pain or discomfort. deep or fast breathing with dizziness. dizziness or lightheadedness. double vision.
2	CARBAMAZEPINE	TEGRITAL - 200mg	Tablet,	Adult: 200- 400 mg TDS	15.90	sedation, dizziness, vertigo, diplopia, ataxia.
		TEGRITAL -100ml	syrup	Paed: 15 -30 mg/kg/day	15.24	
		TEGRITAL -400mg			37.60	
		TEGRITAL-CR - 200mg			23.47	
3	DIVALPROEX	DICORATE ER - 250mg	Tablet	Adult: 10 mg/kg/day	56.00	Somnolence, heart block, fatigue, restlessness, excitation.
4	GABAPENTIN	GABAGESIC - 300mg	Tablet,	Adult: 300 - 60mg OD- TDS.	98.00	Mild sedation, dizziness and unsteadiness.
		GABANTIN - 100mg	Capsule		50.00	
		GABANTIN - 300mg			135.00	
		GABANEURON			112.85	

		GABAPIN-300mg			240.00	
5	GABAPENTIN, METHYLCOBALAMIN	GABANEURON-100mg	Tablet	Dose: 300mg to 600mg about three times a day.	8.14	Dizziness, Sleepiness, Tiredness, Uncoordinated body movements.
6	LACOSAMIDE	LACOSET-100mg	Tablet			Feeling lightheaded, sleepy, having blurred eyesight, or a change in thinking clearly.
		LACOXA-10mg/20ml	Injection	Infuse IV over 15-60 min; 30-60 min preferable, and should be used when a 15 min administration is not required.	201.00	
		LACOXA-50mg	Tablet	Initially, 100 mg twice daily.	59.04	Feeling lightheaded, sleepy, having blurred eyesight.
		LACOXA-100mg	Tablet	300 to 400 mg/day given as 2 divided doses.	103.90	
7	LEVETIRACETAM	LEVIPIL - 500mg	Tablet,	Adult: 500 mg BD 3g OD	118.00	Somnolence, asthenia, dizziness, vertigo, depression, emotional instability, hostility, diplopia.
		LEVIPIL - 5ml	Injection		147.00	
		EPITRA-250mg			59.47	
		EPITRA-500mg			124.32	
		EPITRA-5ml INJECTION			110.88	
		LEVIPIL-100ml SYRUP			382.92	
		LEVIPIL-250mg TABLET			59.45	
		LEVERA-500mg TABLET		500 mg orally or IV twice a day	12.96	
		LEVERA-750mg TABLET(Levetiracetam)		750 mg orally or IV twice a day	204.18	
		LEVEPSY 500 MG TABLET		500 mg orally or IV twice a day	9.91	
		LEVIPIL-750mg TABLET		The maximum dose is 1500 mg twice daily.	202.86	

8	OXCARBAZEPINE	OXETOL-300mg	Tablet	Adult: As monotherapy or adjunctive therapy:Initially,600mg daily in 2 divided doses,may increase if necessary in max increments of 600mg daily at wkly interval. Maintenance dose:600-1200mg daily or up to 2,400 mg daily. Child:≥6 yrs:8-10mg/kg daily in 2 divided doses,may increase if necessary in increments of 10mg/kg daily at wkly interval. Maintenance dose:30mg/kg daily.	88.00	Dizziness, drowsiness, tiredness, nausea/vomiting, stomach/abdominal pain, headache, trouble sleeping, or constipation.
9	PERAMPANEL	TORPANEL - 2mg	Tablet	Once daily.	99.00	Fatigue, anger, nausea, and balance disorder (loss of balance). It may also cause dizziness and sleepiness.
		TORPANEL - 4mg			149.00	Dizziness, sleepiness, drowsiness, tiredness, fatigue, headache, irritability, nausea.
10	PHENYTOIN SODIUM	EPTOIN - 100 MG	Injection	15 - 20 mg/kg IV	164.58	Gum hypertrophy, hirsutism, megaloblastic anemia, HS reaction.
		EPTOIN - 50 MG		Paed: 5-10 mg/kg/day	11.50	
11	PREGABALIN	GABAWIN-25MG	Tablet,	Adult: PO Adjunct in partial seizures Initial: 150 mg/day, may increase to 300 mg/day after a wk. Anxiety Initial:150 mg/day, may increase in mg/day, may increase in	45.00	Somnolence, dizziness, headache, diplopia, blurred vision, vertigo, fatigue, irritability, arthralgia.
		GABAWIN-50MG	Capsule		72.00	
		MAXGALIN-50mg		Adult: PO Adjunct in partial seizures Initial: 150 mg/day, may increase to 300 mg/day after a wk. Anxiety Initial:150 mg/day, may increase in .	8.90	Somnolence, dizziness, headache, diplopia, blurred vision, vertigo, fatigue, irritability, arthralgia.
		PREBALIN ZZ-75MG			143.00	
		PREGABA-50mg	Capsule	Three times a day.	118.50	Blurred vision, Difficulty in paying attention, Dizziness, Dryness in mouth Edema.



12	PREGABALIN & NORTRIPTYLINE	GABAWIN-NT	Tablet	May increase to 300 mg/day after a week. Anxiety Initial: 150mg/day, may increase in increments of 150 mg weekly. Max: 600mg/day. Neuropathic pain Initial: 150 mg/day, may increase to 300 mg/day after 3-7 days. Max: 600 mg/day after a 7-day interval.	15.00	Drowsiness, dizziness, dry mouth, blurred vision, constipation, weight gain, or trouble urinating may occur.
		AXOGURD-NT		Usually 1 to 4 times daily	15.59	
13	PREGABALIN, METHYLCOBALAMIN	MAXGALIN-M-75mg	Tablet	50 mg taken three times per day  Adult: 75–150 mg taken two times per day, or 50–100 mg taken three times per day (for a total of 150–300 mg per day).	14.00	Difficulty in breathing, Chest tightness, Chills Severe diarrhea, Swelling of face, lips, eyelids, tongue, hands and feet, Blurred or double vision Dizziness.
14	SODIUM VALPROATE	ENCORATE 100MG -5ML	Tablet,	Adult: 200-800 mg TDS	30.08	GI upset, drowsiness, ataxia & tremor.
		VALPARIN - 100 ML	Syrup	Paed: 15 - 30 mg/kg/day	67.09	
		VALPARIN -200 MG	Injection		35.53	
		ENCORATE CHRONO - 200MG (C.R)			46.00	
		ENCORATE-200ml	Syrup,	Dosage should start at 600 mg daily increasing by 200 mg at three day intervals until control is achieved. This is generally within the dosage range 1000 mg – 2000 mg per day,	122.21	GI upset, drowsiness, ataxia & tremor Feeling lightheaded, sleepy, having blurred eyesight, or a change in thinking clearly. Avoid driving and doing other tasks or actions that call for you to be alert or have clear eyesight until you see how this drug affects you.
		VALPROL-CR-300mg	Tablet,		6.35	
		VALPROL-CR-200mg	Injection		4.23	
		NEOPRATE-100mg/5ml		20 – 30mg/kg I.V over 5 min	31.00	Stomach pain, feeling or being sick, diarrhoea, feeling tired or sleepy. headache. <u>weight gain.</u>
		EPILEX CHRONO - 500mg		Once or twice daily.	109.40	Headache, Tremors of hands and feet, Involuntary eye movement, Dizziness, Memory impairment, Nausea.

		VALPROL-100mg/5ml		Initial dose: 10 to 15 mg/kg orally or IV daily; increase by 5 to 10 mg/kg per week if necessary according to clinical response Maintenance dose: 10 to 60 mg/kg daily Maximum dose: 60 mg/kg daily.	29.29	Injection site reactions (pain, swelling, redness), Loss of appetite, Nausea, Vomiting, Sleepiness, Hair loss, Weight gain, Liver function abnormal, Tremor.
15	SODIUM VALPROATE + VALPROIC ACID	VALEPTOL-CR-200mg	Tablet	10-15mg/kg/day ,increased by 5-10mg/kg/day at weekly interval.Max dose:60mg/kg/day	45.00	Nausea, Vomiting, Weight gain, Loss of appetite, Swelling of gums, Headache, Sleepiness, Tremor, Hair loss, Liver injury.
		VALEPTOL-CR-300mg			67.00	

### ANTI-FUNGALS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	CLOTRIMAZOLE	CANDID - V -30gm	Tablet, Cream,	Vaginal: Adult:As pessary:100mg daily for 6 days,200mg daily for 3 days or 500mg as a single dose,alternatively apply 1,2 or 10% cream Ear: Adult: Instill 10 drops into the canal of the affected ear(s) once daily Child: 6 month -13 yrs instill 5 drops ,duration depending upon the disease Topical(creams,gel,suspension,solution,powder): Apply onto affected area once daily(nails),bd(skin)	69.25	Erythema, Stinging, blistering,
		CANDID - V3	Capsule, Powder,		55.00	Irritation, redness,
		CANDID - B -20gm	Mouth paint,		89.50	Peeling, swelling, itching,
		CANSOFT-CL (3'S)	Lotion, Ear drops,		73.50	Burning sensation, stomach pain,
		CANDID - 20ml	Vaginal suppositories.		83.50	Nausea, vomiting and fever.
		CANDID -50gm			104.00	
		CANDID - 10ml			53.00	
		CANDID -V6			58.15	
2	FLUCONAZOLE	AF 400mg	Tablet,	Adult: 100 - 400 mg OD, Paed: 6 - 12 mg/kg/ day OD	23.00	GI upset, Headache Taste perversion, HS reaction. Headache, Cutaneous rash, Anaphylaxis, Dry mouth, Alopecia, Seizure, Dizziness, Tremors.
		AF- 150mg	Injection		9.90	
		FORCAN I.V- 100ml			118.50	
3	ITRACONAZOLE	I-TYZA-200mg	Capsule	100-200 mg OD/BID	184.00	Urticaria, anaphylatic .
		I-TYZA-100mg	Capsule	100-200 mg OD/BID	87.50	Urticaria, anaphylatic
4	KETOCONAZOLE	KZ - 75gm		Adult: 200 -400 mg OD	155.00	Urticaria, Anaphylatic
		SCALPE - 75ml	Lotion, Soap, Cream		250.00	Reactions, GI upset, Constipation,
		KZ			185.00	Headache, Dizziness, Hepatitis addison's disease.

		TARRY SHINE-100ml		Adult: As 2% shampoo: Apply on scalp once daily for up to 5 days. For prophylaxis: use once daily for a max of 3 days before exposure to sunlight.	230.00	Application site reactions (burning, irritation, itching and redness).
5	ORNIDAZOLE, FLUCONAZOLE AND ONE AZITHROMYCIN (KIT)	AF-KIT	Tablet	Ornidazole: Adult:0.5-1.5gm daily for 1-7 days Azithromycin:1 gm as a single dose 1 hr before or 2 hrs after food with plenty of water . Fluconazole:50-400mg once a day daily with or without food.	21.00	GI Irritation, Nausea, Vomiting.
6	POSACONAZOLE	POSANAT-100mg	Tablet	Start dose of 300 mg (three 100 mg tablets) twice a day on the first day, then 300 mg (three 100 mg tablets) once a day thereafter. Each dose may be taken without regard to food intake.	4000.00	Diarrhea or vomiting
7	SERTACONAZOLE NITRATE	ONABET - 15gm	Cream	Adult: Apply to affected area bid for 4 wk.	130.00	Swelling, itching, burning sensation, stomach pain,redness, oozing, dry skin.
		ONABET - 30gm			230.00	
		ONABET-B - 10gm			118.00	
8	TERBINAFINE (1% W/W)	TYZA-30gm	Cream, Tablet, Cream	Adult: Apply once or bd to affected area(s) for 1-2 weeks Adult: 250 mg OD	135.00	Skin peeling, Itching.
		TYZA-15gm			69.00	
		TYZA-250mg			108.50	
9	VORICONAZOLE	QUNIL-200mg	Injection	3mg/kg per hour over 1 to 3 hours.	4600.00	Nausea, vomiting, diarrhea,headache, fever.

### ANTI-HISTAMINE

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
------	--------------	------------	-------------	------	------	-----

1	BETAHISTINE (24MG)	VERTIN-OD-24mg TABLET	Tablet	Vertigo, tinnitus and hearing loss associated with Ménière's disease :Adult: Initially 16 mg 3 times a day, dose preferably taken with food; maintenance 24-48 mg daily	335.10	Headache, Nausea, Dyspepsia.
2	BETAHISTINE	VERTIN - 16 MG	Tablet	Adult: 8-16 mg TDS	109.00	GI upset, dizziness, insomnia.
		VERTIN - 8 MG			58.00	
3	BILASTINE + MONTELUKAST	MONTEK-BL	Tablet	Once daily.	175.00	Rhinitis, Headache, Eczema, Urticaria, Abdominal pain/Upper abdominal, pain.
4	CETRIZINE HCL	CETRIZET - 10mg	Tablets	Adult: 10 mg OD, Paed: children >6 yrs 5-10mg	20.00	Dizziness, dry mouth, GI upset, Disturbed coordination, confusion.
		CETZINE-60ml	Syrup		36.14	
		ALERID-10mg	Tablet		17.33	
5	CHLORPHENIRAMINE, PARACETAMOL, AND PHENYLEPHRINE	CORIMINIC-15ml	Drops, Syrup	Drops(1mg/ml):6 months to 11 months: 0.75 mL orally 4 times a day. 1 year to 2 years: 1 mL orally 4 times a day. Oral syrup(4mg/5ml):2 to 5 years: 1.25 ml orally every 4 hours to 6 hours, not to exceed 7.5 mL in 24 hours. 6 to 11 years: 2.5 ml orally every 4 hours to 6 hours, not to exceed 15 mL in 24 hours. 12 years or older: 5 ml orally every 4 hours to 6 hours, not to exceed 30 mL in 24 hours.	54.90	Drowsiness, dizziness, dry mouth/nose/throat, headache, upset stomach, constipation, or trouble sleeping.
		CORIMINIC-60ml				
		T-MINIC-15ml	Drops		63.80	Drowsiness, dizziness, headache, blurred vision, upset stomach, nausea, constipation, or dry mouth/nose/throat.
		T-MINIC-60ml	Syrup		68.20	
6	CINNARIZINE	STUGERON	Tablets	Adult: 25-50 mg TID	130.50	Dizziness, dry mouth, GI upset, Disturbed coordination, confusion.
7	CYPROHEPTADINE	CIPLACTIN	Tablets	4 mg tid. Usual: 12-16 mg/ day in 3-4 divided doses. Max: 32 mg/day.	23.50	Confusion, disturbed coordination,dizziness, excitation, euphoria, hallucinations, headache, hysteria, insomnia, irritability,
8	DESLORATADINE	DAZIT-5mg	Tablets	5 mg once daily.	64.00	Pharyngitis, dry mouth, myalgia, fatigue, somnolence, headache, dizziness, insomnia, dysmenorrhoea.

9	DESLORATADINE + MONTELUKAST	MONDESLOR	Tablets	Once Daily	139.00	Pharyngitis, dry mouth, myalgia, fatigue, somnolence, headache, dizziness, insomnia, dysmenorrhoea.
10	FEXOFENADINE	ALLEGRA - 180mg	Tablets	Adult: 120-180 mg OD-BD, Paed:6-11 yrs: 30 mg BD	129.40	Dizziness, dry mouth, GI upset,
		ALLEGRA 120mg			141.50	Disturbed coordination, confusion.
11	HYDROXYZINE HCL	HYRAX-10	Tablets, Syrup	25 mg at night, up to 25 mg, 3-4 times/day.	17.00	Dizziness, ataxia, fatigue, drowsiness,
		HYRAX - 100ml			44.00	hallucination, seizure, dry mouth,
		ATARAX -10mg			25.17	nervousness, xerostomia, involuntary
		ATARAX - 25mg			45.00	movements, slurred speech.
12	KETOTIFEN	ASTHAFEN-1MG	Tablet	Twice a day	62.30	Irritation, Agitation, Insomnia (difficulty in sleeping), Nervousness, Sedation.
13	PHENIRAMINE MALEATE	PHENIRAMINE MALEAE- 2ml	Tablets, Injections	Adult: 25-50 mg BID Paed: 6-14 yrs: 15-22.5 mg; 1-5 yrs: 15 mg; 4 months to 1 year:7.5mg	3.04	Dizziness, dry mouth, GI upset, disturbed coordination, confusion.
		AVIL - 25mg			5.70	
14	PROMETHAZINE HYDROCHLORIDE	PHENERGAN - 100ml	Tablets, Syrup,	Adult: 25mg every 4-6 hours Paed: 0.25-0.5 mg/kg every 4-6 hours.	23.98	Dizziness, dry mouth, GI upset, disturbed coordination, confusion.
		PHENERGAN - 2ml	Injection		6.92	
	PROMETHAZINE TEOCLATE	AVOMINE-25mg	Tablet	Adult: 25mg at night, increase to 25mg bd if necessary. Alternatively, 10-25mg 2-3 times daily. Child( 2-5yr):5-15mg daily in 1 or 2 divided doses; (>5-10 yr):10-25mg daily in 1 or 2 divided doses.	28.90	Tiredness, Sleepiness.

### ANTI-HYPERTENSIVE & HYPOTENSIVE

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMLODIPINE + HYDROCHLOROTHIOZIDE	AMLONG-H	Tablet	Adult : 5-20 mg OD.	54.50	Headache, pedal edema, palpitation, abdominal discomfort.
2	AMLODIPINE	AMLONG - 10mg	Tablet	2.5 to 5 mg orally once a day.	120.00	Headaches. Make sure you rest and drink plenty of fluids. ...
		AMLONG - 2.5mg			29.85	Feeling dizzy. If amlodipine makes you feel dizzy, stop what you're doing and sit or lie down

		AMLONG - 5mg			49.35	until you feel better. ... Flushing. Try cutting down on coffee, tea and alcohol. ... A pounding heartbeat. ... Swollen ankles.
		STAMLO -2.5mg			31.00	
		STAMLO -5mg			49.05	
3	AMLODIPINE + ATENOLOL	STAMLO BETA	Tablet	Once daily	104.00	Blurred vision.cold hands or feet. difficult or labored breathing. dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly.
		AMLONG - A			88.00	
4	AMLODIPINE+ TELMISARTAN	TELMIDUCE-AM	Tablet	once daily doses of 20 to 80 mg while amlodipine is effective in doses of 2.5 to 10 mg.	61.80	Swelling of ankles or feet Difficulty in breathing Dizziness Back pain Redness of face, neck, arms and chest Muscle pain.
		TELMA AM	Tablet	Once daily	143.40	Bloating or swelling of the face, arms, hands, lower legs, or feet. difficulty breathing. headache. nasal congestion. sore throat.
5	ASPIRIN, ROSUVASTATIN & CLOPIDOGREL	ROSUVA GOLD-20mg	Tablet	Once daily	323.00	Increased bleeding tendency, Abdominal pain, Indigestion Bruise, Nosebleeds.
6	ATENOLOL	ATPARK - 50mg	Tablet	Adult: 25 - 100 mg OD, Paed: 1-2 mg/kg/day in single or 2 divided doses.	33.52	Fatigue, bradycardia, cold extremities, epigastric discomfort.
		ATPARK - 25mg			30.83	
7	BENIDIPINE	BENLONG-4mg	Tablet	2-4 mg once daily, up to 8 mg once daily if needed.	7.80	Headache, facial flushing, vomiting, palpitation, dizziness, lightheadedness and malaise.
8	BISOPROLOL FUMARATE	CONCOR COR-2.5mg	Tablet	Adult:5-10mg once daily(max.20mg/day).	53.35	Nausea, Headache, Fatigue, Constipation, Diarrhoea, Dizziness, Cold extremities.
		CONCOR-5mg	Tablet		72.38	
9	BOSENTAN	LUPIBOSE - 62.5mg	Tablet	62.5 mg bid upto 125 mg BID	762.50	Headache, nasopharyngitis, flushing, fluid retention (e.g. peripheral oedema), hypotension, palpitations, dyspepsia.
10	CARVEDILOL 3.175 MG	CARDIVAS 3.125mg	Tablet	Adult: 12.5 - 25 mg OD	39.00	Fatigue, bradycardia, cold extremities, epigastric discomfort.
		CARDIVAS-12.5mg			82.00	
		CARDIVAS-6.25mg			52.00	

11	CILNIDIPINE	NULONG - 10mg	Tablet	Adult: 5 -20 mg OD	68.50	Headache, pedal edema, palpitation, abdominal discomfort.
		NULONG - 5mg			36.00	
12	CLONIDINE	ARKAMIN	Tablet, Injection	Adult: start with 0.1 mg BID max upto2.4mg/day Paed: 5-7 mcg/kg/day	1.38	Constriction of pupil, drowsiness, bradycardia.
		CLONEON			47.00	
13	DILTIAZEM HYDROCHLORIDE	DILZEM - 30mg	Tablet	Adult:30-120 mg TDS Paed: 1.5 - 2 mg/kg/day	26.15	GI upset, AB block, bradycardia, CHF, edema.
		DILZEM-CD - 90mg			105.00	
		DILZEM SR - 90mg			94.00	
		CREMAGEL-30gm	Gel	Adult:30-120 mg TDS Paed: 1.5 - 2 mg/kg/day.	173.53	
14	ENALPRIL MALEATE	ENVAS - 5mg	Tablet	Adult: 5 -20 mg.	51.44	Dry cough, orthostatic hypotension, GI upset.
		ENVAS - 2.5mg			30.83	
15	EPLERENONE	EPLERITE- 25mg	Tablet	Once daily.	180.00	Dizziness,diarrhea,coughing,flu-like symptoms,fever and body aches,tiredness,gynecomastia.
16	LABETALOL	LABLOL - 4ml	Injection	Adult: 20-80 mg i.v upto max 300 mg.	326.50	Fatigue, bradycardia, cold extremities, epigastric discomfort.
17	LOSARTAN	LOSAR - 25mg	Tablet	Adult: 25 - 100 mg/day in 1-2 divided doses Children > 6yrs: 0.7 -1.4 mg/kg OD	43.37	Hypotension, tachycardia, bradycardia, muscle cramps.
		LOSAR - 50mg			74.65	
		REPACE - 50mg			47.99	
		LOSAR-H			120.40	
18	METOPROLOL SUCCINATE	MET XL - 50mg	Tablet, Injection	Adult: TAB 100 mg OD - 200 mg BD Inj 5-15 mg.	87.16	Fatigue, bradycardia, cold extremities, epigastric discomfort.
		PROLOMET - XL -12.5mg	Tablet		29.00	
		PROLOMET - XL - 25mg			39.12	
		METOLAR - XR - 50mg			105.50	
		METOLAR 12.5mg			27.50	
		METOLAR-5ml			12.31	
		BETALOC-25mg			91.50	
		BETALOC	Injection		15.10	
19	NIFEDEPINE	DEPIN -5mg	Tablet, Capsule	Adult: 30 - 120mg OD	48.00	Headache, pedal edema, palpitation, abdominal discomfort.
		NICARDIA RETARD -10mg			24.70	
		NIFEDINE -10mg			13.08	

		CALCIGARD-5mg		Three times daily	14.20	Swelling of the face, eyes, lips, tongue, hands, arms, feet, ankles, or lower legs Difficulty breathing or swallowing Fainting Rash.
20	NIMODIPINE (30MG)	NIMODIP-30mg	Tablet	Adult: 60mg 4 hrly beginning within 4 days of onset of haemorrhage & continued for 21 consecutive days.	160.00	Dizziness, lightheadedness, flushing, or swelling ankles/feet.
21	OLMESARTAN MEDOXOMIL	OLMETRACK -20mg	Tablet	Adult: 20-40 mg OD	76.69	Hypotension, tachycardia, bradycardia, muscle cramps.
		OLMEZEST-40mg			150.00	
		OMEN-20mg	Tablet	Once daily	127.00	Headache, dizziness, nausea, stomach pain.
22	OLMESARTAN MEDOXOMIL + AMLODIPINE	OLMEZEST-AM	Tablet		100.00	
23	OLMESARTAN MEDOXOMIL + HYDROCHLOROTHIAZIDE	OLMEZEST H - 40mg	Tablet		155.00	
24	PRAZOSIN	PRAZOPRESS-XL-5mg	Tablet	0.5 mg bid or tid	11.80	During cataract surgery, Intraoperative Floppy Iris Syndrome(IFIS)may occur. Postural hypotension, syncope, tachycardia, palpitations, lack of energy dizziness.
		PRAZOPRESS - 1mg			50.00	
25	PROPRANOLOL	CIPLAR-LA-20mg	Tablet	40 - 80 mg BID	12.60	Bradycardia, hypotension, syncope, shock, angina pectoris.Lightheadedness, giddiness, ataxia, dizziness, irritability, sleepiness, hearing loss, and visual disturbances to vivid dreams, hallucinations, and confusion.
		CIPLAR-10mg			19.28	Tiredness,Weakness Raynaud phenomenon Arrhythmia.
		INDERAL- 10mg			12.60	Bradycardia, hypotension, syncope, shock, angina pectoris.Lightheadedness, giddiness, ataxia, dizziness, irritability, sleepiness, hearing loss, and visual disturbances to vivid dreams, hallucinations, and confusion.
		CIPLAR - LA - 40mg			108.69	
26	PROPRANOLOL+ FLUNARIZINE	INDERAL-F-10mg	Tablet	10mgday at bedtime.	80.64	Weight gain,Slow heart rate,Tiredness,Depression, Drowsiness, Abnormal dreams.
27	RAMIPRIL	HOPACE - 5mg	Tablet	Adult: 2.5 - 20mg OD	180.00	Hyperkalemia, angioedema, cough, renal dysfunction.
		HOPACE-1.25mg	Capsules	Adult: 2.5 - 20mg OD.	38.28	



28	ROSUVASTATIN & CLOPIDOGREL	ROSUVAS-CV-10MG	Tablet	Once daily	167.00	Weakness, Headache, Dizziness Muscle pain, Increased glucose level in blood, Increased liver enzymes.
29	TELMISARTAN	TAZLOC-40mg	Tablet	Adult: 20-80 mg/day Initial dose: 40 mg orally once a day Maintenance dose: 40 to 80 mg orally once a day	78.71	Hypotension, tachycardia, bradycardia, muscle cramps.
		TELMA - 20mg			65.60	
		TELSARTAN - 80mg			142.00	
		TELMA - 40mg			8.19	
		ARBITEL - 20mg			45.00	
		ARBITEL - 40mg			81.50	
		TELMIDUCE-20mg			1.50	
		TELMIGET-40mg			5.75	
30	TELMISARTAN + CHLORTHALIDONE	TELMA-CT -40mg	Tablet	Once daily	144.75	
31	TELMISARTAN + HYDROCHLOROTHIAZIDE	TELMA - H -80mg	Tablet	Once daily	269.18	Bladder pain.decreased urine. dizziness, faintness, or lightheadedness when getting up from a lying or sitting position. fast, pounding, or irregular heartbeat or pulse.increased thirst.
		TELMA - H			161.81	
32	TICAGRELOR	TICASPAN-90mg	Tablet	Twice daily.	250.00	Bruising, bleeding more easily, nosebleeds, headache, dizziness, cough, nausea, diarrhea.
		AXCER-90MG			314.30	
33	VERAPAMIL HCL 40MG	CALAPTIN - 40mg	Tablet	Adult: 80 - 160 mg bd	8.03	Headache, pedal edema, palpitation, abdominal discomfort.
34	PHENYLEPHRINE	PHENPRESS-LS-50mg-10ml	Injection	0.1 mg/kg 2 hourly	108.00	Increased blood pressure Blurred vision.
		PHENPRESS 10MG - 1ML	Injection	IM- 2 to 5 mg. IV- 0.1 to 0.5mg by slow infusion.	227.00	Anxiety, reflex bradycardia, tachycardia, arrhythmias,headache, cold extremities/gangrene, hypertension,nausea, vomiting, sweating, weakness, fear, restlessness.
35	PROPRANOLOL	BETACAP-TR-20mg	Capsule	40-80 mg per day	19.48	Tiredness, Cold extremities, Nightmare, Stomach pain, Diarrhoea, Slow heart rate.
		BETACAP-TR-40mg			37.79	

**ANTI-MALARIAL**

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ARTEMETHER & LUMEFANTRINE	LUMERAX - 80mg	Tablet	80 mg/day.	175.00	Mild GI disturbance, dizziness, tinnitus, reduction in reticulocyte and leucocyte counts, nausea, vomiting, abdominal pain, bradycardia, 1st-degree heart block.
2	ARTESUNATE	FALCIGO	Injection	Adult: IV/IM Initial: 2.4 mg/kg. Repeat 12 hr and 24 hr later, then once daily thereafter.	224.82	Transient and reversible , rash, reticulocytopenia, drug fever bradycardia, transient 1st-degree heart block and reversible elevation of serum transaminases.
3	CHLOROQUINE PHOSPHATE	LARIAGO - 250mg	Tablet, Suspension	Initial: 600 mg, then 300 mg 6-8 hr later on day 1. Continue w/ 300 mg/day on days 2 and 3	8.21	Retinopathy, hair loss, photosensitivity, tinnitus, myopathy (long-term therapy). Psychosis, seizures, leucopenia and rarely aplastic anaemia, hepatitis, Glupsets, dizziness, hypokalaemia.
		LARIAGO - 60ml			18.62	
		LARIAGO - DS			8.04	
4	HYDROXY CHLOROQUINE	HCQ	Tablet	200-400 mg/day. Max: 6.5 mg/kg/day	65.49	Retinopathy, hair loss, tinnitus, photosensitivity, myopathy (long- term therapy). Psychosis, seizures, leucopenia and rarely aplastic anaemia, hepatitis.
5	PRIMAQUINE PHOSPHATE	MALIRID - DS	Tablet	15 - 30 mg/day	40.00	Nausea, vomiting, epigastricdistress abdominal cramps, leukopenia, leucocytosis, agranulocytosis, methaemoglobinemia in NADH methaemoglobin reductase-deficient individuals.

### ANTI - PARASITIC

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ALBENDAZOLE	BANDY	Tablet	Child: 15mg/kg daily in 2 divided doses.Max.800mg/day.	16.20	skin rashes, swelling, nausea, vomiting headache, drowsiness, dizziness, and diarrhea.
		ZENTEL	Syrup		16.21	vision changes, yellowing eyes or skin, severe stomach or abdominal pain.
2	DIETHYL CARBAMAZINE CITRATE	HETRAZAN - 100mg	Tablet	Filariasis; Loiasis; Toxocariasis Inital: 1 mg/kg/day, may increase to 6mg/ /kg/day.	52.50	Fever, headache, vomiting, dizziness, drowsiness, nausea, chills.
3	IVERMECTIN	IVERFLAR-12mg TABLET	Tablet	Single dose.	80.00	Diarrhoea, nausea, vomiting, dizziness, pruritus, urticaria. rash. arthralgia. fever. myalgia.

		IVERMECTOL - 12mg	Tablet		59.50	asthenia, postural hypotension, tachycardia, oedema, lymphadenopathy, sore throat, cough, headache, somnolence, transient eosinophilia, raised liver enzyme values.
		IVERMECTOL - 6mg			35.50	
		VERMACT-12mg TABLET	Tablet		372.50	
4	IVERMECTIN + ALBENDAZOLE	BANDY PLUS - 12mg	Tablet	At once.	30.00	Nausea, Vomiting, Diarrhea, Loss of appetite, Increased liver enzymes, Itching, Dizziness, Swelling of lymph nodes.

### ANTI-PARKINSONISM

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
2	CITICHOLINE	CITINERVE - 500mg	Tablet, injection	PO Parkinson's disease; Cerebrovascular disorders; Head injury 200-600 mg/day in divided doses. IV/IM Parkinson's disease;	400.00	Head ache, Diarrhoea.
		CITILIN-250MG - 2ml			106.75	
3	DONEPEZIL	DONECEPT - 5mg	Tablet	5 mg/day in the evening, up to 10 mg/day after 4-6 wk.	94.50	Nausea, vomiting, anorexia, wt loss diarrhoea, insomnia, fatigue, muscle cramps; headache and dizziness; syncope, bradycardia; convulsions;
4	LEVODOPA& CARBIDOPA	SYNDOPA - 110mg	Tablet	Adult: start with 1 tablet ( carb: 10mg, leva: 100mg) TID increase gradually	17.51	GI upset, postural hypotension, cardiac arrhythmia
		SYNDOPA PLUS			23.73	
6	PRAMIPEXOLE	PRAMIPEX-ER-0.375mg	Tablet	0.125 mg PO q8hr initially; gradually titrated upward at	82.00	Dizziness, lightheadedness, drowsiness.
7	SELEGILINE HCL	SELGIN - 5mg	Tablet	Adult: 10mg /day at	46.50	GI upset, postural hypotension, cardiac arrhythmia.
8	TRIHENXYPHENIDYL HCL	PACITANE - 2mg	Tablet	Adult: 2-10 mg/day	47.26	Dryness of mouth, impairment of memory, confusion.

### ANTI-PSYCHOTICS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMISULPRIDE	AMISANT - 200mg	Tablet	Adult: PO 400-800 mg bid. Max: 1.2 g/day.	118.00	Insomnia, wt gain, QT prolongation, hypotension, bradycardia, GI disorders hyperglycaemia; breast pain, gynaecomastia, galactorrhoea.
		SULPITAC - 100mg			87.00	

2	ARIPIRAZOLE	ARPIZOL-2mg	Tablet	Adult: 10 mg or 15 mg once daily	4.90	GI disorders, headache, anxiety, wt gain, insomnia, lightheadedness, drowsiness.
		ARPIZOL - 15mg		Adult: 10 mg or 15 mg once daily	116.00	GI disorders, headache, anxiety, wt gain, insomnia, lightheadedness, drowsiness.
		ARPIZOL-10mg			103.00	
		ARIPAXE-10mg	Tablet	Once daily	82.00	Akathisia (inability to stay still), Anxiety, Blurred vision, Constipation, Dizziness, Fatigue.
		ARIPAXE-15mg	Tablet		111.00	
3	ACAMPROSTATE	ACAMPROL - 333 MG	Tablet	PO $\geq$ 60 kg: 666 mg 3 times/day; <60 kg: 666 mg at breakfast, 333 mg at midday and 333 mg at night. Usual duration: 1 yr.	57.00	GI disturbances (diarrhoea, nausea, vomiting, abdominal pain, asthenia), dermatological (pruritus, rash).
4	CARIPRAZINE	CARISPEC-1.5mg	Tablet	Once daily.	99.00	Dizziness, lightheadedness, drowsiness, nausea, tiredness, excess saliva/drooling, blurred vision, weight gain, or constipation.
5	CLOZAPINE	SIZOPIN -50mg	Tablet	Adult:50-300mg /Day	53.00	Agranulocytosis, blood dyscrasis, metabolic syndrome.
		SIZOPIN - 25mg			32.00	
6	CHLOPROMAZINE	CHLOPROMAZINE-50mg TABLET	Tablet	Adult:25mg tid;may be given as a single 75mg dose at night.Maintenance: 25-100mg tid increased to $\geq$ 1g daily as required in psychotic patients. Child:1-12 yr:500mcg/kg every 4-6hr.Max:>5yr:75mg daily;1-5yr:40mg daily. Elderly:Initially,1/3-1/2 the normal adult dose.	4.85	Drowsiness, dizziness, lightheadedness, dry mouth, blurred vision, tiredness, nausea, constipation, weight gain, and trouble sleeping.
7	HALOPERIDOL	SERENACE 0.25mg	Tablet, Injection	Adult:2-20 mg/day	11.95	Parkinsonism, Autonomic effects.
		SERENACE - 1ml			6.52	
8	LITHIUM CARBONATE	LITHIC SR - 400mg	Tablet	Adult:600-1200mg/day	39.00	Thirst, polyurea, tremors, seizures CNS toxicity, GI upset, renal diabetes insipidus, goitre.
		LITHIC SR - 450mg				
9	LURASIDONE	LURAMAX-80mg	Tablet	Once daily	200.00	Parkinsonism, Akathisia (inability to stay still), Anxiety, Stomach discomfort, Restlessness, Upper abdominal pain Agitation.
10	MODAFINIL	MODALERT-200mg	Tablet	Once daily	269.00	Irritability, Dyspepsia, Increased Liver Enzymes, Abnormal Thoughts Depression, Tachycardia.

11	OLANZAPINE	OLEANZ - 10mg	Tablet	Adult:2.5-10 mg/day	61.67	Agranulocytosis, blood dyscrasis, metabolic syndrome.
		OLEANZ - 5mg			32.40	
		OZAPINE Z-10mg			74.00	
12	QUETIAPINE	QUTIPIN - 50mg	Tablet	Adult:50-400 mg/day	80.00	Sedation, postural hypotension, urinary incontinence.
13	RISPERIDONE	RESPIDON-2mg	Tablet	Adult:2.-12 mg/day	30.00	Increase in prolactin levels, postural hypotension, parkinsonism.
		RISDONE-60ml	Syrup	Adult:2.-12 mg/day	117.60	Increase in prolactin levels , postural hypotension, parkinsonism.
14	ZUCLOPENTHIXOL DECANOATE	CLOPIXOL DEPOT-200 - 1ml	Injection	IM Acute psychosis,As acetate ester: 150 mg,repeat if needed 2-3 days later. Max cumulative dose: 400 mg/course.Max number of inj: 4/course Max duration:2wk.	370.40	Drowsiness, blurred vision, tachycardia, nausea, dizziness, headache, excitement, postural hypotension, hyperprolactinaemia, early phase of treatment; urinary frequency or incontinence; tardive dyskinesia.
15	ZUCLOPENTHIXOL ACETATE	ZUTHIXOL 50mg/1ml (SHORT ACTING)	Injection	50-150mg repeated if necessary after 2 or 3 days.	475.00	Dizziness, lightheadedness, drowsiness, constipation, headache, dry mouth, blurred vision.
16	ZUCLOPENTHIXOL DECANOATE	ZUTHIXOL (LONG ACTING) 200MG/1ML	Injection	200 - 500 mg every one to four weeks.	999.00	

### ANTI-TUSSIVES & EXPECTORANTS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ACEBROPHYLLINE ACETYLCYSTEINE	PULMOCLEAR - 100ml	Syrup	Two times a day.	127.00	Drowsiness, Dizziness, Skin rash, Stomach pain or discomfort, Vomiting.
2	AMBROXOL+ TERBUTALINE + GUAIPHERNESIN	MUCOLITE	Tablet, Syrup, Drops.	Ambroxol: 15 ,mg, Guaiphenesin 50mg, Terbutaline: 1.25 mg.	25.00	Mild GI - effects, Allergic reaction.
		MUCOLITE - 100ml			59.00	
		MUCOLITE SR - 75mg			62.50	
		MUCOLITE - 15ml			19.60	
		AERODIL - 100ml			68.00	
		CHESTON PLUS - 100ml			48.50	
3	AMBROXOL+GUAIFENESIN + LEVOSALBUTAMOL	CHERICOF-LS-100ml	syrup	Adult: 30-120mg daily in 2 -3 divided doses.	71.00	Nausea, Vomiting, Diarrhoea, Stomach upset, Stomach pain, Allergic reaction, Dizziness, Headache, Rash, Urticaria, Tremor, Palpitations, Muscle cramp, Increased heart rate.
		ASCORIL-LS-SYRUP			90.00	

4	AMYL METACRESOL + DEXTROMETHORPHAN	TUSQ-D-LOZENGES	Lozenges	Adult: 1-2 lozenges every 4-6 hrs. Child: 6-12 yrs: 1 lozenge every 4-6 hrs.	39.00	Stomach upset, Sore tongue.
5	BROMHEXINE + HYDROCHLORIDE	BROMHEXINE-8mg	Tablet	Adult & child ≥12 yrs: 8-16mg tid. Child:(2-5yr) 8 mg daily in 2-3 divided doses;(6-11 yr): 4-8 mg tid	45.90	Feeling of fullness in the stomach (bloating), diarrhea, dizziness, headache, indigestion, nausea, sweating and skin rashes.
		BROMHEXINE - 100ml	Tablet, Syrup	Adult 4-8 mg tablet Adults and Children >12 years: 10 mL (2 tsp) 3 times a day. Children 6-12 years: 5 mL (1 tsp) 3 times a day; 2-5 years: 2.5 mL (½ tsp) 3 times a day; < 2 years: Not recommended.	74.00	Headache, Sweating, Skin rashes, drowsiness and dizziness.
		ASCORIL EXPECTORANT - 100ml	Expectorant		76.00	
6	BROMHEXINE + TERBUTALINE	BRONCOREX-50ml	Syrup	Bromhexine-4 mg,two-four times a day Terbutaline Sulphate-1.25 mg,three times per day. Guaiphenesin-50 mg,200 to 400 mg 4 hourly.	57.15	Nausea, Diarrhoea, Bloating, Indigestion, Vomiting, Stomach pain, Sweating, Dizziness, Headache, Skin rash, Urticaria, Tremor, Increased heart rate.
7	BROMHEXINE + GUAIFENESIN + MENTHOL + TERBUTALINE	BRO-ZEDEX-100ml	Syrup	Adults -10ml 2-3 times/day, 3 – 6 years 2.5 ml TID, 6-12 years 5 ml TID	98.00	Nausea, diarrhoea rash, angioedema, pruritus, urticaria, sweating, dizziness, vomiting, headache and pain in the upper part of abdomen.
8	CHLORPHENIRAMINE MALEATE + DEXTROMETHORPHAN	ASCORIL-D -100ml	Syrup	Three times a day	120.00	dizziness, drowsiness; dry mouth, nose, or throat; constipation; blurred vision; or. feeling restless or excited (especially in a child).
9	CHLORPHENIRAMINE MALEATE + DEXTROMETHORPHAN HYDROBROMIDE + PHENYLEPHRINE HYDROCHLORIDE	CHERICOF	syrup	Child:2-6yrs-1/2tsp;6-12-1tsp;Adult-2tsp three to four times/day	76.50	Nausea, Vomiting, Loss of appetite, Headache.
10	CHLORPHENIRAMINE MALEATE + LEVODROPROPIZINE	RESWAS	Syrup	CHLORPHENIRAMINE(2mg/5 ml): 2 times / day; LEVODROPROPIZINE(30mg/5 ml): Adult and children>12 yrs:60 mg 3 times/day;Child: >2 yr: 1 mg/kg tid for ≤7 days.	63.00	Dizziness, GI disturbance

11	DEXTROMETHROPHAN	ALEX COUGH-LOZENGES	Lozenges	Adult & child >12yrs: As loz: suck not more than 10 loz daily Child: As loz(6-12 yr) suck not more than 2 loz within any 4 hr, Max:7 loz daily.	78.00	Sleepiness, Dizziness, Excitation, Confusion, Convulsion, Respiratory depression, Nausea, Vomiting, Diarrhoea, Rash.
12	LEVODROPROPIZINE	RAPITUS-100ML SYRUP	Syrup	(30mg/5ml): Adult and children>12 yrs:60 mg 3 times/day;Child: >2 yr: 1 mg/kg tid for ≤7 days.	90.00	Dizziness, GI disturbance.
13	MENTHOL + CAMPHOR + TERPINEOL + EUCALYPTOL + CHLOROTHYMOL	KARVOL PLUS-CAPSULE	Capsule	Snip tip of cap and squeeze contents into half ltr hot or boiling water or on a tissue/cloth handkerchief and inhale vapours deeply and freely. At night, the inhalant can be applied on the pillow or night dress.	57.30	Nasal irritation, Dryness in the nasal cavity, Sore throat, Burning, Drowsiness, Vomiting, Abdominal pain, Coma.
14	TERBUTALINE + GUAIPHENESIN	BRO - ZEDEX - 100 ML	syrup	Adults -10ml 2-3 times/day, 3 – 6 years 2.5 ml TID, 6-12 years 5 ml TID	75.00	Nausea, diarrhoea rash, angioedema, pruritus, urticaria, sweating, dizziness, vomiting, headache and pain in the upper part of abdomen.
		ASTHAKIND-100ml EXPECTORANT	Expectorant		58.56	

### ANTI-ULCER

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	AMOXYCILLIN, CLARITHROMYCIN & ESOMEPRAZOLE 40 MG	SOMPRAZ-HP KIT(6S)	Tablet	Two times or Three times a day	192.00	Headache, Dizziness, Dryness in mouth, Nausea, Abdominal ache, Constipation, Flatulence.
2	ANTACID GEL	DIGENE GEL - 200ml	Tablets,	" Chew 1 -2 tablets four times a day Syrup:Shake bottle before use. Should be taken undiluted, but the suspension may be followed by a sip of water if	81.90	GI upset, headache, dry mouth confusion, drowsiness
		DIGENE		desired. – 10 to 20ml four times daily – should be taken 15 minutes before meals and at bedtime – do not exceed recommended dosage."	12.00	
		MUCAINE - 200ml			221.00	
3	CINTAPRIDE & PANTOPRAZOLE	CINTODAC	Capsules	Once daily	393.05	Headache, Diarrhea, Stomach pain, Flatulence, Dizziness.
		PAN D	Capsules	Single capsule a day before breakfast in the morning	210.00	Diarrhoea, Dry mouth, Headache, Weakness, Nausea, Vomiting.

4	DOMPERIDONE & ESOMEPRAZOLE	SOMPRAZ-D-20mg m	Capsules	usually 1 to 2 times daily.	8.30	Nausea, bloating, or gas may occur headache, diarrhea, nausea, flatulence, stomachpain, constipation, dry mouth
5	ALUMINIUM HYDROXIDE 300 MG+DIMETHICONE 40 MG+MAGNESIUM HYDROXIDE 250 MG+SORBITOL 1000 MG /5ML	DIOVOL-170ml SYRUP	Syrup	Take 1 to 3 hours after meals as an antacid.	95.00	Bad taste in your mouth. Hard stools.
6	ESOMEPRAZOLE	RACIPER- 40mg	Tablets,	Adult:20-40 mg OD	63.50	GI upset, headache, dry mouth confusion, drowsiness.
		RACIPER-20mg	Capsules		37.50	
		ESOZ - 20mg			64.50	
		ESOZ - 40mg			102.50	
		ESOZOLE ZZ-40mg			111.00	
7	ENTRIC COATED ESOMEPRAZOLE MAGNESIUM AND DOMPERIDONE SR CAPSULES	SOMPRAZ-D - 40mg	Capsule	20 - 40 mg OD	9.90	Increased risk of Clostridium difficile- associated diarrhoea (CDAD) and osteoporosis-related fracture. Headache, rash, pruritus, dizziness, fatigue, cough, back or abdominal pain, arthralgia.
		RACIPER - D			98.00	
8	ESOMEPRAZOLE + LEVOSULPRIDE	SOMPRAZ-L	Capsule	Once Daily	165.00	Diarrhea, nausea, vomiting, stomach pain, flatulence, constipation, and headache.
9	FAMOTIDINE	FAMOTIDINE - 20mg	Tablet	Adult:20-40 mg BD	2.30	GI upset, hypotension, tachycardia, rash.
10	ITOPRIDE HCL	GANATON	Tablet	Adult: PO Antiemetic; As a prokinetic agent 50 mg 3 times/day.	86.50	Rash, diarrhoea, constipation, abdominal pain, headache, sleeping disorders, dizziness, galactorrhea, gynecomastia.
11	LANSOPRAZOLE	JUNIOR LANZOL-15mg	Tablet	1-11 yr ≤30 kg: 15 mg once daily in the morning for up to 12 wk; >30 kg: 30 mg once daily in the morning for up to 12 wk. May increase doses up to 30 mg bid if patient is still more wk of treatment. 12-17symptomatic after 2 or yr 15 mg once daily for up to 8 wk.	56.00	Diarrhoea, abdominal pain, nausea, vomiting, flatulence, constipation, headache, drymouth, peripheral, oedema, dizziness, sleep disturbances, fatigue, paraesthesia, arthralgia, myalgia, rash, pruritus.
		JUNIOR LANZOL-30mg			119.00	
12	LEVOSULPIRIDE & RABEPRAZOLE	RAZO-L TABLET	Tablet	Twice a day	364.75	Nausea, Stomach pain, Diarrhea, Flatulence, Constipation.
13	MAGALDRATE+OXETACAIN+SIMETHICONE	RANIDOM-O-200ml	Syrup	1-2 teaspoon full	119.16	Constipation, diarrhea and intestinal pain.
14	MESALAMINE	MESACOL - 400mg	Tablet	1-4 g/day BD/TDS	73.45	Abdominal pain (if new abdominal pain - consider pancreatitis); headache, nausea; flu; fatigue; fever, rash; sore throat; diarrhoea, joint pain.



15	OMEPRAZOLE + TINIDAZOLE +	HP-KIT	Capsule	omeprazole 20 mg, amoxicillin 750 mg, tinidazole 500 mg; 1 kit		Nausea, Vomiting, Stomach pain, Flatulence, Indigestion, Dizziness, Tiredness, Headache,
16	OMEPRAZOLE SODIUM	OCID-20mg	Capsule	Once per day; Adult:20-40 mg OD; Paed: 0.4-0.8 mg/kg/day	55.62	Headache,Diarrhoea,Stomach pain, flatulence Constipation,Nausea or vomiting.
		OMEZ - 20mg	Capsules,		52.35	GI upset, headache, dry mouth confusion, drowsiness.
		OMEZ	Injection		29.80	
17	PANTOPRAZOLE SODIUM	ZEPOXIN IV-40mg	Injection	Twice a day	53.00	Skin Rash,Constipation.
		PANTAKIND-40MG	Injection		55.00	Headache, Altered sense of taste Fast or slow or irregular heartbeat Diarrhoea, Muscle pain.
		PAN - 40mg	Tablet	40 mg orally once daily 40 mg via IV infusion once a day, given over at least 2 minutes OR over 15 minutes Adult:40 mg OD	80.00	GI upset, headache, dry mouth confusion, drowsiness.
		PAN 20mg	Injection		52.00	
		PANTOCID - 40mg			79.00	
		PANZO-40mg			8.80	
		ZEPOXIN-40mg			46.00	
		PANTOCID-80mg				
PANTOCID I.V-40mg		49.95			Headache, diarrhoea, nausea, abdominal pain, vomiting, flatulence, dizziness, and arthralgia.	
18	PANTOPRAZOLE SODIUM + DOMPERIDONE	PANTOCID DSR	Tablet, Capsules	Adult:40 mg OD	119.00	GI upset, headache, dry mouth confusion, drowsiness
		PANTOCID - D	Injection		77.00	
		PANDOZ DSR TABLET	Tablet		118.70	
19	PANTOPRAZOLE SODIUM LEVOSULPRIDE	PANTOCID-L	Tablet		181.00	
20	RABEPRAZOLE	RAB ZE-20mg	Tablet	Adult:20 mg OD , 20 mg orally 2 times a day	7.03	GI upset, headache, dry mouth confusion, drowsiness.
		ROLES - 20mg	Tablet,	Adult:20 mg OD	59.00	
		RAZO - 20mg			83.00	
21	RABEPRAZOLE + DOMPERIDONE	RAB Z DSR	Tablet	Usually 1 to 2 times daily	9.30	Hoarseness. light-colored stools. rapid weight gain. tingling of the hands or feet.
		ROLES-D	Capsule		77.50	
22	RABEPRAZOLE + LEVOSULPRIDE	ROLES-L	Capsule	Once Daily	179.50	Headache,Nausea,Vomiting Flatulence.
23	RANITIDINE HCL	RANTAC - 150mg	Tablet,	Adult: 150-300mg OD-BD; Injection:50-100mg every 6-8 hours	0.72	GI upset, hypotension, tachycardia, rash.
		RANITIN - 2ml	Injection	Paed: 1-5 mg?kg?dayin 2-4 divided dose njection:1-5mg every 6-8 hours	3.38	
		RANITIN 150mg		17.52		

24	SODIUM BICARBONATE	UDOSIS-500mg	Tablet	Parenteral: 50 to 150 mEq sodium bicarbonate diluted in 1 L of D5W to be intravenously infused at a rate of 1 to 1.5 L/hour. Oral: 325 to 2000 mg orally 1 to 4 times a day. One gram provides 11.9 mEq (mmoL) each of sodium and bicarbonate.	3.00	Headache (continuing) loss of appetite (continuing) mood or mental changes muscle pain or twitching nausea or vomiting.
25	SUCRALFATE	SUCRAFIL - 200ml	Syrup	Adult 1g TID	158.00	Flatulance, Abdominal fullness, GI upset.
		SUCRAFIL - 450ml			370.00	
		SUCRAFIL - O 200ml			195.00	
		SRFIL ENEMA-60ml	Enema	1-2 gm tablet mixed with 4.5-10ml of water in an enema applicator producing a low volume paste. Can be self-administered twice daily for 6 months.	95.00	Rectal bleeding, gas, bloating, diarrhea, stomach cramps, and rare allergic reactions, such as hives and difficulty breathing.
	SUCRA ZZ FIL 200ml	Syrup	2 teaspoonfuls 4 times daily, on an empty stomach at least 1 hour before a meal.	149.50	Dry mouth, upset stomach, gas, and nausea Frequent urge to urinate.	
26	SUCRALFATE + OXETACAINE / OXETHAZAINE	SUCRAFIL-O-200ml	Syrup	Adult: 1 g four times per day on an empty stomach.	213.00	Constipation, Allergic reaction.
27	SUCRALFATE & OXETACAINE	ULCOAT-O-100ml	Suspension	One to two times a day	120.00	Constipation. Dizziness. Sleepiness. Dry Mouth. Nausea.

### ANTI-VIRAL

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ACICLOVIR	ACIVIR DT - 200mg	Tablets,	Adult: 200-800 mg five times daily; Paed: 5-10 mg/kg 5 times daily	81.50	GI upset, increase in liver enzymes, burning, itching, haematological changes, rashes.
		ACIVIR DT - 400mg			67.35	
		ACIVIR DT - 800mg			160.00	
		NEOCLOVIR - 500mg			450.00	
2	AMANTADINE HYDROCHLORIDE	AMANTRIL-100mg	Tablet	Two times a day.	148.20	Dry mouth, constipation, nausea, vomiting, decreased appetite, difficulty falling asleep or staying asleep, abnormal dreams, headache.

3	GANCICLOVIR	VIRSON GEL	Gel	five times a day	111.00	Chills, headache, shortness of breath diarrhea, abdominal and stomach pain, nausea and vomiting, back pain seizures, swelling of legs, itching, decreased urine output.
4	OSELTAMIVIR PHOSPHATE	ANTIFLU-75mg	Capsule	Twice a day for 5 days	726.72	Nausea, vomiting, stomach pain, and diarrhea.
5	REMDESIVIR	REDYX-L100mg	Injection	Once daily for 5 to 10 days	2700.00	Low blood pressure, nausea, vomiting, sweating, and shivering.
		COVIFOR-100mg			5400.00	
		REMDESIVIR-100mg			1568.00	
6	TENOFVIR DISOPROXIL FEMARATE	TENOF-300mg	Tablet	300mg-once daily    paed: 8mg/kg daily	1325.60	Nausea, stomach pain, diarrhea, depression, headache, dizziness, weakness, trouble sleeping, itching or rash.
		TERAVIR-300mg	Tablet	300mg-once daily    paed: 8mg/kg daily	43.67	Nausea, stomach pain, diarrhea, depression, headache, dizziness, weakness, trouble sleeping, itching or rash.
		TENVIR - 300mg	Tablets	Adult>18 years: 300mg OD	1500.00	Flatulence, peripheral neuropathy, insomnia, body fat redistribution, raised liver enzymes.
7	VALCYCLOVIR HCL	VALTOVAL-1gm	Tablets	500mg-2G tid	157.00	Acute renal failure, CNS effects Potentially Fatal: Thrombotic thrombocytopenic purpura/haemolytic uraemic syndrome in patients w/ advanced HIV disease. Acute renal failure, CNS effects Potentially Fatal: Thrombotic thrombocytopenic purpura/haemolytic uraemic syndrome in patients w/ advanced HIV disease.

### CNS - OTHERS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	CAFFEINE CITRATE	APNICAFA-1ml	Injection	Once	280.06	Feeding intolerance Agitation Chills Coma Confusion Depression.
		CAFNEON-20mg/1ml		At once.	280.00	Delirium (acute confusional state) Insomnia (difficulty in sleeping) Frequent urge to urinate Dehydration Fever.

2	EDARAVONE	ARAVON-20ml	Injection	Initial treatment cycle: Administer 60 mg as an IV infusion over 60 minutes qDay for 14 days, followed by a 14-day drug-free period Subsequent treatment cycles: Administer 60 mg as an IV infusion qDay over 60 minutes on 10 of 14 days, followed by a 14-day drug-free period	580.00	Headache red, itchy, or scaly rash
		EDANOX-20ml		Once daily .	691.00	Bruising, Gait disturbance, Headache, Dermatitis, Eczema, Respiratory failure, respiratory ,disorder, low blood oxygen (hypoxia), Glucose in the urine, Tinea infection.
3	ERGOTAMINE, CAFFEINE, PARACETAMOL, PROCHLOPRAZINE	VASOGRain	Tablet	1-2 tabs as per the clinical condition needed.	66.50	Muscle cramps, stiffness, tiredness. Numbness and tingling of extremities Nausea, vomiting, diarrhoea. Ergotism consisting of hypotension gangrene of extremities, anginal pain.
4	PIRACETAM	NOOTROPHIL	Tablet, Injection	PO 2.4 to 7. 2 g/day in 2-3 divided doses  IV/IM 1-2 g 3 times/day.	449.47	Hyperkinesia, nervousness, depression, diarrhoea, rashes. CNS stimulation, sleep disturbances, dizziness, excitement, insomnia, somnolence, wt gain.
		NOOTROPIL - 100mg			231.73	
		NOOTROPIL 15ml - 200mg			118.75	
5	PIRACETAM + GINKGO BILOBA + VINPOCETIN	COGNIVEL			174.40	
6	PIRACETAM & CITICOLINE	STROCIT PLUS	Tablet	Once or twice a day	799.00	Sleeplessness, headache, diarrhea, low or high blood pressure, nausea, blurred vision, chest pain.
		CITISTAR-PM	Tablet		885.90	
7	PRAMIPEXOLE DIHYDROCHLORIDE	PRAMIPEX-0.25mg	Tablet	Three times a day.	7.30	Nausea, dizziness, drowsiness, lightheadedness, trouble sleeping, constipation, headache, or dry mouth may occur.
		PRAMIPEX-0.5mg			11.60	
		PRAMIPEX-1mg			16.00	
		PRAMIPEX-0.125mg			4.70	
8	PREGABALIN (75MG) + NORTRIPTYLINE (10MG)	PREGADOC-NT	Tablet	150 mg/day, may increase to 300 mg/day after 3-7 days. Max: 600 mg/day after a 7-day interval.	159.42	Constipation, Difficulty in urination, Weight gain, Sleepiness, Dizziness, Tiredness, Blurred vision, Dryness in mouth, Increased heart rate, Orthostatic hypotension (sudden lowering of blood pressure on standing), Uncoordinated body movements.

		PREGALIFT-NT		Initial: 150 mg/day, may increase to 300 mg/day after a wk. Max: 600 mg/day.	143.00	Constipation, Difficulty in urination, Weight gain, Sleepiness, Dizziness, Tiredness, Blurred vision, Dryness in mouth, Increased heart rate, Orthostatic hypotension (sudden lowering of blood pressure on standing), Uncoordinated body movements.
9	PYRIDOXINE +CYANOCOBALAMIN+NICOTINAMIDE+THIAMINE +RIBOFLAVIN +PANTHENOL	NEUROBION FORTE-2ml	Injection	500 mcg/day 3 times/wk.	11.22	Itching, Diarrhea, Dizziness, Irritable, Sleepiness, Rash, Eczema, Tingling or numbness, Flushing, Increased blood sugar level.
10	RASAGILINE	RASALECT-1mg	Tablet	0.5 mg PO qDay initially, may increase to 1 mg/day if needed and tolerated.	14.00	Dizziness, drowsiness; sleep problems (insomnia), strange dreams; involuntary muscle movements; loss of appetite, weight loss; indigestion, stomach pain, nausea, vomiting, constipation; joint pain or stiffness; swelling in your hands or feet; dry mouth, cough.
		RASALECT-0.5mg			8.00	
11	ROPINIROLE	ROPARK-0.25mg	Tablet	With or without food, usually 3 times a day.	5.60	Nausea, vomiting, constipation, dizziness, drowsiness, weakness, unusual sweating, headache, and dry mouth may occur.
		ROPARK-0.5mg			8.10	
		ROPARK-1mg			12.50	
		ROPARK-2mg			18.00	
12	TOPIRAMATE	TOPIROL-25mg	Tablet	25-50 mg/day PO initially; increase by 25-50 mg/day at weekly intervals to 100-200 mg q12hr.	7.70	Somnolence, Fever, Cognitive problems Infection, Flushing.
		TOPIROL-50mg			12.20	
		TOPIROL-100mg			21.40	
13	TETRABENAZINE TABLETS	REVOCON	Tablet	12.5 mg PO qDay initially; after 1 week, the dose should be increased to 12.5 mg q12hr.	26.00	Fatigue, Depression, Dizziness, Headache, Shortness of breath, Anxiety.
14	GABAPENTIN, NORTRIPTYLINE	GABANTIN NT	Tablet	3 - 4 times a day or as single dose at night.	390.00	Dizziness, sleepiness, tiredness, headache, nausea, vomiting, blurred vision, dry mouth, dry eye.
15	GABAPENTIN, METHYLCOBALAMIN	GABANEURON-100mg	Tablet	Three times a day	8.14	Nausea and vomiting, Diarrhea, Headache, Skin rash, Chest pain and discomfort, Dizziness.
16	GABAPENTIN, METHYLCOBALAMIN AND ALPHA LIPOIC ACID.	TRIGABANTIN-100mg	Tablet	Initial: 300 mg on the 1st day, 300 mg bid on the 2nd day and 300 mg tid on the 3rd day; alternatively, 300 mg tid on the 1st day. Dose may increase in increments of 300 mg every 2-3 days.	127.00	Dizziness, Sleepiness, Tiredness, Uncoordinated body movements.

17	METHYLCOBALAMIN, AND PREGABALIN.	NEUGABA-M	Tablet	Initial: 150 mg/day, may increase to 300 mg/day after a wk. Max: 600 mg/day.	131.00	Dizziness, Sleepiness, Tiredness, Uncoordinated body movements.
		METHYLPRELIN ZZ	Tablet	Initial: 150 mg/day, may increase to 300 mg/day after a wk. Max: 600 mg/day.	161.00	Dizziness, Sleepiness, Tiredness, Uncoordinated body movements.
18	METHYLCOBALAMIN & ALPHA LIPOIC ACID,FOLIC ACID& PYRIDOXINE HCL	NUROKIND PLUS	Capsule	1-2 capsule /day	84.00	Nausea, Vomiting, Stomach ache, Loss of appetite, Flatulence, Skin rashes, Bloating.
19	METHYLCOBALAMIN+PYRIDOXINE + NICOTINAMIDE	NUROKIND PLUS	Injection	500 mcg/day 3 times/wk	22.00	Injection site related side effects, Flushing, Increased blood sugar level, Blurred vision, Rash, Elevations in liver function tests, Dizziness.
20	MECOBALAMIN,NIACINAMIDE,PYRIDOXINE	RE-XITE PLUS	Capsule	1500 mcg/day in 3 divided doses.	16.80	Constipation, diarrhea, or upset stomach
21	NIACINAMIDE(100 MG), THIAMINE(100 MG), PYRIDOXINE(100 MG),	RENERVE PLUS	Injection	500 mcg/day 3 times/wk	117.00	Pain/redness at the injection site, mild diarrhea, itching, or a feeling of swelling all over the body.
22	PREGABALINE & NORTRIPTYLINE	NOVOMINE-N	Tablet	Two to three times a day.	150.00	Constipation, Difficulty in urination, Weight gain, Sleepiness, Dizziness, Tiredness, Blurred vision, Dryness in mouth, Increased heart rate.
23	PIRACETAM+CITICOLINE	CITINERVE-P	Tablet	7.2 g/day in 2-3 divided doses; may increase by 4.8 g every 3-4 days. Max: 24 g/day in 2-3 divided doses.	540.00	Weight gain, Hyperactivity, Stomach pain, Nervousness, Diarrhoea, Decreased blood pressure.
		CITISTAR-PM			885.90	
24	PIRACETAM + GINKGO BILOBA + VINPOCETIN	COGNIVEL	Tablet	120-240 mg/day in divided doses.	216.00	Weight gain, Stomach upset, Rash, Flushing (sense of warmth in the face, ears, neck and trunk), Nervousness, Abnormality of voluntary movements.
25	BENFOTIAMINE 100 MG	BENALGIS - 100 MG TABLET	Tablet	1tab/ tid	65.85	Nausia, Vomiting, GI Irritation.

### CVS - OTHERS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ACETAZOLAMIDE	DIAMOX - 250mg	Tablet	Adult: 250- 1000 mg/day; Paed: 10 -15mg/kg/day in 3 divided doses	64.10	Malaise, fatigue, depression, excitement, headache, wt loss, GI disturbances, drowsiness, paraesthesia, diuresis, metabolic acidosis,
2	ADENOSINE	ADENOCOR	Injection	Adult: 6 mg IV bolus	255.00	Flushing, headache, palpitation, hypotension.
		ADNEON 6mg - 2ml		Paed: 0.04 - 0.1 mg/kg IV	239.00	

3	AMIODARONE	AMIODON 150mg - 3ml	Injection	Adult: 1-2.2g	71.00	Blue-grey discolouration of skin, photosensitivity, peripheral neuropathy, paraesthesia, myopathy, ataxia Blue-grey discolouration of skin, photosensitivity, peripheral neuropathy, paraesthesia, myopathy, ataxia.
		CORDARONE - 3ml			69.30	
		CORDARONE-100mg	Tablet	100-200mg/day	58.00	
		DURON-150mg/3ml	Injection	Adult: 1-2.2g	64.44	Tiredness, Blurred vision, Photosensitivity, Nausea, Coordination impaired, Peripheral neuropathy (tingling and numbness of feet and hand), Alterations in thyroid hormone levels, Altered taste, Constipation, Abnormal vision, Lung damage, Tremor.
		TACHYRA-100mg	Tablet	Once or Twice a Day	59.04	Blue-grey discolouration of skin, photosensitivity, peripheral neuropathy, paraesthesia, myopathy, ataxia Coughing. Constipation. Tremors. Photosensitivity. Sleep-related disorder Blue-grey discolouration of skin, photosensitivity, peripheral neuropathy, paraesthesia, myopathy, ataxia.
4	CHLORTHALIDONE	CTD - 6.25mg	Tablet	25-50 mg/day up to 100-200 mg/day	40.00	Dry mouth, thirst, weakness, lethargy, drowsiness, restlessness, seizures, oliguria, hypotension, fatigue, muscle cramps and GI disturbances.
5	DIGOXIN	LANOXIN - 0.25mg	Tablet	Rapid digitalisation: Loading dose: 0.75-1.5 mg in the 1st 24 hr Slow digitilisation: 250 mcg 1-2 times/day. Usual maintenance: 125-250 mcg/day.	14.57	Extra beats, anorexia, nausea and vomiting. Diarrhoea in elderly, confusion, dizziness, drowsiness, restlessness, nervousness, agitation and amnesia, visual disturbances.
6	DOBUTAMINE	DOBUTAMINE - 250mg - 5ml	Injection	Acute heart failure 2.5-10 mcg/kg, up to 0.5-40 mcg/kg according to response. Cardiac stress test As 1mg/mL soln: 5 mcg/kg/min for 8 mins, increase dose at 5 mcg/kg/min until 20 mcg/kg/min	295.60	Increased heart rate and BP, ectopic beats, palpitation. Nausea, headache, chest pain, palpitation, dyspnoea, paraesthesia, legcramps.
7	DOPAMINE	DOMIN - 200mg / 5ml	Injection	Initial: 1-5 mcg/kg/min, up to 5-10 mcg/kg/min according to patient's response. For seriously ill patients: up to 20-50 mcg/kg/min may be used.	30.80	Nausea, vomiting, tachycardia, ectopic beats, palpitation, anginal pain, hypotension, vasoconstriction, bradycardia, hypertension.

8	ESMOLOL	NEOTACH-100mg - 10ml	Injection	Loading dose (optional): 500mcg/ kg over 1 min. Maintenance: 50 mcg/kg/min for 4 min.	228.00	Hypotension, bradycardia, heart block, syncope, peripheral ischaemia, pallor, flushing nausea, vomiting, anorexia, dizziness, somnolence, paraesthesia.
		ESOCARD-10ml	Injection		220.00	Nausea, Headache, Fatigue, Constipation, Diarrhoea, Dizziness, Cold extremities.
9	ISOSORBIDE DINITRATE	ISOLAZINE - 20mg	Tablet	Adult: PO Long term management of angina 20-120 mg/day in divided doses. Max: 240 mg/day.Heart failure 30-160 mg/day in divided doses. Max:	6.55	Hypotension, tachycardia, flushing, headache dizziness, palpitation, syncope, confusion.
		SORBITRATE - 10mg			0.78	Nausea, vomiting, abdominal pain. Apprehension, restlessness, weakness and vertigo.
10	ISOSORBIDE MONONITRAT	NITROFIX - 10mg	Tablet	Adult: PO 20 mg 2-3 times/day.	21.40	Hypotension, tachycardia, flushing, dizziness, palpitation, syncope, Nausea, vomiting, abdominal pain, restlessness, weakness and
11	ISOXSUPRINE HYDROCHLORIDE	DUVADILAN	Tablet	10-20 mg 3-4 times/day	83.00	Hypotension, dizziness,palpitation vomiting, abdominal distress, severe rash,flushing, tachycardia.
12	IVABRADINE	IVABRAD - 5mg	Tablet	2.5 - 7.5 mg BD	141.00	Luminous phenomena (phosphen), blurred vision, bradycardia, 1st degree AV-block, ventricular extrasystole, supraventricular extrasystoles, palpitations, headache,dizziness, nausea, constipation, diarrhoea.
13	ISOSORBIDE DINITRATE	SORBITRATE-10mg	Tablet	10-60mg per 4-6hrs	0.78	Headache, dizziness, lightheadedness, nausea, and flushing.
14	LABETALOL	LABEBET-100mg	Tablet	Initially 100 mg twice daily, dose to be increased at intervals of 14 days; usual dose 200 mg twice daily, increased if necessary up to 800 mg daily in 2 divided doses, to be taken with food, higher doses to be given in 3-4 divided doses; maximum	136.00	Excessive tiredness, Tingling scalp or skin, Dizziness, Lightheadedness, Upset stomach, Stuffy nose, Fatigue.
		ALPHADOPA-L			12.20	Dizziness, Allergic reaction, Ejaculation disorder, Increased liver enzymes, Erectile dysfunction, Difficulty in urination.
15	METOLAZONE	ZYTANIX-2.5mg	Tablet	2.5-5 mg/day	119.20	Chest pain, palpitation, necrotising angitis, orthostatic hypotension, syncope, venous thrombosis, vertigo, volume depletion; depression.
16	NEBIVOLOL HCL	NEBILONG 5mg	Tablet	1.25-5 mg once	95.00	Headache, insomnia, chest pain, dyspnoea, rash, and peripheral oedema, hypercholesterolaemia, hyperuricaemia.
17	NICORANDIL	KORANDIL - 5mg	Tablet, Injection,	10-20 mg twice daily	107.00	Headache (usually transitory), flushing, dizziness, nausea, vomiting and weakness. Hypotension and reflex tachycardia at high doses.
		NIKORAN IV-2mg		0.2 mg/kg	94.05	
		NIKORAN-10mg		Twice daily	450.50	
18	NITROGLYCERIN	ANGIPLATT - 2.5mg	Tablet, Injection,	inj: 5-25 mcg/min. Usual: 10-	108.00	Facial flushing, dizziness, tachycardia, throbbing



		ANGIPLATT - 6.5mg	capsules	200mcg/min. Max: 400 mcg/min PO: Extended release Up to 12.8 mg 3 times/day.	135.00	headache and tolerance. Large doses can cause vomiting, restlessness, hypotension, syncope, rarely cyanosis and methaemoglobinaemia, impaired respiration, bradycardia.
		NITROCONTIN - 2.6mg			160.60	
		NITROCONTIN - 6.4mg			189.50	
		NITROGLYCERIN-5ml			41.00	
		NITROCIN-5ml	Injection	0.2 mg/hour to 0.4 mg/hour.	36.81	Blurred vision, Decreased blood pressure, Dizziness, Headache, Increased heart rate, Lightheadedness, Paresthesia (tingling or pricking sensation).
19	NORADRENALINE	NORAD - 2ml	Injection	8-12 mcg/min, up to 8-30 mcg/min in refractory shock. Adjust according to BP response. Maintenance: 2-4 mcg/min.	186.40	Hypertension, headache, bradycar peripheral ischaemia, arrhythmias, skin necrosis, dyspnoea, respiratory difficulty.
		ADRENOR - 2ml			173.00	
20	OXYBUTYNIN (5MG)	TROPEN-5mg	Tablet	5-10mg QID	125.00	Dry mouth, dry eyes, blurred vision, mild constipation, diarrhea, nausea, mild stomach pain or upset, dizziness, drowsiness, weakness, headache, sleep problems.
21	PENTOXIFYLLINE EX.R	TRENTAL - 400mg	Tablet	400 mg 3 times/day	38.81	Nausea, vomiting, dizziness, headache, flushing; angina, palpitations; occasional cardiac arrhythmias; hepatitis, jaundice.
22	POLIDOCANOL	ASKLEROL *3%-2ml	Injection	At once.	134.06	Site pain, irritation, discoloration and site clotting, itching, site warmth.
23	SACUBITRIL & VALSARTAN	CIDMUS-100mg	Tablet	Twice a day	1174.04	Dizziness, Increased potassium level in blood, Fatigue, Hypotension (low blood pressure).
		AZMARDA-50mg	Tablet	Once a day	1092.00	Dizziness, Light-headedness, Nausea, Headache, Fainting, Loose motions Cough.
24	SACUBITRIL 24MG & VALSARTAN 26MG	ARNEY-50mg	Tablet	Twice a day	250.50	Cough, Dizziness, Diarrhea, Fainting, Headache, Nausea, Stomach pain Vertigo (sensation of feeling off-balance) Weakness.
		ARNEY-100mg	Tablet		410.50	
25	SILDENAFIL	ASSURANS - 20mg	Tablet	Adult: PO 50 mg about 1 hr before sexual intercourse.	165.00	Headache, flushing, AV block, migraine, respiratory tract infection, angina pectoris, syncope, tachycardia, postural hypotension, MI, cerebral thrombosis.
26	SOLIFENACIN SUCCINATE + ISOSORBIDE DINITRATE TAMSULOSIN	VELTAM-S-(KIT)	Tablet	3 tablets once a day for 12 weeks	350.00	Constipation, dry mouth, hoarse voice, dry eyes, blurred vision, nausea.
27	TADALAFIL	PULMOPRESS - 20mg	Tablet	10 mg once daily	275.00	Headache, flushing, angina pectoris, respiratory tract infection, AV block, migraine, syncope, tachycardia, postural hypotension, MI, cerebral thrombosis.
		TAZZLE-20mg TABLET			53.80	
28	TOLVAPTAN	NATRISE - 30mg	Tablet	15-60 mgOD	756.00	Polyuria, xerostomia, increased thirst, and polyuria.

		RESODIUM-15mg	Tablet	Initial dose: 15 mg orally once a day, Maintenance dose: Increase the dose to 30 mg once a day, maximum of 60 mg once a day,	575.55	Headache, Dizziness, Frequent urge to urinate, Nocturia (increased urge to urinate during the night), Dryness in mouth, Fatigue, Polyuria, Diarrhoea, Increased thirst.
29	TRIMETAZIDINE	FLAVIDON-MR	Tablet	40-60 mg/day	99.50	GI disturbances, vomiting, nausea.
30	TADALAFIL + DAPOXETINE	UPHOLD	Tablet	1-2 tab/day	256.00	Nausea, Vomiting, Indigestion, Headache, Flushing (sense of warmth in the face, ears, neck and trunk), Dizziness, Nose bleed, Dryness in mouth.
31	TAMSULOSIN (0.4MG)	TAMMAX-0.4mg	Tablet	0.4mg once daily increased to 0.8mg	63.00	headache, dizziness, runny nose, diarrhea, cough, decreased semen, loss of energy or muscle strength, sore throat or trouble swallowing.
		TASULIN-0.4mg	Tablet		9.50	Retrograde ejaculation, Dizziness, Nasal congestion.
32	TAMSULOSIN + DUTASTERIDE	TAMGRESS-D-0.4mg	Capsule	0.4mg once daily increased to 0.8mg	180.00	Impotence, Decreased libido, Breast pain, Ejaculation disorder, Dizziness.
33	TRIMETAZIDINE	CARVIDON-MR	Tablet	40-60 mg/day		GI disturbances, vomiting, nausea.
34	TOLTERODINE (4MG)	TOLU-XR-4mg	Tablet	2mg bid	203.50	Dryness in mouth, Constipation, Headache, Dizziness, Sleepiness, Blurred vision, Dry skin.
35	VASOPRESSIN	CPRESSIN - P	Injection	Adult: IV Initial control of 20 u. variceal bleeding As argipressin: IM/SC Cranial diabetes insipidus As argipressin: 5-20 u 4 hrly.	167.00	Pallor, vomiting, nausea, belching, abdominal cramps, tremour, pounding headache, vertigo, fluid retention, hyponatraemia, hypersensitivity reaction.

### HORMONES & THEIR ANTAGONIST

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	BETAMETHASONE DIPROPIONATE	STERNON - 30ml	Tablet, injection	Adult: 0.5 - 9 mg; Paed: 17.5 mcg/kg ( Adrenocortical insufficiency)	75.00	Peptic ulceration, acne eruptions, glaucoma, osteoporosis.
		BETSONE - 1ml			4.24	
		BETNESOL - 1ml			4.40	
		BETNESOL			8.10	
2	CABERGOLINE	CABGOLIN - 0.25mg	Tablet	0.25 mg orally twice a week. Dosage may be increased by 0.25 mg twice a week.	132.00	Decrease in BP, dizziness, vertigo, headache, nausea, sleeplessness, abdominal pain, dyspepsia, gastritis, weakness, fatigue, constipation, vomiting, breast pain, hot flushes.
3	CARBIMAZOLE	NEO - MERCAZOLE - 10mg	Tablet	Adult: Initial: 15-40 mg/day. Maintenance: 5-15 mg/day. Paed: 1-2 mg/kg /day	397.95	Mild leucopaenia, sore throat, mouth ulcers, fever, bruising, malaise; rarely cholestatic jaundice; nephrotic syndrome

		NEO - MERCAZOLE - 5mg		1 tab. 1-2 mg/kg/day.	204.75	jaundice, hepatic syndrome.
4	CARBOPROST	CABOPROST-1ml	Injection	Initial: 250 mcg. Alternatively, initiate w/ test dose of 100 mcg. May be repeated at 1.5- to 3.5-hr intervals depending on uterine response. May be increased to 500 mcg if uterine contractility is inadequate. Max: 12 mg. Max duration: 2 days.	192.60	Severe pelvic pain, cramping, or vaginal bleeding, high fever, feeling light-headed or short of breath, severe nausea, vomiting, or diarrhea.
5	CETRORELIX ACETATE	SETROSIL	Injection	The usual dose in adults for inducing ovulation is about 0.25 mg given subcutaneously on either day 5 or day 6 of stimulation.	1240.00	Nausea,pain, swelling, redness,Itching,Irritation, Vomiting.
6	CHORIONIC GONADOTROPHIN 5000 I.U	LUPI-HCG -5000I.U	Injection	500-4,000 u 3 times/wk.	438.25	Headache, irritability, restlessness, fatigue, depression, oedema; precocious puberty, gynaecomastia
		OVITRELLE-250mcg/0.5ml	Injection	250 µg should be administered one day following the last dose of the follicle stimulating agent	1732.50	Flatulence, diarrhea, hiccup, ectopic pregnancy, breast pain, intermenstrual bleeding, vaginal hemorrhage, cervical lesion, leukorrhea, ovarian hyperstimulation, uterine disorders, vaginitis, vaginal discomfort, body pain, back pain.
7	CLOMIFENE	SIPHENE - 50mg	Tablet	50 mg/day for 5 days starting on the 5th day of menstrual cycle.	45.75	Ovarian enlargement; abdominal pain and bloating; blurred vision; hot flushes; breast discomfort; depression; multiple or ectopic pregnancies;
		SIPHENE-100mg			70.40	
		CCQ-100mg			186.00	
		CCQ-50mg			192.50	
8	CYPROTERONE & ETHINYLOESTRADIOL	KRIMSON-35	Tablet	Hirsutism 2 mg/day (w/ethinylestradiol)for 21days of the menstrual cycle. Palliative treatment for prostatic carcinoma Initial: 300 mg/day in 2-3 divided doses. Maintenance: 200-300 mg/day.	260.00	Inhibits spermatogenesis, reduces volume of ejaculate, causes infertility, produces abnormal spermatozoa, gynecomastia and enlargement of mammary glands.
9	COENZYME Q10, DEHYDROEPIANDROSTERONE (MICRONIZED) & MELATONIN	OVARES PLUS	Tablet	Heart failure Adult: 50-150 mg daily in 2-3 divided doses. Angina Adult: 150-600 mg daily in divided doses. Periodontal disease Adult: 25 mg bid.	47.00	Drowsiness, change in sleep pattern and headache

10	CONJUGATED ESTROGENS (0.625MG)	PREMARIN-0.625mg	Tablet	.45 mg/day, up to 1.25 mg/day. Attempt to discontinue medication at 3-6-mth intervals.	709.21	Headache, Vaginal bleeding, Fluid retention, Vomiting, Nausea, Stomach cramp, Breast pain, Hair loss.
11	DEFLAZOCORT	DEZERT - 6mg	Tablet	120 mg/day. Maintenance: 3-18 mg/day.	87.00	GI disturbances, musculoskeletal, endocrine, neuropsychiatric, ophthalmic, fluid and electrolyte disturbances; susceptible to infection, impaired healing.
		DFZ - 6mg			85.00	
12	DEXAMETHASONE	DECDAK - 0.5mg	Tablet, Injection	0.75-9 mg/day in 2-4 divided doses.	2.10	Peptic ulceration, acne eruptions, glaucoma, osteoporosis.
		DEXONA - 2ml			5.61	
		AURADEx-8mg	Tablet	6.00		
13	DIENOGEST	DINOGEST-2mg	Tablet	Once a day	602.00	Weight gain, Depressed mood Sleeping difficulties, Nervousness Loss of sex drive, mood changes Headache or migraine, Nausea and vomiting, Abdominal pain Swollen stomach, Acne or hair loss.
14	DINOPROSTONE	DINOFIRST-2mg	Tablet	500 mcg, may repeat 6 hr later if needed. Max: 1.5 mg/24 hr.	54.79	Hypersensitivity reactions, nausea, vomiting, diarrhoea, abdominal pain; flushing, shivering, headache, dizziness; hypertension; convulsions, ECG changes.
		PRIMIGYN - 0.5mg	Gel		259.00	
		CERVIPRIME - 30gm			260.50	
		DECDAN-0.5mg	Tablet	Four times a day	2.00	Weight gain, indigestion, headache, dizziness, rash, fluid retention (swelling in your hands or ankles), sleep problems, mild mood changes and nausea.
15	DYDROGESTERONE	DUPHASTAN-10mg	Tablet	Duphaston 10 mg tablets once or twice daily with or without food.	62.35	Depression, Mood swings, Difficulty in falling asleep, Headache, Dizziness, Drowsiness.
		DROGYNA	Tablet	Once daily.	550.00	Headache, Nausea, Breast pain.
16	DUTASTERIDE	DUTAS	Capsule	Once daily.	350.30	Chest pain and discomfort, Swelling of face and lips, Skin Rash, Abnormal ejaculation, Breast discharge, Headache.
17	ESTRADIOL VALERATE	PROGYNOVA - 1mg	Tablet	10-20 mg OD-TDS	178.00	GI disturbances, genitourinary changes, haematologic disorders, CV and CNS effects, endocrine and metabolic disorders, cholestatic jaundice, local skin reactions, chorea.
		PROGYNOVA-2mg	Tablet	1-2mg per day	384.00	Headache, Nausea, Breast tenderness, Stomach pain.
		EVALON - 15gm	Cream	1 applicator upto the ring /day for 2-3 weeks; then 1 applicator upto the ring twice a week	289.00	

18	EVENING PRIMOSE OIL	PRIMOSA-1gm	Capsule	1 cap /day	242.54	Headache, Diarrhea, Cephalalgia, Upset stomach, Nausea.
		PRIMOSA-500mg		2cap/day	194.81	
19	FLUDROCORTISONE ACETATE	FLORICOT	Tablet	50-300 mcg/day	115.00	Hypertension, sodium and water retention, potassium loss, dizziness, itching, skin rash, headache, convulsions, CHF, muscle weakness, hyperglycaemia.
20	FLUNARIZINE DIHYDROCHLORIDE	SIBELIUM - 10mg	Tablet	5-10 mg/day at bedtime	89.00	Drowsiness. Rarely wt gain, headache, depression, gastric pain, dry mouth, insomnia, extrapyramidal reactions, galactorrhoea.
		SIBELIUM - 5mg			52.50	
		MIGRANEX-10mg	Tablet		4.97	
21	FSH	SITRODIN-75 IU	Injection	The recommended dose is 150 IU/day for 5 days, increased up to 450 IU/day for up to 12 days.	1044.90	Headache, nausea, vomiting, mild stomach/abdominal pain, bloating, redness/pain at the injection site, or breast tenderness/pain may occur.
		FOLISURGE-150IU	Injection	75-150 IU FSH daily and is increased preferably by 37.5 or 75 IU at 7- or preferably 14-day intervals.	3300.00	Allergic reaction, headache, Swelling of breast in male.
22	FOLLITROPIN ALFA	GONALF-300IU	Injection	Two medicines each day for up to five weeks. may be increased every 7 or every 14 days by 37.5 to 75 IU	8020.00	Headache,nausea,vomiting, mild stomach/abdominal pain, pelvic pain or tenderness, bloating.
		GONAL-F-450IU	Injection		11957.00	
23	HUMAN NORMAL IMMOGLOBULIN	GLOBUCEL-10gm/100ml-IV	Injection	500 mg/kg/weeks	15202.95	Back pain,Chills,Flushing.
24	HUMAN MENOPAUSAL GONADOTROPHIN	HUMOG-75IU	Injection	75 IU of hMG was used as an initial dose and this was continued for the first 14 days because adequate follicular development was not achieved. The dose of hMG was subsequently increased to 150 IU for the next 7 days. After 22 days from the start of stimulation.	775.00	Bloating,Stomach pain,Pain in the lower abdomen,Nausea or Vomiting,Rapid weight gain.
25	ISOPHANE INSULIN	HUMAN INSULATARD	Injection	0.3-1 IU/kg/day; may be given in 1-2 divided doses.	149.40	Hypoglycemia, insulin resistance, lipodatrophy, hypokalaemia, blurred vision.
26	IBANDRONIC ACID + CALCIUM CARBONATE	IDROSFOS KIT-TABLET	Tablet	2 -6 mg	519.00	Fever; influenza-like syndrome; electrolyte disturbances;GI disturbances;musculoskeletal pain; headache; rash, allergic dermatitis; anaemia; decreased alkaline phosphatase.
		ROCKBON KIT			468.27	
27	LETROZOLE	LETROZ - 2.5mg	Tablet	2.5 mg once daily	172.00	Hot flushes, arthralgia, nausea, vomiting, fatigue, dizziness, headache, dyspepsia, constipation.

28	LEVONORGESTREL & ETHINYLOGESTRADIOL	OVRAL - L	Tablet	Adult:PO Monophasic combined OC: As tab containing levonogestrel 150-250 mcg and and ethinylestradiol 30 mcg: 1tab once daily. Triphasic combined OC: Levonogestrel 50-125mcg+ ethinylestradiol 30-40 mcg:1 tab once daily.	70.56	Menstrual irregularities; headache, dizziness; breast discomfort; gynaecomastia; depression; disturbance of appetite; wt changes; fluid retention; oedema; changes in libido; hair loss or hirsutism; GI disturbances.
29	LEVONORGESTREL	MIRENA IUS-52mg	IUD	Must be removed at the end of 5th year.	3509.00	Missed periods (amenorrhea), bleeding and spotting between periods, heavier bleeding during the first few weeks after device insertion, abdominal/pelvic pain, ovarian cysts, back pain, headache/migraine, nervousness, dizziness, nausea, vomiting, bloating, breast tenderness or pain, weight gain, changes in hair growth, acne, depression, changes in mood, loss of interest in sex, itching or skin rash, and puffiness in the face, hands, ankles, or feet.
30	MEDROXY PROGESTERONE ACETATE	MEPRATE - 10mg	Tablet	5 to 10 mg orally once a day	60.05	Depression, fluid retention. Fatigue, insomnia, dizziness, headache, nausea;breasttenderness; wt gain/loss, anorexia; cholestatic jaundice; pain at Inj site.
31	MEGESTEROL ACETATE	ENDACE-40mg	Tablet	40 mg 4 times daily or 160 mg once daily.	311.60	Hair loss, increased appetite, mild blood pressure increase, mood changes, nausea or vomiting, swelling of the face, ankles, or feet, weight gain.
32	METHYL PREDNISOLONE	IVEPRED -16mg	Tablet, Injection	2-60 mg/day in 1-4 divided doses	99.00	peptic ulceration, acne eruptions, glaucoma, osteoporosis.
		IVEPRED -8mg	Syrup, Drops		54.00	
		MEDROL - 4mg			41.00	
		IVEPRED -125mg			288.00	
		IVEPRED -40mg			150.00	
		PREDMET -4mg			33.50	
		DEPO-MEDROL - 2ml			1271.00	
		NEODROL - 500mg			270.00	
		SOLU - MEDROL - 1gm			1271.00	
		PREDNISOLONE - 10mg			15.98	
		PREDNISOLONE - 5mg			9.45	
		PREDMET - 10ml			18.90	
		KIDPRED - 60ml			27.17	

		NEODROL-40mg	Injection	10-500 mg/day	48.75	Nausea, vomiting, heartburn, headache, dizziness, trouble sleeping, appetite changes, increased sweating, acne, or pain/redness/swelling at the injection site.
		DEPOPRED-2ml	Injection	The range of initial doses is 0.11 to 1.6 mg/kg/day	67.12	Aggression or anger Decreased urine output Dizziness Headache.
33	MENOTROPHIN	HUMOG-150IU	Injection	Three equal doses of HUMOG HP are given on alternate days	1272.00	In female, a local reaction at the injection site, fever and arthralgia have been observed in rare cases. In the male, a combined treatment with HUMOG HP.
		GMH HP-150IU	Injection	Start within the initial 7 days of the menstrual cycle. 150 IU daily	1485.00	Headache Injection site pain Injection site allergic reaction Abdominal pain Abdominal swelling.
		MENOTAS HP-150IU	Injection	Production of follicular maturation:75 IU each of FSH and LH I.M. daily for 9 to 12 days Stimulation of spermatogenesis. Adults: After 5,000 USP units of CG I.M. three times weekly for 4 to 6 months of treatment; 1 ampule (75 IU FSH/LH) I.M. three times weekly (given with 2,000 USP units CG twice weekly) for at least 4 months	1745.00	Headache, malaise, dizziness, fever,nausea, vomiting, diarrhea, abdominal cramps, bloating.
		IVF-M 150IU	Injection	At once.	1342.00	Allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue. breathing problems. fever or chills.
34	MEDROXY PROGESTERONE ACETATE	EMILY IUS	Single insertion	20 µg per day	2664.00	Depression, fluid retention. Fatigue, insomnia, dizziness, headache, nausea;breasttenderness; wt gain/loss, anorexia; cholestatic jaundice; pain at Inj site.
35	METHYLPREDNISOLONE	NEODROL-1gm	Injection	4 to 32mgday. IV- 1gm every 6 – 8 weeks	871.00	Nausea. Sweating.
		IVEPRED-4mg	Tablet	Four times a day	59.00	Edema (swelling), Thinning of skin.
36	MISOPROSTOL	MISOPROST- 200mg	Tablet	200-800 mcg/day in 2-4 divided doses	87.00	Diarrhoea, abdominal pain, dyspepsia, constipation, flatulence, nausea and vomiting; abnormal vaginal bleeding, cramps, increased uterine contractility, headache.
		ZITOTEC - 200mcg			78.50	
		MISOPROST*-25mcg	Tablet		18.30	

37	MIFEPRISTONE + MISOPROSTOL	INSTA KIT	Tablet	Tablet of Mifepristone, followed by 4 tablets of Misoprostol after 1-3 days.	378.47	Dizziness. Headache. Feeling tired or weak. Belly pain. Upset stomach or throwing up. Many small meals, good mouth care, sucking hard.
		UNWANTED KIT	Tablet	600 mg as a single dose, followed by misoprostol 400 mcg orally or gemeprost 1 mg vaginally 36-48 hr later.	385.89	Nausea, Vomiting, Diarrhoea, Stomach cramp, Uterine contractions, Heavy menstrual periods.
38	NANDROLONE DECANOATE	DECA DURABOLIN - 25mg	Injection	25-100 mg 3-4 wkly	157.00	Male: Postpubertal: Acne, gynaecomastia, bladder irritability, priapism, insomnia, chills, decreased libido, hepatic dysfunction, nausea, diarrhoea, prostatic hyperplasia. Prepubertal: Acne, virilism, chills, insomnia, diarrhoea, hyperpigmentation, nausea. Female: Virilism, insomnia, iron deficiency anaemia, hepatic hypercalcaemia, nausea, diarrhoea, chills, dysfunction.
		DECA DURABOLIN - 50mg			256.00	
39	NORETHINDRONE ACETATE	REGESTRONE - 5mg	Tablet	2.5-10 mg/day	58.60	Mental depression, cholestatic jaundice, porphyria, epilepsy, migraine, headache, breast discomfort, dizziness, nausea and vomiting, changes in libido.
		PRIMOLUT - N			58.73	
40	OCTREOTIDE	OCTRIDE 50mcg - 1ml	Injection	25 mcg/hr for 48 hr	305.00	Local pain, stinging, tingling at site of inj; anorexia, nausea, vomiting, abdominal pain, bloating, flatulence, loose stools, steatorrhea.
41	OXYTOCIN	EVATOCIN - 1ml	Injection	10-40 u by infusion at a rate to control uterine atony.	17.97	Transient hypotension, reflex tachycardia; nasal irritation, rhinorrhoea, lachrymation.
		OXYTOCIN-1ml	Injection		17.79	
42	PROGESTERONE	SUSTEN - 100mg	Capsule	200 - 400 mg/day	108.00	GI disturbances, appetite/wt change, fluid retention, oedema, acne, skin rash, urticaria, depression, headache, fever, fatigue, breast changes, hirsutism, changes in libido, altered menstrual cycles or irregular menstrual bleeding (rare).
		SUSTEN - 200mg			201.00	
		SUSTEN-400mg	Capsule	Once daily in the evening or at bedtime	554.00	
		SUSTEN-SR-200mg	Capsule		415.00	
43	NATURAL MICRONISED PROGESTERONE	HYPERTON 6-5gm	Ointment	max 5mg per oz not exceeding 2oz per month	87.00	Fatigue, Sleepiness, Headache, Abdominal pain.
		AQSUSTEN-25mg/1ml INJECTION	Injection	5 to 10 mg IM once daily for 6 to 8 days	99.00	Depression, fluid retention. Fatigue, insomnia, dizziness, headache, nausea;breasttenderness; wt gain/loss, anorexia; cholestatic jaundice; pain at Inj site.
44	SOLUBLE INSULIN	HUMAN ACTRAPID	Injection	20 u, then 6 microgram/hr until blood glucose drops to 10 mmol/L, when the dose is given 2 hrly.	149.40	Hypoglycemia, insulin resistance, lipoatrophy, hypokalaemia, blurred.
45	SILODOSIN	AVOPROST	Tablet	Once daily.	272.00	Diarrhea, dizziness, low blood pressure, stuffy or runny nose.



46	TESTOSTERONE ENANTHATE	TESTOVIRON-250mg	Injection	Usually every 1 to 4 weeks.	311.60	Nausea, vomiting, headache, skin color changes, increased/decreased sexual interest, oily skin, hair loss, and acne may occur. Pain and redness at the injection site may also occur.
47	THYROXINE SODIUM	ELTROXIN - 50mcg	Tablet	25-100 mcg	68.94	Vomiting & Diarrhoea.
		ELTROXIN- 100mcg			121.60	
		THYRONORM - 100mcg			128.65	
		THYRONORM - 150mcg			187.00	
		THYRONORM - 25mcg			164.00	
		THYRONORM - 50mcg			116.55	
		THYRONORM - 75mcg			126.00	
		THYRONORM-62.5mcg			183.57	
48	TRIAMCINOLONE ACETONIDE	KENACORT - 40mg	Injection ,Paste	Apply 2-3 times daily. Rub in gently.	99.50	Peptic ulceration, acne eruptions, glaucoma, osteoporosis.
		KENACORT - 5gm			48.00	
		KENACORT - 10mg			49.00	

### **HYPOLIPIDEMIC DRUGS**

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR		
1	ATORVASTATIN	ATOCOR - 10mg	Tablet	Adult: 10-80 mg /day children > 6 years same as adults.	70.75	GI disturbance , flu syndrome, back pain, allergic reactions, asthenia.		
		AVAS - 10mg			65.90			
		AZTOR - 10mg			68.43			
		AVAS - 20mg			187.00			
		TG-TOR - 40mg			211.35			
		STORVAS - 10mg			105.93			
		AVAS-80mg			347.00			
		ATORVAT Z-20mg			149.50			
		ATOCOR-5mg			62.25		Once a day	Constipation,Flatulence,Dyspepsia Abdominal pain.
		ATORVAT Z-10mg			61.50		Once daily.	Diarrhea, heartburn, gas, joint pain, forgetfulness or memory loss, confusion.
2	ATORVASTATIN & FENOFIBRATE	ATOCOR - F	Tablet	Once a day	472.50	Muscle pain, Muscle weakness Joint pain, Back pain, Stuffy/runny nose or nose bleeds, Headache.		
3	ATORVASTATIN + ASPIRIN	ECOSPRIN AV-75/20mg	Capsule	One or two capsules once daily.	32.52	Coughing, diarrhea, gas, heartburn, joint pain, forgetfulness,confusion.		
4	ASPIRIN, CLOPIDOGREL AND ROSUVASTATIN	LIPIROSE GOLD-20MG CAPSULE	Capsule	Once a day	198.00	Increased bleeding tendency, nausea, indigestion, diarrhea, stomach pain, constipation, muscle pain, dizziness and headache.		

5	ATORVASTATIN + FENOFIBRATE	TG-GOAL -145mg	Tablet		130.50	Kinase concentrations, flu syndrome Mild to moderate decreases in Hb, haematocrit and leukocyte counts. Potentially Fatal: Hepatitis, cholecystitis, pancreatitis, rhabdomyolysis.
6	ATORVASTATIN (10MG) + CLOPIDOGREL	AVAS C.V-10/75mg	Capsule	10 or 20 mg once daily, may be adjusted at 4-wk interval.	94.01	Stomach pain, Diarrhoea, Indigestion, Constipation, Headache, Dizziness, Muscle pain, Weakness, Increased glucose level in blood, Increased liver enzymes, Increased bleeding tendency.
		AVAS C.V-20/75mg TABLET	Tablet		115.00	
7	ATORVASTATIN-20MG+CLOPIDOGREL-75MG+ASPIRIN	ECOSPRIN GOLD-10mg	Tablet	1tab/day may be increased according to requirements.	81.94	Diarrhoea, Stomach pain, Constipation, Indigestion, Weakness, Increased glucose level in blood, Headache, Dizziness, Muscle pain, Increased liver enzymes, Increased bleeding tendency.
		ECOSPRIN GOLD-20mg			110.50	
		ECOSPRIN-325mg			11.13	
8	ASPIRIN / ACETYLSALICYLIC ACID + ATORVASTATIN + CLOPIDOGREL	CLAVIX GOLD-10mg	Capsule	Loading dose: 4 tab; maintenance: 1 tab/day	57.83	Diarrhoea, Stomach pain, Constipation, Indigestion, Weakness, Increased glucose level in blood, Headache, Dizziness, Muscle pain, Increased liver enzymes, Increased bleeding tendency.
		CLAVIX GOLD-20mg			78.05	
9	FENOFIBRATE	FENOLIP - 145mg	Tablet	Adult: PO Standard micronised formulation: Initial: 67 mg tid or 200 mg once daily, may adjust to 67 mg 2-4 times/day or 267 mg once daily. Non-micronised formulation: Initial: 200-300 mg/day in divided doses. Usual range: 200-400 mg/day. Formulation w/ improved bioavailability: 40-160 mg once daily may be used.	121.00	Abnormal LFTs , increased creatini.
10	ROSUVASTATIN	ROZAVEL - 10mg		Adult: 5-20mg OD	128.00	Headache dizziness, Gi upset, peripheral edema, insomnia, proteinuria.
		ROZAVEL - 20mg			223.00	
		ROZAVEL - 40mg			345.00	
		ROZAVEL - 5mg			68.00	
		ROSEDAY-10mg TABLET	Tablet	Adult: 5-20mg OD.	94.92	Headache dizziness, Gi upset.
		ROSUVAS-20mg(COMBI)	Tablet	Once daily.	1980.00	Headache, Dizziness, Stomach ache, Constipation, Nausea.
		ATORFIT CV-10/75MG	Capsule	Once daily.	183.30	stomach bleeding, gastrointestinal bleeding, nose bleeding), rash, diarrhoea, stomach ulcer, muscle disease.
11	ROSUVASTATIN + FENOFIBRATE	ROZAVEL-F -10mg	Tablet	Once daily.	168.00	Headache,Muscle pain Constipation, Abnormal liver function tests,Joint pain.
		ROZAVEL-F--5mg			10.50	

12	ROSUVASTATIN (10MG) + CLOPIDOGREL (75MG)	ROZALET-10MG	Tablet	Once a day	167.00	Musculoskeletal Pain (Bone, Muscle or Joint), Headache, Nausea or vomiting, Dyspepsia Diarrhoea, Muscle Ache (Read More - Home Remedies for Muscle Aches) Joint pain.
13	METOPROLOL SUCCINATE	SELOKEN XL-50MG SELOKEN XL-25MG	Tablet	Once daily.	61.70 41.60	Drowsiness, dizziness, tiredness, diarrhea, and slow heartbeat.
14	TELMISARTAN (40MG) + AMLODIPINE (5MG)	TAZLOC-AM	Tablet	Once a day	141.55	Sleepiness, Ankle swelling Headache, Flushing (sense of warmth in the face, ears, neck and trunk) Dizziness, Tiredness.

### IMMUNOMODULATORS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR	
1	RITUXIMAB	CYTOMAB* 500mg/50ml	Injection	One Vial of 50ml of concentrate for solution for infusion For intravenous administration after dilution.	28000.00	Low blood pressure Nausea and Vomiting Chills and fever Abdominal pain Decrease in white blood cells and platelets	
		ENFIERA-500mg	Injection		28000.00		Itching
		MABALL-500mg/50ml	Injection	Once weekly for 4 doses	28000.00	Nausea, Vomiting Diarrhea, Lack of energy.	
		MABTAS-100mg	Injection	375 mg/m2 IV once weekly for 4 or 8 doses.	9432.42	Nausea, Rash, Pruritis, Headache Fever.	
		MABTAS N-500mg	Injection	Once weekly for 4 or 8 doses.	24150.00	Diarrhea, back or joint pain, flushing, night sweats, feeling unusually anxious or worried mouth sores.	
		RITUXIREL-500mg/50ml-	Injection	Once weekly for 4 weeks.	28000.00	Fever, chills, body aches, anemia, infections, nausea, diarrhea, swelling in your hands or feet, feeling tired.	
2	CYLOPHOSPHAMIDE	ENDOXAN-50mg	Tablet	Malignancies Low dose regimen: 2-6 mg/kg/wk in divided doses. Moderate dose regimen: 10-15 mg/kg /wk; high dose regimen: 20-40mg/kg	4.00	Congestive heart failure; leucopenia poor wound healing; anorexia. Nausea, vomiting; alopecia; oral mucosal ulceration; anemia thrombocytopenia, nonhaemorrhagic cystitis.	
		ENDOXAN-N 1gm	Injection		280.70		Nausea.
		ENDOXAN-N-500mg			83.57		vomiting.
		CYPHOS-1gm			108.00		loss of appetite or weight. abdominal pain.
		CYPHOS-500mg			82.00		diarrhea. hair loss.

		PHOSMID - 1gm	Injection	Malignancies Low dose regimen: 2-6 mg/kg/wk in divided doses. IV Malignancies Moderate dose regimen: 10-15 mg/kg /wk; high dose regimen: 20-40 mg/kg.	79.00	Congestive heart failure; leucopenia poor wound healing; anorexia. Nausea, vomiting; alopecia; oral mucosal ulceration; anemia thrombocytopenia, nonhaemorrhagic cystitis.
3	IGURATIMOD	IGURATI-25mg	Tablet	Twice a day	260.00	Elevated liver enzymes Low blood platelet count.
4	LEFLUNOMIDE	LEFNO-20mg	Tablet	Once a day	242.00	Nausea or vomiting, Headache Dyspepsia, Rash, Joint pain, Dizziness.
5	METHOTREXATE	FOLITRAX - 5mg	Injection,Tablet	Tab: 2.5 - 30 mg/day IV 10-60 mg/m2	38.27	Ulceration of the mouth and GI disturbances (e.g. stomatitis and diarrhoea), bone marrow depression, hepatotoxicity, renal failure, skin reactions, alopecia, ocular irritation, arachnoiditis.
		FOLITRAX 15mg -1ml			38.50	
		ONCOTREX - 2.5mg			54.64	
		ONCOTREX - 5mg			84.80	
		METHOFAST-50mg/2ml	Injection	Adult: 15-30mg daily for 5 days	38.00	
6	MYCOPHENOLATE MOFETIL	BIOMYF-500mg	Tablet	1- 1.5 g twice daily	750.00	Diarrhoea,vomiting,GIhaemorrhage and perforation; leucopenia;asthenia pain, headache, anaemia, thrombo cytopenia, renal tubular necrosis.
		JBMUNE-500mg	Tablet	Twice a day	700.00	Nausea,Vomiting, Diarrhea, Abdominal pain, Headache High blood pressure, Decreased white blood cell count (neutrophils).
7	RANIBIZUMAB	RANIEYES-2.3MG/0.23ml	Injection	Once a month	18000.00	Blindness. bloody eye. blurred vision or loss of vision. decreased vision or other changes in vision. disturbed color perception. dry eye.
8	TACROLIMUS	TACROVERA-0.1%-15ml	Lotion	Apply a thin layer over the affected skin twice daily.	530.00	High blood pressure. Kidney function that gets worse. Chance of getting an infection. Wash hands often. Stay away from people with infections, colds, or flu. Feeling dizzy. Rise slowly over a few minutes when sitting or lying down. Be careful climbing. Headache. Upset stomach.
9	TOFACITINIB	TOFASHINE	Tablet	Twice a day	384.00	Upper respiratory tract infection, Headache Diarrhea, Nasopharyngitis.
		TOFASHINE XR-11mg	Tablet	Once a day	950.00	

## MUSCLE RELAXANTS, ANAESTHETIC & EMERGENCY

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ARTARICURARIUM	ARTACIL - 2.5mg	Injection	Adults: 3-6 mcg/kg; Paed: 150 mcg/kg.	128.00	Allergic reactions, hypotension, tachycardia, bronchospasm.
2	ACECLOFENAC + PARACETAMOL+ THIICOLCHICOSIDE	HIFENAC-MAX	Tablet	Twice daily	198.00	Diarrhea, Nausea Redness on skin, Vomiting, Running nose, Headache,Insomnia, Stomach cramps, Photosensitivity reactions,Constipation.
3	BACLOFEN	RICLOFEN - 10mg	Tablet	5 mg 3 times/day for 3 days, increased to 10 mg 3 times /day for 3 days.	51.00	Sedation, drowsiness, ataxia, dizziness, headache, confusion, hallucinations, skin reactions, GI symptoms, enuresis.
		LIOFEN-XL-20mg	Tablet		25.00	
		LIOFEN-XL-30mg				
		BACFEN-25mg	Tablet	5 mg three times a day for 3 days 10 mg three times a day for 3 days 15 mg three times a day for 3 days 20 mg three times a day for 3 days.	256.00	Fatigue,Weakness, Drowsiness,Dizziness.
4	BENZOCAINE	MUCOPAIN - 15gm	Gel	Apply up to 4 times/day.	47.00	Skin sensitivity; irritation of conjunctiva, mucosal irritation; reversible brown staining of the teeth; tongue discoloration and burning sensation; transient taste disturbance; parotid gland swelling. Potentially Fatal: Anaphylaxis.
5	BUPIVACAINE HYDROCHLORIDE	ANAWIN - 0.25% - 20ml	Injection	Adult: 225mg with adrenaline & 175 mg without adrenaline by infiltration.	51.00	Bradycardia, hypotension, HTN, nausea, vomiting, diplopia, neuropathy, peripheral nerve injury.
		ANAWIN - 0.5% - 20ml			77.00	
		ANAWIN HEAVY - 4ml			24.99	
6	CHLORZOXAZONE + PARACETAMOL	MYOSPAS	Tablet	250 - 750 mg QID	84.50	GI upset, hypersensitivity reactions,headache.
7	CHLORZOXAZONE + PARACETAMOL+ DICLO	MYOSPAZ FORTE			166.50	
8	ETOMIDATE (20MG/10ML)	TROYMIDATE-10ml	Injection	IV 300 mcg/kg over 30-60 sec.	467.50	Nausea, Rash, Vomiting, Abnormality of voluntary movements, Apnea (absence of breathing), Hyperventilation, Thrombophlebitis, Stridor, Myoclonus, Decreased blood pressure.
9	GLUCOSE & LEVOBUPIVACAINE	LEVO ANAWIN HEAVY-0.5%-4ml	Injection	Bupivacaine & Vasoconstrictor - 2mg / Kg(Upto 200mg).	108.00	Nausea,Vomiting, Paresthesia (tingling or pricking sensation), Slow heart rate Decreased blood pressure, High blood pressure.

10	IBUPROFEN, PARACETAMOL &CHLORZOXAZONE	FLEXON-MR	Tablet	The recommended initial dose is 500 mg 3-4 timesday. May reduce to 250 mg 3-4 timesday subsequently. Max: 750 mg 3-4 timesday.	1.84	Dry mouth, dizziness, drowsiness, lightheadedness, blurred vision, nausea, vomiting, and constipation.
11	ROPIVACAINE-7.5mg	ROPIN HEAVY-0.75% 4ml	Injection	Continuous 72 hour epidural infusion after an epidural block with 5 or 10 mg/mL. Epidural anesthesia with 7.5 mg/mL (0.75%) for cesarean delivery. Brachial plexus block with 7.5 mg/mL (0.75%) ropivacaine. 20 minute IV infusion to volunteers (40 mg)	108.00	Low blood pressure,Nausea Headache,Dizziness, Vomiting.
12	ROPIVACINE	ROPEE-7.5mg-20ml	Injection	75-200 mg (15-30 mL of 0.5%-1% solution)	173.90	Nausea, vomiting; headache, back pain; fever; itching; numbness or tingly feeling.
		ROPICAIN-0.2% 20ml	Injection	3mg/kg-	119.50	Low blood pressure, nausea, vomiting, slow heart rate, fever, chills, pain, postoperative complications, stuffy nose, reduced sensation, anemia, tingling, headache, itching and back pain.
13	SUCCINYLBCHOLINE	SUCOL - 10ml	Injection	Adult: IV 0.6 mg/kg; Paed: IV 1-2 mg; Maintenance: 0.3 -0.6 mg/kg.	67.00	Bradycardia, tachycardia, hypotension,hyperperntension, raised intraocular pressure hyperkalaemia, excessive salivation.
14	TOLPERISONE & PARACETAMOL	MYO-MR PLUS	Tablet	Three times a day	139.75	Upset stomach or throwing up. Harm to the liver may rarely happen.
15	THIOLCHICOSIDE	THIOSPASS-4mg	Capsule	Muscle spasms Adult:	145.20	Drowsiness, abdominal pain, gas, nausea, diarrhea.
		THIOSPAS - 4mg	Tablet		132.00	
16	THIOLCHICOSIDE + ACECLOFENAC	THIOFENAC-Z	Tablet	Initially, 16 mg daily.	223.20	Dizziness.Nausea.,Heartburn.,Diarrhoea. Stomach pain.,Dry mouth.,Indigestion.
		SPERONAC-MR	Tablet	Muscle spasms Adult: Initially, 16 mg daily. Intramuscular Muscle spasms. Adult: Up to 8 mg daily.		Photosensitivity reactions.
		THIOSPAS-A - 4mg		Initially, 16 mg daily.	153.00	Photosensitivity reactions.
		NEORELAX-A - 4mg		Intramuscular	128.00	
		THIOSPAS-A - 8mg		Muscle spasms	233.00	
		NEORELAX-MR		Adult: Up to 8 mg daily.	140.00	
		ZERDOL - TH			130.00	
17	TIZANIDINE	TIZAN - 2mg		Tablet	2 mg once daily	94.00

18	VECURONIUM BROMIDE	NEOVAC - 4mg/2ml	Injection	Adult: 80 - 100 mcg/kg maintenance: 20 -30 mcg/kg	135.00	Muscle weakness, paralysis, muscleatrophy (after long term use), hypersensitivity reactions e.g urticaria and erythema.
		KROVEC-4mg	Injection	Adult: 80 - 100 mcg/kg maintenance: 20 -30 mcg/kg Paed: < 4mth : 10-20mcg/kg > 4 mth: as in adults	135.00	
		SAMVEC-4mg/2ml	Injection	<b>INITIAL DOSE:</b> 0.08 to 0.1 mg/kg by IV bolus  <b>MAINTENANCE DOSE DURING PROLONGED SURGICAL PROCEDURES:</b> 0.01 to 0.015 mg/kg 25 to 40 minutes later, then as frequently as every 12 to 15 minutes <b>CONTINUOUS INFUSION:</b> Initiate with an intubating dose of 80 to 100 mcg/kg followed 20 to 40 minutes later with 1 mcg/kg/minute.	96.02	Skin rash, Increased saliva production, Injection site reactions (pain, swelling, redness), High blood pressure.
19	FENTANYL-50Mcg/ml-2ml INJECTION(Verfen)	VERFEN - 50Mcg/ml	Injection	Initial: 50-200 mcg followed by supplements of 50 mcg. Max: 200 mcg. Admin lower infusion rates of 0.05-0.08 mcg /kg/min. Patients w/ assisted ventilation: Initial: 300-3,500 mcg (up to 50 mcg/kg) followed by supplements of 100-200 mcg depending on response. Loading dose (alternatively via bolus): Approx 1 mcg/kg/min given for the 1st 10 min followed by infusion of approx 100 ng/kg/min	46.00	
20	GLYCOPYRROLATE	PYROLATE - 1ml	Injection	IM/IV 200- 400 mcg every 3-4 hrs	17.80	Xerostomia; loss of taste, nausea, vomiting, constipation, reduced sweating; urinary hesitancy and retention.
21	GLYCOPYRROLATE, NEOS TIGMINE.	MYO-PYROLATE-5ml	Injection	5-10 ml IV over a period of 10-30 secs.	70.20	dryness of the mouth, blurred vision, headaches, clouded thoughts, nervousness, fatigue, dizziness, stomach problems, vomiting and constipation.

22	BUPRENORPHINE	MUTRANS-10mcg	Patches	IM Anaesth premed 300 mcg. IV Perioperative analgesia 300-450 mcg. IM/IV Moderate to severe pain 300-600 mcg 6-8 hrly.	995.00	Nausea, vomiting, dizziness, sweating and drowsiness.
		BUPRIGESIC-5mcg		Once every 7 days	570.00	Headache. stomach pain. constipation. difficulty falling asleep or staying asleep. mouth numbness or redness.
		BUPRIGESIC-10mcg			1094.00	tongue pain
		BUPREGESIC-1ml INJECTION	Injection	IM Anaesth premed 300 mcg. IV Perioperative analgesia 300-450 mcg. IM/IV Moderate to severe pain 300-600 mcg 6-8 hrly.	21.00	Nausea, vomiting, dizziness, sweating and drowsiness.
23	KETAMINE	ANEKET-250mg-5ml	Injection	Adult: IV 1-4.5 mg/kg as inj or infuse 0.5-2 mg/kg. IM 6.5-13 mg/kg.	52.65	
		KETMIN-250MG-5ML	Injection	1-4.5 mg/kg slow IV once.	58.66	Nausea, Agitation, Double Vision, Erythema, Vomiting, Nightmare, Abnormal Dreams.
24	LIDOCAINE	LOX - 10 % - Spray	Spray, Injection,	Adult: 1-1.5 mg/kg	393.00	Arrhythmia, bradycardia, arterial spasms, CV collapse, oedema, flushing, heart block, hypotension, sinusnodesuppression, agitation.
		LOX - 4% TOPICAL - 30ml	Gel, Cream	Paed: 0.5-1mg/kg	34.00	
		ANORELIFE - 30gm			107.00	
		TOPLAP-10gm Cream			210.00	
25	LIGNOCAINE HYDROCHLORIDE	LOX - 2% JELLY 30gm	Jelly, injection	Adult: 300mg Paed: 3mg/kg	40.00	Arrhythmia, bradycardia, arterial spasms, CV collapse, oedema, flushing, heart block, hypotension, sinusnodesuppression, agitation.
		LOX-2% - 30ml			33.50	
		LOXICARD - 2% - 50ml			55.00	
		LOX-2% AND ADRENALINE-30ml			29.75	
26	LIDOCAINE TOPICAL + PRILOCAINE TOPICAL	PRILOX-5gm	cream	Max: 20 g in 24 hr	99.00	Dizziness, drowsiness, vomiting, blurred vision and slurred speech.
27	MIDAZOLAM	MEZOLAM - 1ml	Injection	0.05 - 0.15 mg/kg	27.70	Decreased respiratory rate, hypotension, bradycardia, GI upset.
		MEZOLAM - 5ml			32.50	
28	MORPHINE SULPHATE-10mg TABLET	VERMOR - 10mg	Tablet	5-20 mg 4 hrly. Extended-release: 5-20 mg 12 hrly.	5.00	
29	MORPHINE-10mg/ml-1ml INJECTION(Vermor)	VERMOR - 10mg/ml	Injection	5-10 mg at 1-2 mg/min followed by a further 5-10 mg as necessary	22.00	
30	PETHIDINE-50mg/ml-1ml INJECTION(Verpat)	VERPAT - 50mg/ml	Injection	Moderate to severe acute pain 25-100 mg IM/SC inj or 25-50 mg by slow IV inj repeated after 4 hr	49.35	
31	PENTAZOCINE	MAXUWIN-1ml INJECTION	Injection	30 mg every 3 to 4 hours.	35.00	Vomiting,Nausea.
32	PROPOFOL	NEOROF 10mg -10ml	Injection	Adult: 1.5 - 2.5 mg/kg IV upto 15 mg/kg/hr Paed: 3 - 3.5 mg/kg	98.00	Nausea, vomiting, hiccough, ophthalmoplegia, anaphylaxis.



33	SEVOFLURANE (99.97%)	SEVORANE-250ml	Liquid	0.5-3% with or without Nitrous oxide.		Vomiting, Nausea, Hallucination, Abnormal dreams, Nightmare, Confusion, Agitation, Generalized tonic-clonic seizure, Double vision, Increased blood pressure, Tachycardia.
		SEVOTROY-250ml	Liquid		7682.71	
34	HYDROXYETHYL STARCH(HES) (6GM) + SODIUM CHLORIDE (0.9GM)	VOLUVEN FREEFLEX-6%-500ml	I.v fluid	IV Blood loss & hematocrit up to approx 20 mL/kg/day equiv to 1.2 g hydroxyethyl starch/kg/day Patient weighing 70 kg 1,500 mL. Septic shock due to burns Adult Infusion rate approx 25 mL/kg/min & 12.5-25 mL/min or 1,000 mL in 40-80 min.	531.18	Nausea, vomiting, constipation, lightheadedness, dizziness, drowsiness, increased sweating, urinary retention, headache or dry mouth.
35	ATROPINE SULPHATE	ADRENALINE - 1ml	Injection	0.2-0.5 ml of 1: 1000 solution ( 0.2-0.5 mg)	15.00	Cardiac arrhythmias, rise in BP, angina, tachycardia, anxiety.
		VASOCON			20.00	
		TROPINE - 1ml	Injection	Adult: IM/IV 0.6-m 2mg; Paed: < 5kg: 0.2 mg/kg > 5kg : 0.1-0.2 mg/kg.	4.60	Rapid pulse, flushing, shortness of breath, urinary retention, GI upset.
		ATROPINE SULPHATE -100ml			98.00	
36	ANTI SNAKE VENOM	ANTI SNAKE VENOM - 10ml	Injection	cobra antivenom 0.6 mg/1 mL, bungarus caeruleus venom antiserum 0.45 mg, daboia russellii antivenin 0.6 mg, echis carinatus venom antiserum 0.45 mg.	950.00	HS reaction.
37	BOVINE LIPID EXTRACT SURFACTANT SUSPENSION	NEOSURF-5ml	Suspension	adult and children more than 12 yrs: 2cap/3times daily.	9202.00	Drowsiness, dizziness, hypotension or a headache.
38	MAGNESIUM SULPHATE	MAGNEON - 2ml	Injection	Adult: im/iv 1-2 G Paed: IM/IV 25-100mg/kg/ dose.	11.00	Flushing, sweating, hypotension, hypothermia.
		MAGNESIUM (SULP) PACKET - 20gm			10.00	
39	MAGNESIUM CITRATE, AND PYRIDOXINE.	ALKASTON B6-200ml Syrup	Oral solution	less than 6yrs 0.5 mL /Kg orally upto max/; 200mL repeat every 4-6hrs util stool are clear ; 6-12yrs - 100-150mL/DAY once;Adult -240mL once/day.	137.00	Bowel movement within 6 hours after taking the medicine; pain with bowel movements, rectal bleeding; watery diarrhea, nausea, vomiting, severe stomach pain;painful or difficult urination;
40	MAGNESIUM CITRATE 375 MG+POTASSIUM CITRATE	KMG-CIT WITH B6-200ml Syrup	Oral solution	less than 6yrs 0.5 mL /Kg orally upto max/; 200mL repeat every 4-6hrs util stool are clear ; 6-12yrs - 100-150mL/DAY once;Adult -240mL once/day.		Bloating Confusion Anxiety Uneven heart rate.
41	NEOSTIGMINE METHYLSULPHATE	MYOSTIGMIN - 1ml	Injection	Adult:IM/IV/SC: 0.04-0.05mg/kg; Paed: IM/IV/SC-0.01-0.04mg/kg.	4.45	Salivation, GI upset, drowsiness, visual changes, arrythmia.

42	POTASSIUM CHLORIDE	POTCL 10ml	Injection, syrup	Adult: IV infusion 10-20 meq/hr	23.60	Abdominal pain with cramps, diarrhea, flatulence, hyperkalemia.
		POTKLOR - 200ml		Paed: IV infusion 10-20 meq/hr	68.68	
43	POTASSIUM CITRATE-1100MG/5ML+MAGNESIUM CITRATE	KMAC-200ml Syrup	Oral solution	less than 6yrs 0.5 mL /Kg orally upto max/; 200mL repeat every 4-6hrs util stool are clear ; 6-12yrs - 100-150mL/DAY once;Adult -240mL once/day.	134.00	Bloating Confusion Anxiety Uneven heart rate.
44	PRALIDOXIME		Injection	Adult: IV/IM Used in combination w/ atropine. As Cl,iodide or mesilate: Upon atropinisation, administer 1-2 g,repeat dose after 1 hr, then 8-12 hrly if needed. Severe: 200-500 mg/hr via continuous infusion, titrate according to response.	170.00	Drowsiness, dizziness, visual disturbances, nausea, tachycardia, headache, hyperventilation and muscle weakness.
45	SODIUM BICARBONATE	AUXISODA-IV - 100ml	Injection, Tablet	Adult:1 mEq/kg then 0.5 mEq/kg at 10 min interval; Paed: 1-2 mEq/ kg/ dose.	140.00	Metabolic alkalosis, hypernatremia.
		AUXISODA-500mg			27.50	
		SODAE (8.4 %)			35.00	
		SODIUM - BI-CARBONATE - 10ml			20.00	
46	CALCIUM CHLORIDE + MAGNESIUM CHLORIDE + POTASSIUM CHLORIDE AND SODIUM CHLORIDE	PLEGIOCARD-20ml	Solution	Two to four times a day	182.38	Diarrhea,Flatulence,Stomach cramp,Increased potassium level in blood,Increased production of urine.

### NSAIDS & GOUT

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ACETAMINOPHEN	DOLOPAR - 500mg	Tablet, Syrup, Injections, Drops Suppositories, Suspension.	Adult dose : 500, 650 mg; Paedaitric : 10-15 mg/kg.	35.50	Thrombocytopenia, Pancytopenia, Neutropenia.
		DOLO 650mg			33.46	
		NEOMOL - 170mg			33.50	
		NEOMOL - 250mg			58.00	
		NEOMOL - 80mg			27.00	
		NEOMOL I.V - 100ml			297.00	
		P-250 - 60ml			35.10	
		FEVASTIN - 2ml			9.69	
		CALPOL- 15ml			24.00	
		CALPOL - 500mg			16.30	
		WIKORYL			40.50	

		SINAREST			38.20	
		SINAREST			50.33	
		DOLO-100ml INFUSION			268.00	
		CROCIN-60ml SUSPENSION			32.25	
2	ACETAMINOPHEN + DOMPERIDON	GRENIL	Tablet	2 tab at the onset of attack, may repeat 4 hrly if needed. Max: 4 g paracetamol/day.	42.50	Irregularity in heart beat A feeling of dizziness or faintness Difficulty in breathing, swelling of the throat or the face Menstrual irregularities Discharge of breast milk from the nipple Swelling of breasts in males.
3	ACETAMINOPHEN + PARACETAMOL	ACTON-OR	Tablet	352-650 mg every 4-6 hours.	46.00	Low fever with nausea, stomach pain, and loss of appetite; dark urine, clay-colored stools; or jaundice.
4	ACETAMINOPHEN + TRAMADOL	TRAFEN Z	Tablet	2 tablets orally every 4 to 6 hours.	69.50	Constipation, tiredness, increased sweating, dizziness, drowsiness, weakness, stomach pain, loss of appetite.
		ULTRACET	Tablet	2 tab 6 hrly. Max: 8 tab/day. Max duration: 5 days.	130.00	Dizziness, drowsiness, tired feeling; stomach pain, nausea, loss of appetite; diarrhea, constipation; or sweating.
5	ACECLOFENAC	DOLOKIND	Tablet	Twice daily.	26.55	Diarrhea, nausea, headache, indigestion, heartburn, abdominal pain, flatulence, feeling sick, dizziness and rash.
		ACECLO-Z-FENAC-100mg	Tablet	Twice daily.	51.00	Abdominal pain, Constipation, Diarrhea, Nausea and vomiting, Skin rash .
6	ACECLOFENAC, PARACETAMOL WITH SERRATIOPEPTIDASE	HIFENAC-D-TABLET	Tablet	The recommended dose is one tablet to be taken twice daily.	127.50	Dizziness •nausea (feeling sick) •diarrhoea •increased liver enzymes in the blood.
7	ACECLOFENAC + DROTAVERIN	DROLGAN	Tablet		87.00	
8	ACECLOFENAC+RABEPRAZOLE CISSUS QUADRANGULARIS EXTRACT MESALAMINE	CISSACT	Tablet	100mg bd 2tabs/bd per day 2.4 g/day in divided doses. Maintenance of remission: 1.2-2.4 g/day in divided doses.	154.00	Headache, intestinal gas, dry mouth, diarrhea, and insomnia.
		MESACOL-400mg	Tablet		75.93	Mild nausea, vomiting, stomach cramps, diarrhea, gas, fever, sore throat, or other flu symptoms, constipation, headache or dizziness, tired feeling or skin rash.
9	ASPIRIN	ECOSPRIN - 150mg	Tablets	Adult dose : 75- 150 mg; 300 -	7.00	Prolonged bleeding time.
		ECOSPRIN - 75mg			4.26	GI disturbances.
		ECOSPRIN AV - 75mg			34.30	
		DISPRIN			4.70	
14	DROTAVERINE	DROTIKIND - 2ml	Injection, Tablet	40-80 mg 3 times/day.	16.50	Vertigo, nausea, vomiting, dry mouth.
		DROTIN			16.21	

		DROTIN DS			70.03	
		DROTIN			40.69	
		DROTIN - M			99.73	
11	DROTAVERINE HYDROCHLORIDE & PARACETAMOL	DROTIN PLUS	Tablet	40-80 mg 3 timesday.	12.61	Difficulty in breathing, Allergic skin reaction, Swelling of face, lips, eyelids, tongue, hands and feet, Fall in blood pressure.
12	DICLOFENAC + PARACETAMOL & SERRATIOPEPTIDASE	ENZOFLAM	Tablet	Twice a day	152.00	Nausea,Vomiting,Stomach pain,Indigestion,Heartburn Loss of appetite,Diarrhea
13	DICLOFENAC SODIUM	VOVERAN EMULGEL - 30gm	Tablet, Injection, Gel,	Adult dose : 50mg.	78.90	Oedema, Syncope, Nausea, Diarrhoea.
		OXALGIN NANOGEL - 30gm	Sprays, suppositories, Liniment.		90.00	
		NOBEL GEL - 30gm			50.60	
		VOLITRA PLUS			83.00	
		VOVERAN - 50mg			74.21	
		JUSTIN SUPPOSITORIES - 12.5mg			35.00	
		JUSTIN SUPPOSITORIES - 25mg			37.00	
		JUSTIN SUPPOSITOTIES - 100mg			52.90	
		VOVERAN - 1ml			19.47	
		DICLOGESIC			32.85	
		LYSER - D			154.00	
		FENLONG MR LINIMENT - 30ml			50.00	
FENLONG LINIMENT-30ml		54.00				
14	DICLOFENAC SODIUM + METAXALONE	FLEXURA - D	Tablet	Three to four times a day	115.00	nausea, vomiting, upset stomach, drowsiness, headache and nervousness.
15	ALCOHOL 10 %V/V+DICLOFENAC 4 %W/V	VOLITRA APS SPRAY-30ml	Spray	100-200 mgday in divided doses.	235.00	Itching,Headache,Drowsiness.
16	DICYCLOMINE	COLIMEX	Injection, Tablet	10-20 mg 3 times/day. IM 20 mg 4 times/day for 1-2 days.	29.65	Difficulty in accommodation, exacerbation of glaucoma; tachycardia, palpitations, arrhythmias; urinary retention; restlessness; confusion, excitement, hallucination and delirium.
		CYCLOPAM - 10ml	Syrup, Drops		31.90	
		CYCLOPAM - 30ml			36.30	
		CYCLOPAM			36.90	
17	DISODIUM HYDROGEN CITRATE	ALKASOL-100ml	Syrup, Drops	15 to 30 ml two to three times a day.	115.14	Stomach pain, Tiredness, Diarrhea, Nausea, Vomiting, Frequent urge to urinate.
18	ETORICOXIB	ETOCOX - 90mg	Tablet	Adult: 60-90 mg	66.30	Constipation, Nausea, Vomiting.
		ETOSHINE - 60mg			76.00	
		ETROBAX-60mg			114.50	
19	ETORICOXIB + THIOLCHOSIDE	ETORIFLEX-MR-4mg	Tablet	Adult: 60- 4 mg	165.00	Constipation, Nausea, Vomiting.
		ETORIFLEX-MR-8mg			209.00	
		ETOSHINE - MR			169.00	
20	ETODOLAC	ETOVA - 400mg	Tablet	600-1,000 mg/day	58.50	Constipation, Nausea, Vomiting.
		ETOVA - P - 400mg			78.65	

21	ETODOLAC + THIOCOLCHICOSIDE	ETOVA-MR			157.00	Constipation, Nausea, Vomiting.
22	FEBUXOSTAT	FEBUSTAT - 40mg	Tablet	40 - 80 mg OD	102.00	GI Distrubances, Arthritis, Anemia, Hepatitis, Heptomegaly Eczema.
23	IBUPROFEN	BRUFEN - 200mg			4.30	
		BRUFEN - 400mg			7.80	
		IBUGESIC - 100ml			22.50	
24	IBUPROFEN + PARA	COMBIFLAM	Tablets, Syrup	Adult : 200 - 600 mg	11.80	GI Irritation, Nausea, Diarrhoea.
		COMBIFLAM			21.90	
25	INDOMETACIN	MICROCID - 25mg	Capsule	Adult : 25mg bid ro tid - 200mg/day	33.00	CHF, Tachycardia, Chest pain, Arrhythmia, Palpitations, HTN, Pulmonary HTN, Odema, GI disrubances.
		MICROCID-SR - 75mg			85.00	
26	KETOROLAC	KETOROL - 1ml	Tablets , Injection	Tablet 20- 40 mg/day  10-30 mg 4-6 hrly	20.00	Drowsiness, Dizziness, Headache, Mental and Sensory changes, Psychotic reaction, Sweeting, Dry mouth, Thirst, Fever. Convulsions, Myalgia, Aseptic meningitis, HTN, Dyspnoea, Pulmonary odema, Bradycardia, Chest pain, Palpitations, Fluid retention.
		KETOROL-10mg			40.30	
27	LEFLUNOMIDE	SUPRIDOL - 1ml	Tablet	100 mg	123.38	HTN, Diarrhoea, Upper respiratory tract infection, respiratory tract infection, Thrombocytopenia, Leukopenia, Alopecia, Dry skin, Eczema, Erythema multiforme
		LEFRA - 10mg			101.81	
28	MEFENAMIC ACID + DICYCLOMINE	MEFTAL SPAS	Tablet		31.00	Dizziness, Dryness in mouth, Blurred vision Nausea, Sleepiness
29	MESALAZINE + MESALAMINE	MESACOL-OD	Tablet	Once daily		Flatulence, Hair loss, Headache, Itching, Diarrhea, Nausea, Stomach pain / epigastric pain.
30	NAPROXEN	NAPROSYN-250mg	Tablet	0.5-1 g/day in 1-2 divided doses.	20.50	GI disturbances, arthritis, Anemia, Hepatitis, Heptomegaly eczema.
31	NEPAFENAC	NEPAFLAM-5ml	Drops	1 drop, 0.3% should be applied to the affected eye one-time-daily beginning 1 day prior to cataract surgery, continued on the day of surgery and through the first 2 weeks of the postoperative period. An additional drop should be administered 30 to 120 minutes prior to surgery.	137.00	Application site irritation.
32	PIROXICAM	DOLONEX - 20mg -1ml	Tablets, Injection	Adult: 10 -30 mg	18.00	Upset stomach, Constipation.
		DOLONEX - DT - 20mg			79.66	
33	PARACETAMOL	PARA -Z-650mg	Tablet	Adults is 1 gram (1000 mg) per dose and 4 grams (4000 mg) per day.	37.40	Adults is 1 gram (1000 mg) per dose and 4 grams (4000 mg) per day.

		P-250mg	Tablet	Every 4 to 6 hours.	13.20	Nausea and Vomiting, Stomach pain, Diarrhoea, Indigestion, Skin Rash.
		KABIMOL-50ml	Infusion	10 to 15 mg/kg every 4 to 6 hours.	238.00	Nausea, Stomach pain, Dark coloured urine.
		PARACETAMOL-250m/60ml	Syrup	Every 6 hrs	44.35	An allergic reaction, which can cause a rash and swelling.
34	PARACETAMOL CAFFEINE PHENAZONE	DART	Tablet	Every 4 to 6 hours.	32.90	Nausea, stomach pain, loss of appetite, upper stomach pain, dark urine, clay colored stool, allergic reactions, skin rash and kidney impairment.
35	PARACETAMOL + ACECLOFENAC	HIFENAC - 100 MG		Adults: 100mg	31.50	Nausea, Vomiting, Perforation, GI ulceration, Bleeding, Bronchospasm, Coagulopathy.
		HIFENAC -D			68.50	
		DOLOKIN PLUS		1 tablet should be taken 1 - 2 times daily	23.47	
		PARAFENAC-Z	Tablet	352-650 mg every 4-6 hours.	4.50	
		ACTON-OR	Tablet		46.00	
36	PIROXICAM (20MG)	UGESIC-20mg	Tablet	20 mg/day as a single or in divided doses as necessary.	106.50	Nausea, vomiting, constipation, lightheadedness, dizziness, or drowsiness.
37	RIZATRIPTAN BENZOATE ORALLY DISINTEGRATING	RIZANET-10mg	Tablet	Rizatriptan dosage is 5 mg single dose in patients weighing less than 40 Kg and 10 mg single dose in patients weighing 40 kg or more.	53.00	Chest pain Drowsiness Dizziness Shortness of breath Feeling of warmth or heat Irregular heartbeat Itching Numbness Dry mouth Nausea and Vomiting Unusual tiredness and weakness.
		RIZANET-5mg			36.00	
38	SERRATIOPEPTIDASE	LYSER FORTE	Tablet	Adult: PO 5-10 mg 3 times/day.	77.00	Skin rash, diarrhoea, anorexia,
39	TRAMADOL	SUPRIDOL - 1 ML	Injection, Capsule	PO 50-100 mg 4-6 hrly,	13.20	GI disturbances, arthritis, Anemia, Hepatitis, Heptomegaly eczema
		DOMADOL	Tablet		69.89	
40	TRAMADOL + DICLOFENAC	DURAPAIN	Tablet	IV/IM 100 mg, then 50 mg every 1 10-20 mins if needed, up to 250 mg for the 1st hour.	119.00	
41	TAPENTADOL	TYDOL - 50mg	Tablet	Adult: PO 5-10 mg 3 times/day	118.00	Bladder pain bloody or cloudy urine, body aches , chills.
42	TRYPSIN	CHYMORAL FORTE	Tablet	2-3tab/ 3times daily	265.00	Tightness in the chest, Swelling of the mouth, Difficulty breathing, Temporary stinging sensation, Burning sensation.
		RUTOHEAL			154.00	
43	TRYPSIN, CHYMOTRYPSIN and SERRATIOPEPTIDASE	CHYMOCIP-TABLET	Tablet	2-3tab/ 3times daily	274.00	Tightness in the chest, Swelling of the mouth, Difficulty breathing, Temporary stinging sensation, Burning sensation.
44	TRYPSIN+BROMELAIN+RUTOSIDETRIHYDRATE+ DICLOFENAC	RUTOHEAL-D TABLET	Tablet	2-3tab/ 3times daily.	217.80	Nausea, Vomiting, Heartburn, Indigestion, Diarrhea.

## SEDATIVES AND HYPNOTICS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ALPRAZOLAM	ALPRAX - 0.25mg	Tablet	Adult: 0.25 - 0.5 mg BID	16.50	Anorexia, loss of memory & coordination, tachycardia, hypotension
		ALPRAX - 0.5mg			35.00	
		ALPRAX-1mg	Tablet		94.75	
2	CLONAZEPAM	ZAPIZ-0.25mg	Tablet	Adult: 1.5 mg/day Paed: 0.01 - 0.03 mg/kg/day		Anorexia, loss of memory & coordination, tachycardia, hypotension
		CLOZE ZZ-0.5mg	Tablet		32.70	
3	CHLORDIAZEPOXIDE	LIBRIUM - 10mg	Tablet	Adult:20 - 40 mg OD Paed: 5-10 mg OD	45.00	Anorexia, loss of memory & coordination, tachycardia, hypotension.
		LIBRAX			54.15	
		LIBRIUM-25mg			147.70	
4	CLOBAZAM	FRISIUM 10mg	Tablet	Adult: 20-30 mg/day, up to 60 mg/day Paed: 3-12 yr: 125 mcg/kg bid increased every 5 days. Usual maintenance dose: 250 mcg/kg bid. Max: 500 mcg/kg bid.	139.67	Anorexia, loss of memory & coordination, tachycardia, hypotension.
		FRISIUM 5mg			70.16	
5	DIAZEPAM	VALIUM - 5mg	Tablet	Adult: 2-20 mg BID to QID	15.23	Anorexia, loss of memory & coordination, tachycardia, hypotension.
6	DEXMEDITOMIDINE	DEXTOMID 100mcg - 1ml	Injection	Adult: a loading infusion of one mcg/kg over 10 minutes.a maintanance infusion of 0.2 to 0.7 mcg/kg/hour	326.00	bradycardia, hypotension, nausea, agitation, tachycardia, hypertension,xerostomia.
		DEXTOMID 50mcg - 0.5ml			195.00	
7	LORAZEPAM	NEULORA-2mg TABLET	Tablet, injection	Adult: 2-3 mg BD to TID 4 mg IV over 2-5 min Paed: 0.1 mg/ kg IV over 2-5 min	31.40	Anorexia, loss of memory & coordination, tachycardia, hypotension
		LOPEZ - 2ml			17.12	
		NEULORA-2mg			15.50	
		ATIVAN - 1mg			82.50	
8	MELATONIN+ZOLPIDEM	ZOLCALM-10mg	Tablet	3-5 mg given at bedtime 12.5 mg once daily immediately before bedtime.		Anorexia, loss of memory & coordination, tachycardia, hypotension.
		ZOLCALM-5mg			59.00	
9	NITRAZEPAM	NITROSUN-5mg	Tablet	Once daily at bedtime.	37.00	Drowsiness, dizziness, lightheadedness, fatigue, or loss of coordination may occur. Headache, stomach upset, mental confusion, diarrhea, nausea, or vomiting rarely occur.
10	PHENOBARBITONE	GARDENAL - 30mg	Tablet, Syrup	Adult: 30-60 mg OD- TDS	46.62	Sedation, HS reaction, dependence
		GARDENAL - 100ml			46.20	
		PHENOBARBITONE-1ml	Injection		100-200 mg 60-90 min	19.45
11	ZOLPIDEM	NITREST - 5mg	Tablet	Adult: 5-20 mgOD	59.00	Anorexia, loss of memory & coordination, tachycardia, hypotension

		SOBRIUM-5mg	Tablet	Women: 5 mg orally once a day immediately prior to bedtime -Men: 5 to 10 mg orally once a day immediately prior to bedtime Maintenance dose: 5 to 10 mg orally once a day immediately prior to bedtime Maximum dose: 10 mg/day	5.42	Unsteadiness, Confusion, Dizziness, Fast heartbeat, Drowsiness, Blurred vision, Dry mouth Headache, Loss of appetite Nausea or Vomiting, loss of memory & coordination, tachycardia, hypotension.
12	DOXYLAMINE SUCCINATE, PYRIDOXINE HYDROCHLORIDE	DOXINATE	Tablet	25 mg every 4-6 hr. Max:150 mg daily.	4.32	Acute dystonic reactions and long-lasting impaired consciousness in child. CNS depression including slight drowsiness to deep sleep, lassitude, dizziness, incoordination. Headache.
13	TRICLOFOS (500MG/5ML)	PEDICLORYL-30ml	Syrup	Adult: 1 - 2 g at night Children: up to 1 year: 25 - 30 mg / kg 1 - 5 year: 0.25 - 0.5 g 6 - 12 year: 0.5 - 1 g	75.00	dizziness, irritability, and vomiting.
14	BUPRENORPHINE	MUTRANS-5mcg	Patches	Once daily.	545.00	Constipation,Headache,increased sweating, tiredness or drowsiness (especially after a dose) loss of appetite, nausea and vomiting,abdominal pain, skin rashes, itching or hives.

## VACCINE

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	BACTERIUM SALMONELLA TYPHI	BIOTYPH-2.5ml	Injection	0.5 ML	650.00	Headache, fever, rash, itching, abdominal pain, nausea, diarrhea, vomiting, anaphylactic reaction,
2	DIPHTHERIA, TETANUS, PERTUSSIS (ACELLULAR, COMPONENT), HEPATITIS B (RDNA), POLIOMYELITIS (INACTIVATED) AND HAEMOPHILUS INFLUENZAE TYPE B CONJUGATE VACCINE (ADSORBED)	HEXAXIM	Injection	0.5 ML	3900.00	Injection-site pain, injection-site erythema, injection-site swelling Irritability Pyrexia (body temperature $\geq 38.0^{\circ}\text{C}$ ).
3	INACTIVATED INFLUENZA VACCINE	INFLUGEN	Injection	0.5ML	720.00	Aches pain, head ache, fever.
		INFLUVAC TETRA-0.5ml	Injection	0.5 ml.	1840.00	Body aches, Low-grade fever, Headache, Nausea, Fatigue, Fainting in rare cases.



4	PNEUMOCOCCAL POLYSACCHARIDE VACCINE	PNEUMOVAX 23 - 0.5ml	Injection	0.5 ML	1365.00	Aches pain, Increased temperature Tenderness.
5	ROTAVIRUS	ROTATEQ-2ml	Drops	3 doses at ages 2 months, 4 months, and 6 months.	1175.00	Mild fever, vomiting, runny nose, sore throat, wheezing or coughing.
6	TETANUS TOXOID	T.T - 0.5ml	Injection	0.5 ML	10.00	Aches pain, Increased temperature Tenderness.
7	TETANUS IMMUNOGLOBULIN	TETGLOB-250mg	Injection	Prophylaxis: 250 Units IM (single dose) Active tetanus: 3000-6000 Units	1200.00	Aches pain, Increased temperature Tenderness.
		TETGLOB-500mg			1474.90	
8	TETANUS & DIPHTHERIA	TD VAC-(T.T+DIP)-5ml	Injection	0.5 ML.	170.00	Aches pain, Increased temperature Tenderness.
9	VARICELLA-ZOSTER	NEXIPOX	Injection	Each 0.5ml vial/dose	1779.00	Redness, itching, tenderness, swelling, bruising, or a lump where the shot was given; low fever; or mild skin rash that looks like chickenpox (can occur up to 1 month after vaccination).
10	VARICELLA VACCINE (LIVE) ATTENUATED (1350PFU)	VARIPED	Injection	0.5ml At Once.	2220.00	Injection site reactions (pain, swelling, redness), Fever. Rash.
11	PURIFIED VI POLYSACCHARIDE TYPHOID VACCINE (25MCG)	TYPBAR-2.5ML VIAL	Injection	Single dose	577.50	Feeling of discomfort Nausea Diarrhea Fever Headache
12	YELLOW FEVER VIRUS (LIVE, ATTENUATED) (1000IU)	STAMARIL-0.5ml (YELLOW FEVER)	Injection	A single, 0.5 millilitre dose to adults and children from 6 months of age. The first dose should be given at least 10 days before protection from yellow fever is needed	2230.00	Fever, weakness, drowsiness, unusual crying, and loss of appetite.

### MISCELLANEOUS

S.No	GENERIC NAME	BRAND NAME	FORMULATION	DOSE	COST	ADR
1	ACOTIAMIDE (100MG)	ACOTRUST-100mg TABLET	Tablet	Dose:100 mg tid	190.00	Headache, Diarrhoea.
2	ACESULFAME POTASSIUM, CALCIUM SODIUM PHOSPHOSILICATE, CARBOMER, GLYCERIN, PEG 400, SILICA, SODIUM LAURYL SULPHATE AND TITANIUM DIOXIDE	VANTEJ-50gm PASTE	Paste	Calcium sodium phosphosilicate in Vantej toothpaste imparts significant level of dentinal occlusion with durable occlusive deposits following four days of twice daily brushing.	116.00	Puffiness of face, Cough syrup, Urge to vomit, Abdominal cramps, Wind, Diarrhea, Rash, Allergic reaction, Glycerin laxative-anal, Bowel irritation.

3	FOCRORHIZA + CUMINUM CYMINUM	APTIVATE-175ml SYRUP	Syrup	Children under age 6 years are given a dosage of 12mg daily on spilted basis and children above 6 years of age are given a dosage of 16mg each day.	84.50	Nausea, Stomach Upset, Skin Rash, Acute Toxicity.
4	ANTI Rh - D IMMUNO GLOBULIN(MONOCLONAL)	RHOCLONE - 300MCG (ANTI-D)	Injection	300 mcg	2466.90	Pain, Discomfort.
5	AZATHIOPRINE TABLETS	AZORAN - 50MG	Tablets	1-3 mg/kg/day.	114.50	Fever, chills; bone marrow characterised by leucopenia, depression thrombocytopenia or anaemia; anorexia, nausea, diarrhoea; arthralgias; secondary infections.
6	A 37% AQUEOUS (WATER) SOLUTION OF FORMALDEHYDE, PUNGENT GAS	FORMALIN-10LTR-020	Solution	Disinfectant	700.00	Coughing,wheezing,nausea,skin irritation.
7	BARIUM SULPHATE ORAL SUSPENSION 1 LIT	BARIUM SULPHATE SUSPENSION 1 LIT ESKAY	Suspension	5-10 g dissolved in 250 mL of water.	260.00	Bloating, Constipation
8	BISACODYL	DULCOFLEX	Tablet, Suppository	5-10 mg/day at night po As supp/enema: 10 mg/day in the morning.	11.21	Abdominal discomfort (colic, cramps).Suppositories may cause irritation and proctitis.
		DULCOFLEX - ADULT			19.05	
9	CALCITONIN	OSTOSPRAY	Spray	200 -400 IU daily	1020.00	Nasal Irritation.
10	COLLOIDAL SILVER 32 PPM	SILVEREX HEAL-15gm OINMENT	Ointment	Apply the gel on the wound/ affected area at least once daily.	93.00	Nausea, vomiting, diarrhoea; hypersensitivity, skin reactions; haematuria; crystalluria; thrombocytopenia, leucopenia, eosinophilia.
		MEGAHEAL - 15gm	Cream	Apply sufficient quantity.	105.00	Itching of skin, Skin rash, Redness of skin.
11	CISSUS QUADRANGULARIS, MORINGA OLEIFERA AND VITAMIN D3	CISBIND-TABLET	Tablet	once a day in the morning with or without food.	15.00	Dry mouth, muscle pain, increased thirst, loss of appetite, vomiting, metallic taste, upset stomach, constipation, difficult urination.
12	COLLAGEN PARTICLES	CO-MUPIMET-5ml COLLAGENS	Granules	Make a paste/solution of BIO-SEAL in sterile saline/DM water and use with the help of a syringe. Cover the wound with any conventional absorbent dressing of choice.	660.00	Itching, redness.
13	COLLOIDAL SILVER	HYDROHEAL-AM-15gm GEL	Gel	Gel to the effected area of skin 2 to 3 times daily.	96.00	Drowsiness, dizziness, hypotension or a headache.
14	COENZYME Q10	KQ-300mg TABLET	Tablet	Once a day	778.00	Very upset stomach or throwing up. Very loose stools (diarrhea).
15	COENZYME Q10, DEHYDROEPIANDROSTER ONE (MICRONIZED) & MELATONIN	OVARES PLUS-TABLET		500 mcg of selenium daily.	47.00	Drowsiness, change in sleep pattern and headache

16	DI SODIUM HYDROGEN CITRATE	CITRALKA-100ml	Syrup	PO 15-30 mL bid-tid.	65.14	Mild diuresis, stomach cramps, flatulence, diarrhoea, GI ulceration, hypokalaemia,
17	DEXTROSE AND ELECTROLYTES LIKE SODIUM, POTASSIUM, CHLORIDE, AND CITRATE IONS	ORS LIQUID-200ml	Liquid	Taken orally with a glass of water.	29.20	Dizziness, unusual weakness, swelling of ankles/feet, mental/mood changes.
18	DIATRIZOATE MEGLUMINE & DIATRIZOATE SODIUM	UROGRAFFIN - 76 %	Injection	<p>Radiographic Examination of Segments of the Gastrointestinal Tract</p> <p>Oral Administration: Adult oral dosage may range from 30 to 90 mL (11 to 33 g iodine), depending on the nature of the examination and the size of the patient. For infants and children less than 5 years of age, 30 mL (11 g iodine) are usually adequate; for children 5 to 10 years of age, the suggested dose is 60 mL (22 g iodine). These pediatric doses may be diluted 1:1, if desired, with water, carbonated beverage, milk, or mineral oil. When used in infants, the solution may be given in a nursing bottle.</p> <p>Pediatric doses may also be used in dehydrated and/or debilitated adult patients. A 1:1 dilution is also recommended when the contrast medium is used in elderly cachectic individuals.</p> <p>For very young (under 10 kg) and debilitated children the dose should be diluted: 1 part Urografin 76% (Diatrizoate Meglumine and Diatrizoate Sodium Solution) in 3 parts water is recommended.</p> <p>For Enemas or Enterostomy</p>	206.00	Stomach cramps, Rash, Heart burn.

19	DIATRIZOATE SODIUM	GASTROPAQUE - 30ML	Oral Contarast.	<p>Oral Administration: Adult oral dosage range 30 to 90 mL (11 to 33 g iodine), For infants &amp; children &lt; 5 years of age, 30 mL (11 g iodine); for children 5 to 10 years of age, the suggested dose is 60 mL (22 g iodine). These pediatric doses may be diluted 1:1, if desired, with water, carbonated beverage, milk, or mineral oil. When used in infants, the solution may be given in a nursing bottle. Pediatric doses may also be used in dehydrated and/or debilitated adult patients. A 1:1 dilution is also recommended when the contrast medium is used in elderly cachectic individuals. For very young (under 10 kg) and debilitated children the dose should be diluted: 1 part Gastropaque (Diatrizoate Meglumine and Diatrizoate Sodium Solution) in 3 parts water is recommended.</p> <p>For Enemas or Enterostomy Instillations: Gastropaque should be diluted when it is used for enemas and enterostomy instillations. When used as an enema, the suggested dilution for adults is 240 mL (88 g iodine) in 1000 mL of water.</p>	310.00	
20	DIASTASE+PEPSIN	ARISTOZYME-SYRUP	Syrup	take 5 ml of Diastase + Pepsin, immediately after meals (for adults).	81.00	Irritation in Stomach, Upset stomach, Irritation in throat, Damage in throat, Blisters, Allergic reactions.
21	DAPSONE	DAPSONE	Tablet	<p>Multibacillary leprosy 100 mg/day for 12 mth.</p> <p>Paucibacillary leprosy 100 mg/day for 6 mth.</p>	241.50	Anaemia, peripheral neuropathy, haemolysis and methaemo globinaemia (dose-related), nephrotic syndrome, psychological changes, hepatitis.
22		HAND SANITIZER-500 ML (PUMP)	Solution		250.00	
23	ESSENTIAL AMINO ACIDS AND VITAMINS	ASTYMIN - 200 ML	Syrup	Single Dose : 15 ml	152.50	Nausia, Vomiting, Haed ache.

24	FIBER	FYBOGEL-100gm POWDER	Powder	Adults and children over 12yrs: 1 measureful (5.4gm) in morning and at bedtime or 2 measureful at bedtime. Children between 6-12yrs: Half to one measureful in the morning and at bedtime or 1-2 measureful at bedtime. Children below 6yrs: Not recommended except under medical advice.	130.00	wind and bloating.
25	FINASTERIDE 1 MG	FINAX-TABLET	Tablet	Benign prostatic hyperplasia :▶ Adult: 5 mg daily, review treatment at 3–6 months and then every 6–12 months (may require several months treatment before benefit is obtained) Androgenetic alopecia in men :▶ Adult: 1 mg daily.	195.00	Impotence, loss of interest in sex, or trouble having an orgasm, abnormal ejaculation, swelling in your hands or feet, swelling or tenderness in your breasts, dizziness, weakness, feeling like you might pass out, headache, runny nose.
26	FLAVOXATE HCL	URISPAS - 200 MG	Tablet	200 mg tid.	234.91	Increased intraocular pressure; difficulty in ocular accommodation; blurred vision; dry mouth & throat GI disturbances; nervousness;vertigo headache.
27	GLUCOSAMINE & MSM & D	JOINTACE-DN	Tablet	500 mg 3-4 times daily	233.00	Heart burn, epigastric pain/tenderness, diarrhea, nausea, dyspepsia, constipation, abdominal pain, palpitations, drowsiness, skin reaction, headache, indigestion.
		REJOINT			173.00	
28	GLUCOSE	DEX-75GM	Powder	82.5 gms Glucodex – 75 in 250 – 300 ml water.	69.00	Confusion, a light-headed feeling, like you might pass out, fever, swelling in your hands or feet, sweating, pale skin, severe shortness of breath, chest pain.
29	HONEY AND CLOVE	KOFLET-H-LOZENGES	Lozenges	1 loz-three to four times a day.	24.00	clinically safe.
30	HUMAN PLASMA PROTEIN	HUMAN ALBUMIN 5 % -250 ML	Injection	Minimum effective dose in adults is 250-500 mL.	6350.00	

31	HUMAN NORMAL IMMUNOGLOBULIN	HISTOGLOB	Injection	<p>Adult: IV Prevention of infections after bone marrow transplantation 500 mg/kg/wk. Increase platelet count in patients w/ idiopathic thrombocytopenic purpura 400 mg/kg/day for 2-5 consecutive days. Kawasaki disease 1.6-2 g/kg in divided doses over 2-5 days. Guillain-Barre syndrome 400 mg/kg/day for 5 consecutive days, repeat 4 wkly if needed. Allogenic bone marrow transplantation 500 mg/kg/wk, starting 7 days before transplantation and for up to 3 mth after transplantation. Primary antibodies deficiency Initial: 400-800 mg/kg, then 200 mg/kg 3 wkly.</p> <p>Maintenance: 200-800 mg/kg/mth. For secondary immunodeficiency syndromes: 200-400 mg/kg 3-4 wkly. IM Hepatitis A 500 mg. Measles attack in immunocompromised patients Prevention: 750 mg w/in 6 days after exposure. To modify an attack: 250 mg. Primary rubella in pregnant women 750 mg.</p>	91.00	
32	HYALURONIC ACID	HYALGAN PFS 20MG - 2ML	Injection	20 mg	1450.00	
		HYNIDASE			114.24	
		ENTODASE-INJECTION	Injection	Subcutaneous Fluid Administration (Hypodermoclysis) :For infants and children less than 3 years old, the volume of a single clysis should be limited to 200 mL; and in premature infants or during the neonatal period, the daily dosage should not exceed 25 mL/kg of body weight;	131.00	Acute urticaria, Hormone imbalance, Allergic reaction, Slow growth in children and teenagers, Angioedema (swelling of deeper layers of skin), Dizziness, Injection site reaction, Edema (swelling).

				the rate of administration should not be greater than 2 mL per minute. For older patients, the rate and volume of administration should not exceed those employed for intravenous infusion. Subcutaneous Urography : 75 Units of Entodase (hyaluronidase injection) is injected subcutaneously over each scapula, followed by injection of the contrast medium at the same sites. Absorption and dispersion of other injected drugs may be enhanced by adding 50 – 300 Units, most typically 150 Units of Entodase Entodase to the injection solution.		
33	HYDROCORTISONE ACETATE CREAM 1%W/W	CUTISOFT-10GM CORT - S - 100 MG	Cream, Injection	0.1-2.5% oint/cream/lotion: Apply as directed.Intra-articular	71.00 41.00	Sodium and fluid retention. Potassium and calcium depletion. Muscle wasting,
34	HYDROXYPROPYL METHYL	HICOOL - 5 ML EYE DROPS APPAVISC SOLUTION - 5 ML	Eye drops	Drops may be used as often as needed.over the eye and squeeze out 1 or 2 drops as needed. Look down and gently close your eye for 1 or 2 minutes,Ointments are usually used 1 to 2 times daily as needed. If using an ointment once a day, it may be best to use it at bedtime.	69.50 60.00	
35	HYOSCINE BUTYLBROMID	HYOCIMAX HYOCIMAX	Injection, Tablet	PO Genito-urinary spasm; GI tract spasm As butylbromide: 20 mg 4 times/day. Motion sickness As hydrobromide: 300 mcg 30 mins before journey, then 300 mcg 6 hrly if needed.	11.39 32.87	Flushing, postural hypotension, tachycardia, fibrillation. Rarely psychotic reactions. Dizziness, drowsiness, fatigue, headache, memory loss.
36	HERBAL PREPARATION	HADJOD-TABLET	Tablet	2 tablets twice daily	147.00	Headache, intestinal gas, dry mouth, diarrhea, and insomnia.
37	HAND SANITIZER-500 ML (PUMP)	HAND SANITIZER-500 ML	Liquid		250.00	

38	IODIXANOL	VISIPAQUE-320MG -- 100ML	Injection	<p>Intra-arterial administration (arteriography): Carotid arteries: 10-14 mL, Vertebral arteries: 10-12 mL, Right coronary artery: 3-8 mL, Left coronary artery: 3-10 mL, Left ventricle: 20-45 mL, Renal arteries: 8-18 mL, Aortography: 30-70 mL, Major aorta branch: 10-70 mL, Peripheral arteries: 15-30 mL, Aortofemoral runoffs: 20-90 mL; Intra-arterial administration (IA-DSA):</p> <p style="text-align: center;">Carotid or vertebral arteries: 5-8 mL, Aortography: 10-50 mL, Major aorta branch: 2-10 mL, Aortofemoral runoffs: 6-15 mL, Peripheral arteries: 3-15 mL; IV administration CECT of head or body: 75-150 mL bolus, then 100-150 mL infusion not to exceed 150 mL, Excretory urography: 1 mL/kg, not to exceed 100 mL, Venography: 50-150 mL per lower extremity.</p>	2750.00	
39	IOHEXOL	OMNIPAQUE - 300--50ML	Injection	<p><b>Adult:</b> Inj As iohexol soln containing 6-350 mg iodine/ml: Dose and strength used dependson the procedure and route of administration.</p>	742.00	Flushing or a sensation of heat; pain, extravasation, thrombophlebitis at the inj site; nausea, vomiting, headache, and dizziness; urticaria, pruritus, pallor, sweating, metallic taste, weakness, coughing, rhinitis, sneezing, lachrymation, visual disturbances.
40	KOJIC ACID(12%W /W) , OCTINOXATE(7.5%W /W) , ARBUTIN(1.5%W /W)	KOJIVIT PLUS-15gm GEL	Gel	<p>Take small quantity of cream and rub on face in circular motion till it disappears and dry completely.Use this cream every time you wash your face to notice visible effects.</p>	325.00	Feeling tired or weak (fatigue), stomach pain, loose stools (diarrhea), headache, nausea, or blurred vision.



41	LACTITOL MONOHYDRATE	LACTIHEP PLUS-250ml SYRUP	Syrup	2 tablespoon 8-15ml during night.	211.00	Dehydration
		LACTIHEP - 200ML	Syrup	the usual recommended dosage in adults is about 15-30 ml per day. The medication should be taken with meals. If no adverse reactions show then the dosage maybe increased to 60 ml per day.	196.00	Abdominaldistension, Flatulence, Nausea, Vomiting, Dyspepsia, Cramps, Epigastric pain, Abdominal discomfort, Anal pruritus, dyspepsia.
42	LACTULOSE	DUPHALAC - 200 ML	Solution, Syrup, Enema	<b>Adult: PO Constipation</b> Initial: 10-20 g (15-30 mL)/day Max: 45 mL (or 40g of the reconstituted oral formulation)/day. Hepatic encephalopathy 60-100 g (90-150 mL)/day in 3 divided doses; adjust accordingly. Rectal Hepaticencephalopathy Mix 200 g (300 mL w/ 700 mL water or 0.9% NaCl as a retention enema. Retain enema for 30-60 mins; repeat 4-6 hrly until oral medication can be administered.	223.50	Diarrhoea, Nausia, Vomiting.
	LIVO LUK - 60 ML	68.90				
	DUPHALAC-250ml SOLUTION	262.31				
43	LIQUID PARAFFIN	LIQUID PARAFFIN - 100 ML	Suspension, Syrup	Adults is 10-30ml when required. children over the age of 3 years old is 5-20ml	60.00	No common side effects seen.
	CREMAFFIN - 200 ML	106.37				
	CREMAFFIN PLUS	123.85				
44	LOPERAMIDE HYDROCHLORIDE	LOPERMIDE	Tablet	6-8 mg/day.	16.87	Abdominal pain, distention, and discomfort; paralytic ileus; constipation, dry mouth, drowsiness, dizziness, fatigue, rash.
45	LACTATION PREPARATION	GALACT GRANULES-200gm	Granules	Ayurvedic preparation containing asparagus racemosus 15 g, pueraria tuberosa 1 g, anethum sowa 1 g, tribulus terrestris 2.5 g, glycyrrhiza glabra 1 g, cuminum cyminum 1 g, lepidium sativum 1 g. Recommended dosage of medicine is 2 teaspoonfuls or 10 grams twice a day taken with milk.	290.00	Occasional stomach upset, Diarrhea, Tiredness, Absence of a menstrual period in women, Headache, Water and sodium retention, Decreased sexual interest in men.

46	L-ARGININE , PROANTHOCYANIDIN	ARG9 - 5GM SACHET	Powder	Mix the contents of the sachet in half a glass of water and one sachet daily.	30.00	Stomach burn.
		ARGIPREG - 6.5GM SACHET			29.46	
47	L-ORNITHINE & L-ASPARTATE	HEPACOR - 10 ML	Injection & Tablet	Adult: 1 - 2tablets 3 times daily each tablet contain 150 mg of L-Ornithine L-aspartate ,Adult Dosage 300 mg 300 (300) 12 hourly IV For 3-4 Weeks.	213.00	Vomiting,Transient nausea.
		HEPAMERZ - 10ML			270.00	
		HEPAMERZ			122.00	
48	LACTULOSE SOLUTION	LOOZE-200ml LIQUID	Liquid	Adult: PO Constipation Initial: 10-20 g (15-30 mL)/day Max: 45 mL (or 40g of the reconstituted oral formulation)/day.	216.30	Diarrhoea, Nausia, Vomiting
		LACTULOS ZE-SOLUTION	Solution,Syrup,Enema	Adult: PO Constipation Initial: 10-20 g (15-30 mL)/day Max: 45 mL (or 40g of the reconstituted oral formulation)/day.	230.50	Diarrhoea, Nausia, Vomiting.
49	LACTIC ACID, SORBITOL, COCAMIDOPROPYL BETAINE, POLYQUATERNIUM-7, TEA TREE OIL, HIPPOPHAE RHAMNOIDES.	INTIWASH-50ml	Liquid	During bathing daily, it should be used every time when there is any chance of vaginal pH imbalance, like every time after urination, after unprotected sex, during periods and even during pregnancy. Any woman under the age group of 15-55 can use Intiwash.	90.00	Drowsiness, dizziness, hypotension or a headache.

50	LACTOBACILLUS ACIDOPHILUS FIDOBACTERIUM	PROVIDAC-CAPSULE	Capsule	<p>daily in 3-4 divided doses.ADULTS: BY MOUTH: For hayfever: At least 2 billion colony-forming units of lactobacillus daily for 7 weeks has been used along with 10 mg of loratadine once daily for 5 weeks.</p> <p>For preventing diarrhea caused by antibiotics: Many different lactobacillus species have been studied. In most cases, lactobacillus is given in daily doses that provide 10-100 billion colony-forming units daily. Lower doses of 100 million colony-forming units have also been used. Usually treatment is started within 2 days of beginning antibiotic treatment and continued for at least 3 days after antibiotic treatment is complete.</p> <p>For eczema (atopic dermatitis): lactobacillus has been given to pregnant women during the last month of pregnancy. Typically, lactobacillus is given alone or along with other probiotic species in doses of 100 million to 10 billion colony-forming units.</p> <p>For a condition associated with an increased risk for developing</p>	275.60	Stomach pain and discomfort, bloating, and incomplete bowel movements.
51	L-CARNOSINE	IGNICAR-200mg	Tablet	Once a day.	495.00	Stomach upset, Headache, Nausea, Vomiting.
52	LIGNOCAINE(LIDOCAINE) +METRONIDAZOLE +SUCRALFATE	CREMAGEL-H-20gm GEL	Gel	Two to three times daily.	126.50	Burning sensation Tingling Rash
53	MAGNESIUM CITRATE & POTASSIUM CITRATE	CITALIFT-200ml	Syrup	Diluted with a glass of water and taken after meals/bedtime.	136.00	Abdominal discomfort, Diarrhoea, Vomiting, Nausea.
54	MIRABEGRON	EXENA - 50mg	Tablet	Once a day.	417.30	Increased heart rate (tachycardia) Infection of the structures that carry urine (urinary tract infections), Nausea,Constipation, Headache, dizziness, Diarrhoea.

55	MIRABEGRON (25MG) + SOLIFENACIN (5MG)	MEGATAS-S 25MG	Tablet	Once daily .	375.00	High blood pressure, Nasopharyngitis (inflammation of the throat and nasal passages), Urinary tract infection, Headache, Dryness in mouth, Constipation, Increased heart rate, Blurred vision.
56	MEPHENESIN METHYL NICOTINE & CAPSICUM OLEORESIN	RELAXYL - 30GM	Ointment	Apply half inch ointment over affected area gently 3-4 times a day.	50.25	Skin irritation & rash.
57	METHYLERGEOMETRINE M	MEM - 1 ML	Injection	200 mcg 3-4 times/day	14.50	Headache, dizziness, hallucinations; tinnitus; nausea, vomiting, foul taste, diarrhoea; hypertension, temporary chest pain, palpitations, bradycardia.
58	MEBEVERINE HCL	COLOSPA - 135mg	Tablet	100 or 135 mg 3 times/day.	111.87	GI disturbances, dizziness, headache, insomnia, anorexia, decreased heart rate, hypersensitivity reactions.
59	MENTHOL,METHYL SALICYLATE,DICLOFENAC DIETHYLAMMONIUM,LINS EED OIL	VOLINI-40gm SPRAY	Spray	Spray over the affected area local application whenever needed.	125.00	Rash, No side effects reported, Drowsiness, Anaphylactic reactions, Allergic contact dermatitis, Pruritus, Dermatitis.
60	MALTODEXTRIN, SUNFLOWER SEED OIL, CALCIUM CAEINATE, WHEY PROTEIN ISOLATE, ISOMALTULOSE, FRUCTOSE, SOY PROTEIN ISOLATE, RAPESEED OIL, GUM ARABIC, OLIGOFRUCTOSE, MINERALS, ANTIOXIDANT, VITAMINS, CARNITINE, TAURINE AS ACTIVE INGREDIENTS.	PROHANCE-D VANILLA-400gm P	Powder	50 gm Daily or as per requirement.	600.00	Nausea, or unusual or unpleasant taste in your mouth.
61	POVIDONE -IODINE	WOKADINE	Ointment, Pessaries,	<b>Topical:</b> Clean affected area. Apply small amount topically to affected area 1—3 times daily. May cover with sterile bandage. Do not use for longer than 1 week. <b>Pessary:</b> one pessary for a period of 7 days <b>Gargle:</b> Adults, the elderly and children over 6 years of age: Use undiluted with	82.00	Skin irritation, Severe allergic reaction
		BETADINE GARGLE - 50 ML	Powder, Gargle,		83.50	
		BETADINE - V - PESSARIES	Solution		105.50	
		BETADINE SOLUTION 10% - 100ML			106.05	
		DREZ - S - 10 GM			53.24	
		DREZ - S - 15 GM			77.00	

62	POVIDONE IODINE (10% W/W)	WOKADINE-15gm OINTMENT	Ointment	Clean affected area. Apply small amount topically to affected area 1—3 times daily. May cover with sterile bandage. Do not use for longer than 1 week.	98.50	No common side effects seen.
63	POVIDONE IODINE (7.5% W/V)	POVIDINE IODINE SCRUB	Solution	Clean affected area. Apply small amount topically to affected area 1—3 times daily. May cover with sterile bandage. Do not use for longer than 1 week.	201.60	Skin irritation, Allergic Skin Reaction, Acneiform eruptions, Thyroid imbalances.
64	PRUCALOPRIDE TABLETS	PRUEASE-1mg TABLET	Tablet	Take one tablet daily.	16.35	Headache, Nausea, Diarrhea, Abdominal pain.
		PRUEASE-2 mg TABLET	Tablet	2 milligrams per day.	22.90	
65	POVIDONE IODINE-5%W/W	CIPLADINE OINMENT-15GM	Oinment	Apply sufficient quantity.	8.44	Local swelling, itchiness, redness, small blisters, and irritation.
66	PAPAIN+UREA	AUTOLYSE-15gm	Ointment	Clean the affected area with saline or with mild wound cleansing solution.Do not use hydrogen peroxide.Apply ointment directly to the lesion cover with appropriate dressing and secure into place.Wash the lesion each time when dressing is changed.	159.00	Mild stinging or burning of the skin where the medicine is applied.
67	POTASSIUM BINDER	K-BIND - 15GM	Powder	Used once/daily.	112.00	Vomiting, Stomach irritation, Nausea, Constipation, Loss of appetite.
68	PROTEIN	PROSURE-400gm POWDER	Powder	The recommended daily intake is 2 servings/day. 2 servings a day should be consumed for at least 3 weeks to see results. ProSure tastes best if served chilled. The supplement should be shaken well. Tube Feeding: Patients on tube feeding can receive the benefits of ProSure by providing 2 servings (240 mL) in conjunction with a standard tube feeding formula. In a tube feeding regimen, providing 2 servings of ProSure a day is suggested to achieve the recommended level of 2 g EPA,	789.00	Asthma, Allergic rhinitis with hives, Eczema, Increased bowel movements, Nausea, Thirst, Bloating, Cramps, Reduced appetite, Tiredness.

		PROTINEX-250gm POWDER	Powder	1 serving of 25 gm (2 to 3 heaped tablespoons) per day in 200 ml of milk.	315.00	
		ENSURE VANILLA-400gm POWDER	Powder	Directions for use: to prepare a 230 ml feeding, put 195 ml of cold water in a glass. While stirring, gradually add 6 level scoops (scoop enclosed) of Ensure powder and mix until dissolved.	480.00	
69	POLYETHYLENE GLYCOL 3350 AND ISPAGHULA HUSK GRANULES	PEGFIBER-154.812gm GRANULES	Granules	Bowel preparation: Oral: Adults: 1-2 lts of dissolved solution. Max 4 lts (Peglec soln is prepared by dissolving 1 pack of content in 2 ltrs of boiled and cooled water).	210.00	Nausea, abdominal cramping.
70	POLYETHYLENE GLYCOL ELECTROLYTES FOR ORAL SOLUTION	PEGLEC POWDER-137.15gm	Powder	Adults: 1-2 lts of dissolved solution. Max 4 lts (Peglec soln is prepared by dissolving 1 pack of content in 2 ltrs of boiled and cooled water).	316.80	Vomiting, Abdominal cramps, Stomach pain or swelling, Unusual tiredness or weakness, Frequent urge to urinate, Swelling of feet or lower legs, Muscle pain or twitching, Rapid breathing, Unpleasant taste, Mood or mental changes.
71	POLYETHYLENE GLYCOL 3350 POWDER ORAL SOLUTION	PEGMOVE-POWDER	Powder	Mix in 1/2 to 1 cup (120 to 240 ml) of liquid like water, juice, soda, coffee, or tea and drink it right away.	173.00	Stomach pain, Nausea, Diarrhoea.
72	RACECADOTRIL	ZEDOTT -100mg	Capsule	Three times a day.	212.20	Vomiting, nausea, constipation, abdominal pain, thirst, vertigo and headache.
		REDOTIL	Capsule	<b>Adult:</b> PO 100 mg 3 times/day for $\leq 7$ days.	134.00	Vomiting, nausea, constipation, thirst, abdominal pain, vertigo and headache.
73	RECOMBINANT TISSUES PL	ELAXIM - 30 MG	Injection	Usual Adult Dose for Myocardial Infarction less than 60 kg: 30 mg IV bolus administered over 5 seconds. 60 to less than 70 kg: 35 mg IV bolus administered over 5 seconds, 70 to less than 80 kg: 40 mg IV bolus administered over 5 seconds, 80 to less than 90 kg: 45 mg IV bolus administered over 5 seconds, 90 kg or greater: 50 mg IV bolus administered over 5 seconds.	30074.00	

74	SILODOSIN	RAPILIF - 8MG	Tablet	4 - 8mg OD	19.90	Retrograde ejaculation ,Prostate-specific antigen increased, Nasopharyngitis, nasal congestion, sinusitis, , priapism, pharyngeal edema.
75	SILYMARIN	SILYBON - 140 MG	Tablet	Adult dosage is 140 mg 2-3 times daily.	107.00	
		SILYBON - 70 MG			87.00	
76	SILDENAFIL (25MG)	PENEGRA-25mg TABLET	Tablet	Pulmonary arterial hypertension ▶ BY MOUTH:Adult: 20 mg 3 times a day Erectile dysfunction ▶ BY MOUTH ▶ Adult: Initially 50 mg, to be taken approximately 1 hour before sexual activity, adjusted according to response to 25–100 mg (max. per dose 100 mg) as required, to be taken as a single dose; maximum 1 dose per day	109.80	Headache. flushing of the face.
77	SODIUM ACID PHOSPHATE & SODIUM PHOSPHATE	PROCTOCLYSIS ENEMIA - 100 ML		Constipation (using Phosphates Enema BP Formula B)  Bowel evacuation before abdominal radiological procedures, endoscopy, and surgery (using Phosphates Enema BP Formula B) ▶ BY RECTUM ▶ Child 3–6 years: 45–65 mL once daily ▶ Child 7–11 years: 65–100 mL once daily ▶ Child 12–17 years: 100–128 mL once daily ▶ Adult: 128 mL daily	48.00	
78	SOLIFENACIN	URGISO-5MG	Tablet	Once a day.	263.00	Dry mouth, blurred vision, constipation, nausea, dyspepsia (indigestion), urinary tract infections, blurred vision, abdominal pain, burping or belching, heartburn, and stomach upset.

79	SODIUM PICOSULPHATE	CREMALAX-10mg TABLET	Tablet	Constipation :Child 1 month–3 years: 2.5–10 mg once daily, adjusted according to response ▶ Child 4–17 years: 2.5–20 mg once daily, adjusted according to response ▶ Adult: 5–10 mg once daily, dose to be taken at night	104.25	Vomiting, Headache, Nausea.
80	SULFASALAZINE DELAYED - RELEASE	SAZO - 500 MG	Tablet	1-2 g 4 times/day	42.25	Headache, anorexia, nausea, vomiting, diarrhoea, abdominal discomfort, photosensitivity, crystalluria, reversible oligospermia.
81	SOY PROTEIN ISOLATE, CASEINATE, CHROMIUM, ZINC, IRON, IODINE, VITAMIN C, B1, B2, B12, PANTOTHENIC ACID, VITAMIN A AND MANGANESE	VIDAVANCE VANILLA-200GM	Powder	2 scoops everyday	400.00	Nausea and stomach pain, runny nose, headache, metallic taste.
		VIDAVANCE VANILLA-400GM	Powder		755.00	
82	SOLUTION OF SORBITOL AND TRICHOLINE CITRATE	SORBILINE-100ml SYRUP	Solution	Hepatitis: 15ml t.i.d. following meals. CHILDREN: 5-10ml. Constipation: 10-20ml before breakfast. CHILDREN: 5-10ml before breakfast.	69.87	Gas, Diarrhea, Debilitation, Urge to vomit, Dizziness, Anal irritation, Abdominal cramping, Abdominal bloating, Dizzines, Acute allergic reaction.
83	TAURINE + ACETYLCYSTEINE	RENOSAVE	Tablet	Once daily.	150.00	Nausea, Vomiting, Rash, Fever, Runny nose, Diarrhea, Abdominal pain, Skin irritation, Throat irritation, Drowsiness, Chest tightness.
84	TENDOCARE TABLET CONTAINS COLLAGEN PEPTIDE(40MG), CHONDROITIN SULFATE (200MG), SODIUM HYALURONATE (30MG) AND VITAMIN C (35MG)	TENDOCARE-TABLET	Tablet	1 tablet once or twice per day	20.33	Nausea or vomiting Headache Bloating Nausea Gastrointestinal symptoms Wheezing.
85	TOCOTRIENOL	TOCOFIL-CAPSULE	Capsule	1 capsule a day		Feeling tired or weak (fatigue), stomach pain, loose stools (diarrhea), headache, nausea, or blurred vision.
86	TROXERUTIN & CALCIUM DOBESILATE	OXERUTE CD	Tablet, Cream	600-1,200 mg/day.	163.00	GI disturbances, headache and flushing.
		OXERUTE - 30 GM			118.00	



87	TROXERUTIN,CALCIUM DOBESILATE,ZINC, PHENYLEP	PILORUTE-30gm CREAM	Cream	Apply on the affected area of the skin after cleaning and drying it well. It is usually recommended to use about 2-3 times a day.	115.50	Burning or stinging sensation, dizziness, drowsiness, changing body temperatures, blurred vision or a pounding/ringing in your ears.
88	URSODEOXYCHOLIC ACID	UDILIV - 150 MG	Tablet	6 - 15 mg/kg/day in 2-3 divided doses.	136.00	Diarrhoea, pruritus, nausea, vomiting gallstone calcification.
		UDILIV - 300MG			238.00	
89	URSODEOXYCHOLIC ACID / URSODIOL (300MG)	URSOCOL-300mg TABLET	Tablet	13-15 mg/kg/day administered in two to four divided doses with food.	240.00	Upset stomach, indigestion, or vomiting. Dizziness.
90	URSODEOXYCHOLIC ACID+SILYMARIN	UDIMARIN FORTE-TABLET	Tablet	450 mg/daily.		Diarrhea, Sticky stools.
91		JOHNSON'S COTTON BUDS-15'S			20.00	
92		JOHNSON'S COTTON BUDS-30'S			36.00	
93		NEW MASK-BABY(CIPLA)			225.00	Dry/irritated throat, hoarseness, voice changes, bad taste in the mouth, runny nose, or nosebleeds.
94		NOVOFINE NEEDLE			12.50	Redness, swelling, or itching at injection sites.
95		STAY FREE (DRY WINGS)			30.00	
96		STREPSILS			2.92	
97	1 X 2ML DENSITY GRADIENT + 1 X 5ML WASH MEDIA	DOUBLE LAYER DENSITY GRADIENT MEDIA(5ml+2ml)	CULTURE MEDIA		590.00	
98	DEVICE	PENLANCET	GLUCOMETER DEVICE		200.00	
99	GRADIENT	CELL MED DENSITY GRADIENT 40%&80%	GRADIENT		363.00	
100	MEDIA	CELL MED SPERM WASH MEDIA-4.5ML	MEDIA		227.00	
101	DUTASTERIDE & TAMSULOSIN	DUTAS-T	Capsule	Once a day	572.25	Headache Sexual Dysfunction Itching Of Skin Runny Nose
102	DIADZEIN, GENISTEIN & PALMITOYLETHANOLAMIDE	PALMIGES	Tablet	Twice a day	302.00	Constipation Bloating Nausea Vomiting Fever Itching
103	L-GLUTATHIONE	THIOTRES-500mg	Tablet	One or Two tablets a day	1197.00	Skin rash. Fever. Diarrhoea. Dry mouth. Difficulty in breathing & Cough
104	SILODOSIN & DUTASTERIDE	SILDOO-D 8mg	Tablet	500 mcg daily	399.25	Orgasm with reduced or no semen Trouble getting or keeping an erection (impotence) Decrease in sex drive (libido) Ejaculation problems Enlarged or painful breasts Dizziness

105	TOLVAPTAN	HYPONAT-O-15mg	Tablet	Once a day	679.95	Frequent urge to urinate Nocturia (increased urge to urinate during the night) Dryness in mouth Increased thirst
106	MIRABEGRON & SOLIFENAC	EXENA-S	Tablet	Once a day	275.00	Constipation, Headache, Dry mouth Blurred vision, Vomiting
107	MIRABEGRON & SILODOSIN	RAPILIF-M-25mg	Tablet	8 mg orally once daily	235.00	Retrograde ejaculation,Dizziness Diarrhea,Orthostatic hypotension (sudden lowering of blood pressure on standing) Headache.
		RAPILIF-M-50mg	Tablet		285.00	
108	AVANAFIL	AVANAIR-100mg	Tablet	Once a day	220.00	Nausea, vomiting, headache, skin color changes, increased/decreased sexual interest, oily skin, hair loss, and acne.
		AVANAIR-200mg	Tablet		320.00	
109	DEFLAZACORT & DAMSULOSIN	TAMSUKEM PLUS	Tablet	Once a day	800.00	Runny nose,Ejaculation disorder,Dizziness
110	MENTHOL	SENZICOOL	Lozenges	Every 2 hours as needed	80.00	hiccups, heartburn, and nausea.
111	ULINASTATIN	U-TRYP-100000IU	Injection	1,000,000 IU every 8 h and tapered down to 500,000 IU every 8 h after 4-7 days	4264.70	Chills, Chest pain, Nausea, Vomiting, Abdominal pain, Fast or slow heartbeat, Itching of skin.
112	COMPARTMENT A: CALCIUM CHLORIDE, MAGNESIUM CHLORIDE, DEXTROSE, OR D- GLUCOSE MONOHYDRATE,LACTIC ACID, SODIUM CHLORIDE, POTASSIUM CHLORIDE, SODIUM BICARBONATE,  COMPARTMENT B: SODIUM CHLORIDE,SODIUM BICARBONATE,POTASSIU M CHLORIDE,CALCIUM, MAGNESIUM,SODIUM, CHLORIDE,LACTATE BICARBONATE,POTASSIU M DEXTROSE,THEORETICAL	PRISMASOL BAG-5LITRE	Tablet	Immediately before use, remove the overwrap from the bag and mix the solutions in the two different compartments. After removing the overwrap, inspect the bag for leakage by pressing firmly on the bag. Discard the bag if any leakage is detected since sterility cannot be assured. As soon as the overwrap is removed, the reconstitution of compartments A and B should be done and the mixed solution should be used immediately.  After removal of the overwrap, the solution is stable for 24 hours including the duration of the treatment	2724.00	Hives, difficulty breathing, swelling of your face, lips, tongue, or throat, severe dizziness, rapid and shallow breathing,nausea, vomiting

113	SOLIFENACIN SUCCINATE	VESILIFE-5mg	Tablet	Once a day	270.00	Dry mouth Blurred vision Nausea Constipation, indigestion, stomach pain Sense of fullness in the stomach, burping, heartburn.
114	ALLANTOIN-0.01G + ALLIUM CEPA EXTRACT- 0.1G + HEPARIN SODIUM- 50IU	SCAREND-15gm	Gel	A small quantity apply affected area	289.85	Pruritus (severe itching of the skin),Erythema (redness),Telangiectasia (visible dilation of blood vessels),Scar atrophy (scar lower than skin surface)
115	TURMERIC	CURCUMIN-250mg	Tablet	Once a day	460.00	stomach upset, nausea, dizziness, or diarrhea.
116	L-ARGININE CHONDROITIN SULPHATE SODIUM COLLAGEN PEPTIDES TYPE I SODIUM HYALURONATE MAGNESIUM STEARATE ASCORBIC ACID SILICONE DIOXIDE STABILISER BINDER MICROCRYSTALLINE CELLULOSE GLAZING AGENT	TENDOCARE FORTE	Tablet	Three times a day	600.00	Allergic reactions,Blue color or flushing or redness of skin,Cough,Bloating.
117	ADEMETHIONINE	CIRROSAM-400MG	Tablet	Twice a day	1082.00	Nausea,Diarrhea,Upset stomach,Constipation,Dizziness.
118	IOHEXOL	OPTISCAN-350-50ml VIAL	Injection	Single dose	1400.00	Diarrhea, nausea, vomiting, stomach pain, gas, or headache
		OPTISCAN-350X100ml-VIAL	Injection	Single dose	2780.00	
119	SODIUM CHLORIDE	FREEFLEX-500ML INFUSION(SOL	Injection	Single dose	87.83	Pain at the site of injection Dehydration of internal organs particularly in the brain which may lead to blood clots or internal bleeding Signs of allergic reactions (such as low blood pressure, increased body temperature tremors, chills hives, rashes and itchy skin).
120	LIPIDS 3.8% SOYA OIL FISH OIL - RICH IN OMEGA- 3 ACIDS MEDIUM CHAIN TRIGLYCERIDES OLIVE OIL GLUCOSE 12.7% GLUCOSE MONOHYDRATE ELECTROLYTES 0.7%	SMOFKABIVEN - 1970ml(INTRA VENOUS INFUSION).	Injection	12-24 hours. Maximum daily dose	5459.00	nausea and vomiting,flatulence abdominal pain,swelling of hands, ankles or feet,fast heart beat difficulty in breathinglack of appetite.

## EYE, EAR, NOSE, MOUTH & THROAT

1	CLOTRIMAZOLE	CANDID - 10ml	Drops,		53.00	
2	CIPROFLOXACIN + DEXAMETHASONE	CIPLOX-D-10ml EYE DROPS	Drops,	Four drops into the affected ear twice daily, for seven days.	18.15	Bone marrow depression, HS reaction, Grey baby syndrome, Superinfections.
3	NEPAFENAC	NEPALACT	Drops	1 drop, 0.3% should be applied to the affected eye one-time-	159.00	Application site irritation.
4	TOBRAMYCIN	TOBAREN - 5ml	Drops	Local treatment of infections ▶ TO THE EYE ▶ Child 1–17 years: Apply twice daily for 6–8 days ▶ Adult: Apply twice daily for 6–8 days Local treatment of infections (severe infection) ▶ TO THE EYE ▶ Child 1–17 years: Apply 4 times a day for first day, then apply twice daily for 5–7 days ▶ Adult: Apply 4 times a day for first day, then apply twice daily for 5–7 days.	61.78	
5	TIMOLOL MALEATE	TIMOLET-0.5-5ml	Drops	Two to three times a day.	66.89	Blurred vision, Burning or stinging sensation at application site, Cataract, Headache, Itching, Dizziness.
		IOTIM-0.5%-5ml	Drops	Reduction of intra-ocular pressure in primary open-angle glaucoma Adult: Apply twice daily.	55.18	Burning eyes, Stinging in the eyes.
6	TRETINOIN	RETINO-A-20 gm	Cream	usually once daily at bedtime	135.00	Itching, redness.
7	SODIUM CHLORIDE (5% W/V)	SOLINE-10ml	Eye Drops	1 or 2 drops every 3 or 4 hours.	92.00	Burning sensation in eye
8	SODIUM CHLORIDE	NASIVION-S-10ml	Drops	Instil 2-3 drops in each nostril 3-4 times daily	36.19	Allergic contact dermatitis, Nausea, Vomiting, Diarrhea, Muscular twitching, Inflammation of the gastrointestinal tract, Congestion, Eye irritation, Restlessness, Convulsions.
		NASOCLEAR-20ml	Drops	2 drops thrice daily for 1 week	40.00	Emesis, Collapse, Nausea, Vomiting, Diarrhea, Muscular twitching, Inflammation of the gastrointestinal tract, Congestion, Eye irritation, Restlessness.

9	SODIUM CHLORIDE \$ SALINE 0.9	NASOCLEAR MIST SPRAY-100ml	Spray	Adults and Children: Apply 2-4 sprays in each nostril 3 times a day.	276.00	Confusion, constipation, dizziness (mild), drowsiness (mild), headache, nausea or vomiting, stomach pain.
10	SODIUM CARBOXYMETHYL CELLULOSE	LUBI-FRESH-10ml	Drops	drops can be used as often as required.	98.00	Stinging in the eyes, Burning eyes, Eye redness, Allergic reaction in eye.
11	BENZOCAINE + CHLORBUTOL + PARADICHLORO BENZENE + TURPENTINE OIL	SOLIWAX-10ml	Drops	Instill 5-10 drops into the affected ear 2-3 times a day and allow contact for 10-20 minutes and overnight. Continue for 2-3 days, if necessary.	111.00	Burning/stinging/redness/new pain in or around the ears.
12	LOTEPREDNOL ETABONATE	LOT-5ml	Drops	One to two drops.	141.25	Watery eyes, Irritation in the eyes, Eye itching, Foreign body sensation.
13	OXYMETAZOLINE HCL	NASIVION (ADULT) - 0.05 %	Drops	Adults and children 6 years of age: 2 to 3 drops of in each nostril twice daily, in the morning and evening. Children 2-5 years of age: 2 to 3 drops of in each nostril twice daily in the morning and evening.	63.00	Nasal irritation.
		NASIVION (PAED) - 0.025 %			46.00	
		OTRIVIN-ADULT	Spray	Adult Dose for Nasal Congestion: 2 or 3 sprays in each nostril not more often than every 10 to 12 hours  Pediatric Dose for Nasal Congestion: 6 years or older : 2 or 3 sprays in each nostril not more often than every 10 to 12 hours.	66.50	Temporary burning, stinging, dryness in the nose, runny nose, and sneezing.
14	PARADICHLORO BENZENE BENZOCAINE, CHLORBUTOL	OTOREX-10ml	Drops	2 or 3 drops should be instilled into the ear three or four times daily; alternatively, a gauze wick kept saturated with the drops may be inserted into the external auditory meatus.	68.72	
15	PREDNISOLONE (TOPICAL) (1% W/V)	PRED FORTE-10ml	Drops	Adult: PO Allergic and inflammatory disorders 5-60 mg/day, in 2-4 divided doses.	34.18	Eye irritation, Watery eyes, Burning sensation.

16	FLUOROMETHOLONE	FML-5ml	Eye drops	TO THE EYE ► Child 2–17 years: Apply every 1 hour for 24–48 hours, then reduced to 2–4 times a day ► Adult: Apply every 1 hour for 24–48 hours, then reduced to 2–4 times a day	87.17	
17	HOMATROPINE HBR	HOMIDE DROPS-5ml	Drops	apply it usually 2 to 3 times daily, up to every 3 to 4 hours.	30.61	Burning/stinging/redness of the eye, eye irritation, or temporary blurred.
18	HYPROMELLOSE + SODIUM CHLORIDE + POTASSIUM CHLORIDE + SODIUM ACETATE + CALCIUM CHLORIDE + SODIUM CITRATE + MAGNESIUM CHLORIDE	LACRIGEL-5gm	Ointment	The usual dose is 1 or 2 drops in each affected eye 3 times daily.	72.30	Mild eye burning or irritation, itching or redness of your eyes, watery eyes, blurred vision or unpleasant taste in your mouth.
19	POTASSIUM NITRATE 3 % + SODIUM FLUORIDE	SENQUEL-AD-100ml	Mouthwash	Recommend that you swish the product around in your mouth for 30 seconds to one minute, then spit it out.	55.50	Dizziness, laboured breathing, abdominal pain, blue fingernails, lips or skin, diarrhoea. a headache, convulsions, confusion, nausea, unconsciousness.
20	POTASSIUM NITRATE 5 % W/W + SODIUM MONOFLUORO PHOSPHATE	SENQUEL-F-50gm*	Paste	For external use .once daily.	71.00	Abdominal bloating, Dental fluorosis.
21	TRAVOPROST	LUPITROS-3ml	Drops	once a day in the evening.	180.50	Eye irritation
22	CHLORHEXIDINE GLUCONATE	CLOHEX ADS-200ml	Mouthwash	Swish for 30 seconds to 1 minute and spit it. Do not swallow. Avoid eating, drinking, or smoking for at least 30 minutes after rinsing.	149.00	Respiratory tract infection severe, Skin peeling severe rare, Hypersensitivity severe rare, Inflammation of gum severe, Tooth pain, Dry mouth, Staining of tooth, Altered sense of taste.
23	CALCIUM & VITAMIN D	JUSDEE 400-30ml	Drops	10mg/day	74.26	Nausea/vomiting, constipation, loss of appetite, increased thirst, increased urination, mental/mood changes, unusual tiredness.
24	PILOCARPINE NITRATE 0.5%	CARPINOL-1ml	Injection	Once a day	18.00	Headache, Itching or burning Diarrhoea, Tremor Nausea or vomiting, Chills Flushing, Weakness.
25	CARBOXYMETHYLCELLULOSE	REFRESH-10ml LIQUIGEL	Drops	drops may be used as often as needed. squeeze out 1 or 2 drops as needed. Look down and gently close your eye for 1 or 2 minutes.	180.07	Eye pain, change in vision, continued eye redness/irritation.
26	FLURBIPROFEN SODIUM	FLUR DROPS - 10ml	Drops	Instill 1 drop into the eye 4 times/day for 1-3 wk starting at 24 hr after surgery.	111.00	Irritation, itching, redness of eyes.

27	CARBOXY METHYL CELLULOSE	JUST TEARS-10ml	Eye drops	1-4 applications/day.	81.00	Eye irritation, Burning sensation.
		LACRIMOS 0.5% -10ml			79.00	
		LUBREX -10ml			89.89	
28	CHLORHEXIDINE	HEXIDINE-160ml	Mouth wash	Rinse w/ 10 mL for about 1 min bid. As spray: Apply bid. Max: 12 actuations bid. As gel: Brush teeth w/ 1 inch of gel once or bid for about 1 min.	69.00	Dry mouth, Mouth irritation.
		REXIDIN-120ml			72.00	
		CLOHEX PLUS-150ml			86.50	
29	CHOLINE SALICYLATE	ZYTEE-10ml	Gel	Apply gel on affected area once/day	70.00	Dry mouth, Mouth irritation
		DOLO-CT GEL-10gm			49.00	
30	CHLORAMPHENECOL	CANDIBIOTIC PLUS-5ml	Capsule	Adult: 250 -500 mg 6 hrly Paed: 25-50mg/kg/day.	62.00	Swelling, Dermatitis. pruritis in the ear.
31	GATIFLOXACIN	HI GATI - 5gm	Eye Drop, Eye ointment	Adult and child (above 1 year): Instill one drop upto 8 times daily for first 2 days.	51.00	Eye discharge, Itching eye, Tearing.
32	GATIFLOXACIN+PREDNISOLONE	GATE-P	Eye Drop	One / two drop in the affected eyes	33.50	Fluid retention, alteration in glucose tolerance, increased appetite, high blood pressure.
33	MOXIFLOXACIN HCL	MOXIBLU-5ml	Tablets, Injection, Drops, Ointment	Adult: 400mg OD	59.00	QT prolongation, phototoxicity, GI upset
		MOXAP			98.00	
		MOXICIP			64.00	
		MOXIGRAM-LX-5ml			142.00	
34	MOXIFLOXACIN (0.5 %W/V) + DEXAMETHASONE (0.1 %W/V)	MOXIGRAM-DM-5ml	Eye Drop	Two to three times a day.	106.96	Eye irritation, Eye discomfort, Blurred vision.
35	MOXIFLOXACIN & PREDNISOLONE	ANZMOX-PD-10ml	Drops	Instill one or two drops in the affected eye 2 to 4 times a day	38.00	Blurred or decreased vision. change in vision. chest tightness. hives, itching, skin rash.
36	NEOMYCIN	NEOSPIRIN	Eye Ointment		40.00	
37	OLAPATADINE HCL	OLOPAT OD-3ml	Eye drops	Adult & Child: ≥3yrs: As 0.2% solution : 1 drop in the affected eye(s) once daily.	119.00	Eye irritation, burning sensation.
		OLOPAT-5ml	Drops			Headache, blurred vision, burning/stinging/redness/dryness of the eye, eyelid swelling, or a feeling as if something is in your eye.
38	NATAMYCIN (5%)	NATAFORCE	Eye Drops	Adult: As 5% ophth susp: Instill 1 drop in the conjunctival sac every 1-2 hr, reduce to 1 drop 6-8 times daily after 3-4 days.	111.09	mild eye irritation or discomfort (redness stinging or burning)

## DERMATOLOGICAL (EXTERNAL PREPARATION)

1	BETAMETHASONE + NEOMYCIN	BETNOVATE-N-20gm	Cream	Two or three times a day.	330.30	Skin itchiness, drying or cracking of skin, acne, changes in skin colour, persistent infections.
2	BETNOVATE VALERATE CREAM I,P	BETNOVATE - 20gm	Cream	apply to affected area 2-3 times per day	20.00	Itching, redness.
3	CLOTRIMAZOLE	CANDID - B -20gm	Mouth paint,		89.50	Peeling, swelling, itching,
		CANDID -50gm			104.00	
4	ESTRADIOL VALERATE	EVALON - 15gm	Cream	1 applicator upto the ring /day for 2-3 weeks;then 1 applicator upto the ring twice a week	289.00	
5	GANCICLOVIR	VIRSON GEL	Gel	five times a day	111.00	Chills, headache, shortness of breath diarrhea, abdominal and stomach pain, nausea and vomiting, back pain seizures, swelling of legs, itching, decreased urine output.
6	KETOCONAZOLE	KZ - 75gm		Adult: 200 -400 mg OD	155.00	Urticaria, Anaphylatic Reactions, GI upset, Constipation, Headache, Dizziness, Hepatitis addison's disease.
		SCALPE - 75ml	Lotion, Soap, Cream		250.00	
		KZ			185.00	
		TARRY SHINE-100ml		Adult: As 2% shampoo: Apply on scalp once daily for up to 5 days. For prophylaxis: use once daily for a max of 3 days before exposure to sunlight.	230.00	Application site reactions (burning, irritation, itching and redness).
7	SERTACONAZOLE NITRATE	ONABET - 15gm	Cream	Adult: Apply to affected area bid for 4 wk.	130.00	Swelling, itching, burning sensation, stomach pain,redness, oozing, dry skin.
		ONABET - 30gm			230.00	
		ONABET-B - 10gm			118.00	
8	TERBINAFINE (1% W/W)	TYZA-30gm	Cream, Tablet, Cream	Adult: Apply once or bd to affected area(s) for 1-2 weeks Adult: 250 mg OD.	135.00	Skin peeling, Itching.
		TYZA-15gm			69.00	Headache, GI upset, Taste disturbance, insomnia, Abdominal pain, Diarrhoea, Cutaneous rash, URTI, Visual disturbances, cutaneous angioedema.
9	ADAPALANE	ADIFF-15gm	Gel	Adult: Apply thinly onto affected areas once daily at night after cleansing.	130.00	Burning , Itching, Redness.
10	ADAPALANE + CLINDAMYCIN	DERIVA CMS -15gm				
11	ALOE VERA GEL, CALAMINE ,LT LIQ PARAFFIN	CALAPURE-A-50ml	Lotion	apply 1-4 times/day	85.00	Allergic skin reaction.
		CALOSOFT-AF--50ml			89.00	



12	KETOCONAZOLE	KERAGLO AD-75ml	shampoo	Apply to wet hair, massage into scalp, leave on hair for 5 mins and rinse thoroughly with water.	273.90	Nausea, headache, diarrhea, stomach pain and dizziness.
13	ALOEVERA TOPICAL, SQUALENE TOPICAL AND VITAMIN E	VENUSIA-75gm	cream	apply over the body both morning and evening.	265.00	Erythema multiforme, No known side effects.
14	ALOE VERA. LIGHT LIQUID PARAFFIN. GLYCERINE. CALAMINE. PRAMOXINE HYDROCHLORIDE.	VENUSIA SOFT-100ml	Lotion	Gently spread it onto the dry-itchy area of your skin, use it 2-3 times a day.	333.00	Dry skin, skin reddening, Skin burning, Skin itching, skin rash.
15	ALMOND BUTTER TOPICAL, CETYL ALCOHOL TOPICAL, COCOA BUTTER TOPICAL, CYCLOMETHICONE TOPICAL, DIMETHICONE TOPICAL, DISODIUM EDETATE TOPICAL, EMULSIFYING WAX TOPICAL, FRAGRANCE TOPICAL, GLYCERIN TOPICAL, MANGO BUTTER TOPICAL, METHYL PARABEN TOPICAL, PROPYLENE GLYCOL TOPICAL, PURIFIED WATER TOPICAL, SHEA BUTTER TOPICAL, STEARIC ACID TOPICAL AND ZINC OXIDE TOPICAL	VENUSIA MAX-150gm	cream	topical application all over the body after bath apply twice a day.	355.00	Burning of skin, Redness of skin, Inflammation of chinchilla middle ears, Contact dermatitis, Mild lactic acidosis, Burning, Stinging, Erythema, Skin irritation, Pruritus.

	BUTTER TOPICAL, METHYL PARABEN TOPICAL, PROPYLENE GLYCOL TOPICAL, PURIFIED WATER TOPICAL, SHEA BUTTER TOPICAL, STEARIC ACID TOPICAL AND ZINC OXIDE TOPICAL	VENUSIA MAX-300gm	Lotion.	topical applicaltion all over the body after bath apply twice a day.	468.00	
16	MINOXIDIL(10.0 %)	TUGAIN 10% FOAM-60gm	Foam	Adults—Apply half a capful to th	1345.30	Headache Burning sensation Excessive hair growth Itching of skin, rash, hives (raised rashes) Dry skin
17	FUSIDIC ACID 2.0 W/W	MOMATE-F -10gm	Cream		145.00	
18	AQUA, CETYL ALCOHOL, PROPYLENE GLYCOL, SODIUM LAURYL SULPHATE, STEARYL ALCOHOL, POLYSORBATE 20, METHYL PARABEN, PROPYL PARABEN, BUTYL PARABEN.	CETAPHIL(oily skin)-125ml	Face Wash	Twice a day	545.00	oozing, heat, swelling, redness, or pain.
19	PURIFIED WATER, ALOE VERA, PROPYLENE GLYCOL, EMULSIFYING WAX, SQUALENE, GLYCERIN, CYCLOMETHICONE, STEARIC ACID, VITAMIN E ACETATE, CETYL ALCOHOL, DIMETHICONE (10 CST), SODIUM METHYL PARABEN, TROLAMINE, DISODIUM EDTA, FRAGRANCE AL 012982, SODIUM PROPYL PARABEN, BUTYLATED HYDROXYANISOLE	VENUSIA-100gm	Cream	Apply Daily	438.50	Skin Irritation, Stinging

20	CYCLOMETHICONE, GLYCERINE, SHEA BUTTER, COCOA BUTTER, MANGO BUTTER, CETANOL, STEARIC ACID, DIMETHICONE AND ZINC OXIDE	VENUSIA DERM-75gm	Cream	Apply Daily	274.00	Skin Irritation, Stinging
21	WATER, CETEARYL ALCOHOL, ETHANOLAMINE, STEARTRIMONIUM CHLORIDE, STEARAMIDE DEA OTHER INGREDIENTS : CETEARETH-20, TOLUENE-2, 5 DIAMINE SULFATE, STEARETH-21, STEARAMIDOPROPYL DIMETHYLAMINE, DIMETHICONE, AMODIMETHICONE, BOSEWELLIA SERRATA EXTRACT, MINERAL OIL, RESORCINOL, OLETH-2, LANOLIN, PORTULACA OLERACEA EXTRACT, BEHENTRIMONIUM CHLORIDE, GLYCERYL STEARATE SE, SORBITAN SESQUIOLEATE, SODIUM SULFITE, M-AMINOPHENOL, CYSTEIN HCL, ALLANTOIN, O-AMINOPHENOL, TETRASODIUM EDTA, POLYQUATERNIUM-10, PANTHENOL, ASCORBIC ACID, FRAGRANCE, OLEA	CUTICOLOR HAIR COLOUR	Hair colour	Once in 15 days	1433.50	Causing allergy.
22	UREA+LACTIC ACID	MOISTUREX-100gm	Cream	A thick layer Cream is applied	200.00	Rash, diarrhoea, constipation, abdominal
23	TACROLIMUS	TACORYZA-0.01-10gm	Cream	Apply a thin layer to the affected areas of skin, usually twice daily	388.70	Application site reactions (burning, irritation, itching and redness).

		TACROTOR-0.03%-10gm	Ointment		154.40	
		TACROZ FORTE-0.1%-10gm	Ointment		461.50	
		TACROZ-0.03%-10gm	Ointment		177.00	
24	PODOPHYLLUM RESIN BENZOIN ALOES SOLUTION	PODOWART-10ml PAINT	Solution	Twice a day and once a week ,Clean and dry the skin area to be treated. do not wash the treated area after immediately applying Podowart Paint Lotion.	154.00	Severe burning, stinging, itching, redness, swelling, or skin changes where the medicine was applied, numbness, burning, pain, or tingly feeling, easy bruising or bleeding, weakness, fever or severe constipation or stomach cramps.
25	PERMETHRIN TOPICAL & 50MG/GM	PERLICE-120gm	Cream	Apply Perlice Creme/Rinse 1% Cream to the hair, or skin, depending on the location of infection. Avoid its contact with eyes. The same treatment can be repeated after seven days if required.	196.22	Mild burning orstinging.
		PERMITE-60gm		Apply the medicine from your head to the soles of your feet, including under your nails and in skin folds such as between the toes, as directed. Massage the cream into the skin. Do not use more medication than prescribed. Wash off the cream after 8-14 hours by showering or taking a bath.Demonstrable living mites after 14 days indicate that retreatment is necessary.Permethrin cream is safe and effective in pediatric patients two months of age and older.	106.17	
26	TEA TREE OIL, ALOE VERA, AND ALLANTOIL	ACNESTAL-75gm	Soap	Apply soap on face and shoulder areas Leave it for a minute Wash with water and gently dry the area Can be used for 2- 3 times a day.	88.00	Peeling, itching, irritation, and reddened.
27	SALICYLIC ACID	SALICYLIX - SF - 6 % -50gm	Ointment, Shampoo	1 to 2 times daily for up to 2 weeks for corns and calluses and 12 weeks for warts.use as a shampoo, wet hair and scalp. Lather well and leave on as you have been told. Rinse fully.	94.00	Itching, redness.
		DERSOL - 24% -25gm			99.00	
		AKRALIC - 12 %			65.00	
		AKRALIC - 6 %			45.00	

28	SALICYLIC ACID AND LACTIC ACID	SALACTIN-10ml	Lotion.	First wash the affected area. You may soak it in warm water for about 5 minutes to soften it. Dry thoroughly. Removing the dead skin helps the medication to work better. Do not try to rub the wart, corn, or callus off. Using the applicator provided, apply a thin coat of the medication (one drop at a time) to cover the entire wart, corn, or callus. To minimize irritation, be careful not to get the liquid on surrounding skin. Let dry. Repeat this procedure 1 to 2 times daily as needed until the wart, corn, or callus is removed. For warts, you may use this product for up to 12 weeks. For corns and calluses, you may use it for up to 2 weeks. Wash hands after use.	271.00	Rash, diarrhoea, constipation, abdominal pain, headache, sleeping disorders, dizziness, galactorrhea, gynecomastia.
29	SALICYLIC ACID TOPICAL,SULFUR TOPICAL.	ACNIL SOAP	Soap	Adults: .Wet the face or body with water Apply the soap and gently work up the lather. Rinse thoroughly. Can be used for 2- 3 times a day.	60.00	Skin irritation, Inflammation of skin, Rash, Contact dermatitis, Irritation to the gastrointestinal tract, Nausea, Vomiting, Diarrhea.
30	SILVER NITRATE	SILVEREX-20gm	Gel	Apply the gel on the wound/ affected area at least once daily	99.00	Nausea, vomiting, diarrhoea; hypersensitivity, skin reactions; haematuria; crystalluria; thrombocytopenia, leucopenia, eosinophilia.
31	SULPHONATED SURFACTAN	ACNE AID-100gm	Soap	Make a lather with warm water and massage into the affected skin using the finger tips or a facecloth. Rinse and repeat as necessary.	125.80	Peeling, itching, irritation, and reddened.
32	SILVER NITRATE	SILVEREX-25gm	Cream	on the affected area 2-3 times per week for 2-3 weeks	83.50	Itching, redness.
33	SUNSCREEN LOTION	PHOTOBAN-30-100gm AQUAGEL	Gel	apply over the face and neck,half an hour before before going out into sunlight.	339.50	Dermatitis, Contact dermatitis, Burning of skin, Irritation of skin, Skin irritation.

34	SODIUM TALLOWATE, SODIUM COCOATE, WATER, GYLCE RIN, FRAGRANCE,SODIUM CHLORIDE, SODIUM HYDROXIDE, TETRASODIUM ETIDRONATE, O-TOLYL BIGUANIDE,TITANIUM DIOXIDE,	TETMOSOL-100gm	Soap	For external application. Apply soap till lather comes,rinse with water.	61.00	itching, skin irritation, inflammation, rashes, redness on skin.
35	PIRFENIDONE (200MG)	FIBERDONE-200mg	Tablet	Adults for treating Idiopathic Pulmonary Fibrosis is 267 mg orally thrice a day over a period of 1-7 days and 534 mg from days 8-14.	192.95	Unusual tiredness, weakness, dizziness upon standing, diarrhea, weight loss, menstrual period changes, decreased sexual interest or ability, enlarged/tender breasts in men.
36	PIMECROLIMUS TOPICAL	PICON-1%-10gm	Cream	Apply a thin layer to the affected areas of skin, usually twice daily.	400.00	Itching, rash, and redness.
37	SODIUM FUSIDATE	CHITOMESH-SF-10gm	Cream, Ointment	Apply a small amount to the affected area 3 or 4 times daily.	100.00	Itching, redness.
		FUCIDIN-15gm			136.50	
38	PALM & COCONUT OIL, SHEA BUTTER, ALOE VERA	GLADDEW BABY-75gm	Soap	Wet the face or body with water Apply the soap and gently work up the lather. Rinse thoroughly	109.00	Drowsiness, dizziness, hypotension or a headache.
		GLADDEW-75gm	Soap		105.00	
39	MOMETASONE FUROATE	MOMATE-40gm	Ointment	Thin film of medication to the affected area once daily.	317.50	Rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.
		MOTIMESH-5gm	Cream	Apply a thin film of Mometasone Furoate Cream to the affected skin areas once daily.may be used in pediatric patients 2 years of age or older.	75.00	Application site reactions (burning, irritation, itching and redness).
		MOMATE-5gm	Cream	Apply a thin layer of this medicine on the affected area once daily	170.00	Headache, dry or sore throat, unpleasant taste and smell, sneezing, a fungal infection in the mouth.
40	MOMETASONE FUROATE 0.1 %W/V	HH SONE-10ml	Lotion	Apply affected area one or two times a day.	158.00	burning –stinging –itching –tingling
		MOMATE - 30ml	Cream, Lotion, Nasal spray	Prophylaxis and treatment of allergic rhinitis ► BY INTRANASAL ADMINISTRATION ► Child 6–11 years: 50 micrograms daily, dose to be	250.00	Itching, redness.
		MOMATE -15gm			171.50	
		MOMEFLO -NASAL SPRAY			202.50	
		HH SONE-10gm			116.00	
		HH FUDIC -10 GM			208.00	

41	MINOXIDIL	MINICHEK-5%-60ml SOLUTION	Solution	Clean and dry the scalp area before applying the medication. You may apply this product to damp hair. To use the solution, fill the applicator with 1 milliliter of medication (to the 1 milliliter line), or use 20 drops. Part your hair in the area of thinning and apply the solution evenly to the affected area of the scalp. Gently rub in. Allow the solution to dry completely before using other styling products (e.g., gels, mousse) or before going to bed.	650.00	Application site irritation, Itching, Headache.
		ZENOXIDIL-2%-60ml GEL	Topical solution	<p>FOAM men Androgenetic alopecia ► TO THE SKIN ► Adult: Apply 0.5 capful twice daily, to be applied to the affected areas of scalp; discontinue if no improvement after 16 weeks</p> <p>SOLUTION Androgenetic alopecia ► TO THE SKIN ► Adult: Apply 1 mL twice daily, to be applied to the affected areas of scalp; discontinue if no improvement after 1 year</p> <p>FOR WOMEN REGULAR STRENGTH Androgenetic alopecia ► TO THE SKIN ► Adult: Apply 1 mL twice daily, to be applied to the affected areas of scalp; discontinue if no improvement after 1 year</p>	225.00	Burning, stinging, or redness at the application site.

		MINTOP 2% - 60 ML LOTION	Lotion	apply 1 ml to the scalp bd	233.00	Itching, redness.
		MINTOP FORTE 5% - LOTION			528.00	
42	GLYCERINE, VEG OILS, REFINED PERFUMES	GLYCEP SOAP-75gm	Soap	apply soap to wet body and facetill lather comes,then rinse with water.	60.00	No common side effects seen.
		GLYCEROL-100ml	liquid	adult laxative for constipation: The common dose of glycerol is a 2-3 grams in suppository form or a 5-15 mL enema. For children younger than six years old, the dose is a 1-1.7 grams as a suppository or a 2-5 mL enema.	40.00	Headaches,dizziness, bloating, nausea, vomiting, t hirst, and diarrhea.
43	MONOSULFIRAM	SCABELICE MEDICATED-75GM	Soap	Once or two times day	48.40	
44	PERMETHRIN	PERLICE - 60GM CREAM	Cream		119.15	
45	MINOXIDIL & FINASTERIDE	MINI CHEK-F-5%-60ml SOLUTION(Minoxidil & Finasteride)	Solution	1ml of liquid solution applied directly to the scalp two times per day	946.05	Burning, irritation, itching and redness.
46	HALOBETASOL PROPIONATE	HALOX-30ml LOTION	Lotion.	One to two times daily	199.00	Itchiness, dryness, redness, burning and stinging sensation, stretch marks, increased hair growth, folliculitis, hives, warm skin, acne and skin discoloration.
47	METHOXSALEN TOPICAL SOULUTION	OCTAMOP-30ml LOTION	Lotion	It is applied on the skin along with ultraviolet-A irradiation.	118.00	Chewing sugar-free gum may help. Sunburn. Skin irritation. Itching.
48	ISOTRETINOIN	TUFCANE 10 MG	Capsules	Adult: PO Initial: 0.5-1 mg/kg/day in 2 divided doses.	114.30	Dryness of mucous membranes, dryness of skin with scaling, fragility, erythema, cheilitis, pruritus, epistaxis, conjunctivitis, dry sore mouth and palmo-plantar exfoliation.
		TUFCANE 20 MG			142.00	
49	CLOBETASOL PROPIONATE &FUSIDIC ACID CREAM	CLONATE - F - CREAM	Cream	Apply thinly onto the affected area once daily or bid, reduced if necessary. Max dose: 50 g wkly. Max duration of therapy: 4 wk.	145.00	Irritation, Itching, Burning
50	CLOBETASOL PROPIONATE	TENOVATE - 15 GM	Ointment	Apply thinly onto the affected area once daily or bid, reduced if necessary. Max dose: 50 g wkly. Max duration of therapy: 4 wk.	56.30	Irritation, Itching, Burning
51	MODIFIED LANOLIN	NIP CARE OINTMENT	Ointment		118.00	Itching, redness.



52	GLYCERIN HYDROGENATED POLYISOBUTENE CETEARYL ALCOHOL CETEARETH-20 MACADAMIA NUT OIL DIMETHICONE TOCOPHERYL ACETATE STEAROXYTRIMETHYLSIL ANE (AND) STEARYL ALCOHOL PANTHENOL FARNESOL BENZYL ALCOHOL PHENOXYETHANOL	CETAPHIL DAM-100gm LOTION	Lotion	Recommended for gentle daily cleansing of damaged skin associated with common skin conditions and following dermatological treatments	373.00	Allergic reaction: hives; difficult breathing; swelling of your face, lips, tongue, or throat.
53	AZELAIC ACID	AZIDERM-10% - 15GM	Cream, Soap	Apply thinly into the affected areas bid (morning and evening) after cleansing for 6months.	159.00	Itching, redness.
		AZAC SOAP - 75 GM			70.00	
54	ALPHA LIPOIC ACID BENFOTIAMINE CHROMIUM PICOLINATE FOLIC ACID INOSITOL MECOBALAMIN	BENFOMET FORTE	Tablet	Once daily.	255.00	Diarrhea, Nausea, Vomiting, Stomach upset, Constipation.
55	ALMOND, ESSENTIAL OIL AND VITAMIN E	KERA-XL HAIR GROWTH-60ml	Hair Growth Serum		1395.00	Abdominal cramps. Photosensitivity. Rash. Burning. Excess of creatinine in urine.
56	GLYCOLIC ACID 12%	GLYCO - A - 30 GM	Cream	The first night leave the Glycolic Cream® on for 10 minutes then wash off in cool water.	208.00	Itching, redness.
		GLYCO-6% - 30GM			130.00	
57	SLS, GLYCERIN, LANOLIN, DISODIUM EDTA	AQUASOFT-75GM SOAP	Soap			Itching, redness
58	FUCIDIC ACID & BETA METHASONE VALERATE	FUCIBET - 15 GM	Cream	Adult : Topical Apply to the affected area(s) bid. Max duration: 2 wk.	43.70	Itching, redness.
59	BENZALKONIUM CHLORIDE+ZINC OXIDE	HAPPY NAP-20gm CREAM	Cream	Put a thin layer on the affected skin(diaper rash)1-3 times a day. Do not wash the treated area after immediately applying Happynap Cream.	92.00	Hives; difficulty breathing; swelling of your face, lips, tongue, or throat.
60	BENZALKONIUM & ZINC	HAPPY NAP - 15GM CREAM	Cream	Apply twice or thrice daily.	65.00	Itching, redness

61	BENZOYL PEROXIDE 5%	BENZAC AC - 20 GM GEL	Gel	Apply once daily or twice to cover affected areas after washing.	125.95	Itching, redness
		BENZAC-AC 5%-20gm GEL	Gel	Apply to affected skin two or three times a day.	166.90	Skin reactions such as peeling, itching, irritation, and reddened skin may occur, especially at the start of treatment.
62	CALAMINE	LINICAL- 100ML LOTION	Lotion	Topical Apply 1-4 times/day.	110.00	Itching, redness.
		VENUSIA SOFT-75ML LOTION			275.00	
63	CICLOPIROX +ZINC PYRITH	8X - 100ML SHAMPOO	Shampoo	Wet your hair, take approximately 1 teaspoon of 8x shampoo (or 5ml) apply onto the scalp. Or u can use for long and thick hair 2 teaspoons of 8x shampoo (10ml).Apply it equally onto the affected area and massage into scalp, Leave it for 3 minutes. The eyes must be protected and rinse off with water.The procedure should be repeated twice in a week for 1 month. There should be minimum 3 days of gap per each cycle.	273.35	Burning sensation, redness, swelling, irregular heartbeat, rash, dandruff, headache and skin problems.

64	CETEARETH,PURIFIED WATER, GLYCERIN, HYDROGENATED POLYISOBUTENE, CETEARYL ALCOHOL,	CETAPHIL DAM-30gm LOTION	Lotion.	<p><u>Cetaphil Lotion Intense gel</u> Apply &amp; gently rub into the scar once a day for 8 wk (on new scars) &amp; 3-6 mth (on existing scars).</p> <p><u>Cetaphil Lotion Proactive gel</u> : Apply &amp; gently massage into the scar at least bid for 8 wk (on new scars) &amp; 3-6 mth (on existing scars).</p> <p><u>Cetaphil Lotion PM Overnight cream</u> : Apply &amp; gently rub into scar once nightly for 8 wk (on new scars) &amp; 3-6 mth (on existing scars).</p> <p><u>Cetaphil Lotion Kids cream Scars</u> : Apply a thin coat to the scar &amp; gently massage in tid for 8 wk (on new scars) &amp; tid for 3-6 mth (on existing scars).</p> <p>Sunscreen : Apply liberally 15 min before sun exposure. Reapply at least every 2 hrs.</p>	152.00	Stinging, contact dermatitis, and burning.
65	CLOBETASOL-0.05% W/W+SALICYLIC ACID-3% W/W	CLOP-S-20gm LOTION	Lotion	You usually put Clobetasol propionate on your scalp in the morning and at night.	191.40	Skin peeling Skin thinning Application site reactions (burning, irritation, itching and redness).
66	COAL TAR & SALICYLIC ACID	HYDROMAX CT-100gm	Cream	Massage the solution into wet hair. Rinse, then apply again and massage in hair for about 5 mins before rinsing. May be used daily.	270.00	Itching, redness.
67	CALCIPOTRIOL	PASITREX-20gm OINMENT	Ointment	Apply 1-2 timesday	330.00	Skin Irritation.
68	HALOBETASOL PROPIONATE	HALONOVA-15gm CREAM	Cream	Topical- Apply a thin layer to the affected skin once or twice daily.	129.80	Dry skin, redness, skin scar, white patches, vesicles and rash.

69	HALOBETASOL PROPIONATE	HALOVATE-S - 15GM	Lotion, Ointment	Adult Dose/ 12 years older .Apply a thin layer to affected area once or twice a dayTreatment should be limited to two weeks.	179.00	Itching, redness.
		ULTRAVEX - S3			127.00	
		ULTRAVEX - S6			169.50	
70	DERMADEW SOAP SOAP: COCOA BUTTER , COCONUT OIL , PALM OIL , GLYCERIN , ALOE VERA , OLIVE FRUIT EXTRACT.	DERMADEW-75gm SOAP	Soap	Wet the face or body with water Apply Dermadew soap Gently work up the lather Rinse thoroughly.	142.00	Drowsiness, dizziness, hypotension or a headache.
71	DESONIDE	DESOWEN - 10GM	Cream	Apply as thin film to the affected area bid/tid depending on the severity of the condition.	126.99	Itching, redness
72	FLUOCINOLONE TOPICAL	SEBOWASH-100ml SHAMPOO	shampoo	Apply about one ounce of the shampoo to the scalp once daily by working into a lather. Allow it to remain on the scalp for 5 minutes.	175.00	Application site reactions (burning, irritation, itching and redness).
73	MUPIROCIN + METRONIDAZOLE	MUPIMET-5gm OINTMENT	Ointment	Applied to the affected area three times daily	109.00	Mild burning or stinging
74	MUPIROCIN + METRONIDAZOLE NITROFURANTOIN	CO-MUPIMET-5ml COLLAGENS	Collagens	Adults: Clean the affected area and applied to the lesion three times a day for 10 days(bacterial infection).	450.00	Dry skin, Application site reactions (burning, irritation, itching and redness).
75	NADIFLOXACIN	NADIBACT - 10GM CREAM	Cream	Nadifloxacin cream 1 % w/w: Applied to the lesion two times a day till the lesion resolved	63.50	Sweating, Dryness of skin
		NADOXIN CREAM - 10 GM			84.50	
76	BECLOMETHASONE+LIDO CAINE+PHENYLEPHRINE	ANOVATE-20gm	Cream	Applied to the affected area once or twice daily.	91.75	Application site irritation.
77	FRAMYCETIN SULPHATE	SOFRAMYCIN	Cream	Adult Dose	31.29	Itching, redness.
78	POLYMYXIN + NEOMYCIN + BACITRACIN + HYDROCOTISONE	NEOSPORIN - H-5GM	Ointment	Adult and children (>2 years): Clean the affected area and apply ointment bd	44.50	Nausea, Diarrhoea, Vomiting
79	OZENOXACIN	OZEWID-1%-5gm	Cream	Twice a day	154.00	Rash,Application Site Itching Swelling Of Face, Lips, Eyelids, Tongue, Hands And Feet Seborrhoeic Dermatitis.

80	TRICLOSAN	SOAPEX CLASSIC-75gm SOAP	Soapex	Used as bathing soap daily for antibacterial action. Wet the face and body with water Apply the soap and gently work up the lather. Rinse thoroughly.	113.20	
81	PERMETHRIN	HHMITE-75gm	Soap	Twice daily	138.00	Itching, Scalp irritation. Skin irritation.
82	LULICONAZOLE	LULICAN-10gm	Cream	Apply an amount sufficient to cover the affected area.	150.00	Burning, irritation, itching and redness.
		LULICAN-10ml	Lotion		238.50	
83	MICONAZOLE	DK - GEL	Gel	Adult & children: Apply thinly over affected area bid for 2-6 week. Continue treatment for at least 1 wk after symptoms have disappeared.	116.00	Redness, severe blistering, irritation of skin, itching, peeling, dry skin, dry mouth, sore tongue, tooth pain, headache.
84	ALOE VERA GEL LIGHT LIQUID PARAFFIN GLYCERIN CALAMINE PRAMOXINE HYDROCHLORIDE	ATARAX ANTI-ITCH-100ml	Lotion	Apply sufficient quantity.	403.50	Drowsiness, dizziness, blurred vision, constipation, or dry mouth.

### MULTIVITAMINS, MINERALS AND IRON

1	IRON SUCROSE	MICROFER-100mg	Injection	100 mg diluted in a maximum of 100 mL of 0.9% NaCl.	300.10	Bloating or swelling of the face, arms, hands, lower legs, or feet, chest pain or tightness in the chest, difficult or labored breathing, headache, slow or fast heartbeat, tingling of the hands or feet, unusual tiredness or weakness.
2	NATIVE TYPE 11 COLLAGEN CAPSULES	GERIJOINT-40mg	Capsule	2.5 mg daily for up to 24 weeks	41.00	Upset stomach or throwing up. Constipation. Change in color of stool to green. Diarrhea. Belly pain.

3	SODIUM CHLORIDE IP 0.526 G, SODIUM GLUCONATE USP 0.502 G, SODIUM ACETATE TRIHYDRATE IP 0.368 G, POTASSIUM CHLORIDE IP 0.037 G, MAGNESIUM CHLORIDE HEXAHYDRATE IP 0.030 G, WATER FOR INJECTIONS	KABILYTE-500ML IV	Injection		312.00	Headache, Nausea, Vomiting Diarrhoea, Rashes.
4	TERIPARATIDE	ROCKBON-750mcg/3ml INJ. With Needles (30's)	Injection	25 micrograms administered once daily.	7139.00	Pain, swelling, redness.
		BONISTA CATRIDGE	Injection	20mcg /day	7117.50	Joint pain, Injection site reactions (pain, swelling, redness), Nausea, Leg cramps.
5	FOLIC ACID, METHYLCOBALAMIN, VITAMIN B1.	NUROKIND GOLD	Tablet	1tab once or twice /daily	89.00	Drowsiness, dizziness, hypotension or a headache
6	ADEMETIONINE / S- ADENOSYL METHIONINE	NUSAM-200mg	Tablet	1tab OD/day, Adults: 1-2 tablets/day.	285.00	Headache, Feeling anxious or nervous, Vomiting, Abdominal pain, Diarrhea, Insomnia, Increased sweating, Nausea, Dizziness.
		NUSAM-400mg			450.00	
7	NICOTINAMIDE PYRIDOXINE HYDROCHLORIDE, CYANOCOBALAMIN, ZINC SULPHATE, LYSINE HYDROCHLORIDE, ALCOHOL, NON-SYRUPY BASE, ABSOLUTE ALCOHOL I.P.	BETONIN AST-200ML	Syrup	two tablespoons of Betonin Syrup	176.79	Allergic rejection, Acute gastritis, Abdominal cramps, Allergic sensitization.
8	VITAMIN A 1.5 MG, VITAMIN D 10 MG, THIAMINE MONONITRATE (B1) 1.5 MG, RIBOFLAVIN (B2) 1.7 MG, ASCORBIC ACID 60 MG, NIACINAMIDE 20 MG, PYRIDOXINE (B6) 2 MG, FOLIC ACID 0.4 MG, CALCIUM PANTOTHENATE 10.92 MG, CYANOCOBALAMIN (B12) 6 MG, VITAMIN E 15.00 IU, FERROUS SULPHATE 50 MG, POTASSIUM IODIDE 196 MG, POTASSIUM SULPHATE 11.14 MG, MANGANESE SULPHATE MONOHYDRATE 1 MG, CUPRIC SULPHATE PENTAHYDRATE 2 MG, ZINC SULPHATE 37.04 MG	FULFIL FORTE	Capsule	Once a day	119.80	Acne, Hot Flashes, Injection Site Pain Weight Gain, Prostate Enlargement Increased Red Blood Cells.

9	IRON SUCROSE-100MG/5ML	OROFER-S -200mg	Injection	50-100 mg, taken three times daily	573.44	Muscle cramps,Nausea,Vomiting Taste changes,Diarrhoea,Constipation Headache,Cough.
10	NATURAL GINSENG AND MAGNESIUM, IMPROVE MENTAL ALERTNESS AND CONCENTRATION, AND INCREASE THE ABILITY TO MANAGE STRESS, DAILY ESSENTIAL VITAMINS, MULTIVITAMINS AND MINERALS SUCH AS VITAMIN C, VITAMIN D AND ZINC HELP IN IMPROVING IMMUNITY PREVENTING FREQUENT ILLNESS, BOOSTS ENERGY, IMPROVES STAMINA AND SUPPORTS IMMUNITY, ITS KEY INGREDIENTS, NATURAL GINSENG, VITAMIN B COMPLEX AND IRON, HELP SUPPORT DAILY ENERGY NEEDS, AND FIGHT TIREDNESS THROUGHOUT THE DAY	REVITAL-H WOMAN	Tablet	Once a day	375.00	Mild nausea and vomiting.
11	THIAMINE NITRATE, PYRIDOXINE HCL, FOLIC ACID, ELEMENTAL ZINC, ELEMENTAL MAGNESIUM, ELEMENTAL SELENIUM AND ELEMENTAL COPPER.	ALCOMAX-5ml	Injection		77.50	Kidney stones may rarely happen.
12	MECOBALAMIN JP PYRIDOXINE HYDROCHLORIDE IP FOLIC ACID IP VITAMIN D3 IP (AS STABILISED)	HOMIN-D3	Tablet	Twice a day	120.00	Confusion,Seizures (uncontrollable jerking of limbs),Depression,Erythema Conjunctivitis,Neuropathy
13	CALCIUM CARBONATE (1250MG) + VITAMIN D3 (2000IU) + METHYLCOBALAMIN (1500MCG) + L-METHYL FOLATE CALCIUM (1MG) + PYRIDOXAL-5-PHOSPHATE (20MG)	OSTEOFIT HD	Tablet	2 Tablets before meals, daily 3 times	358.50	Dryness in mouth, Nausea, Constipation Abdominal pain, Headache
14	VIT.C(ASCORBIC ACID) AND ZINC	CIMUNE CHEWABLE TABLET(Vit.C+Zinc)	Tablet	Once Daily	92.00	

15	VITAMIN E 400 MG	EVION-400mg	Capsule	one capsule daily	24.58	Nausea, diarrhea, stomach cramps, tiredness, dizziness, blurred vision, and headache may rarely occur.
16	VITAMIN D3 ORAL SOLUTION	ARACHITOL-NANO-5ml ORAL SOLUTION	Oral Solution	Adults, elderly and children over 12 years of age – Swallow 2 twice daily, ideally two in the morning and two in the evening. Children under 12 – Do not give Vitamin D3 to children under 12 years.	65.12	Constipation, wind, feeling sick, stomach ache, diarrhoea •Skin rash •Hypercalcaemia.
17	SAFFLOWER OIL 362.2 MG, FISH OIL 87.7 MG, BORAGE OIL 51.0 MG, ASCORBIC ACID 50.0 MG, ZINC SULPHATE 2.0 MG, VITAMIN E 10.0 IU (AS DL-ALPHA TOCOPHERYL ACETATE). OMEGA 3: EICOSAPENTAENOIC ACID (EPA) 15.8 MG, DOCOSAHEXAENOIC ACID (DHA) 10.5 MG. OMEGA 6: GAMMA LINOLENIC ACID (GLA) 10.2 MG, LINOLENIC ACID (LA) 280.8 MG.	EICOSA	Capsule	One capsule daily	11.00	Digestive tract, liver or kidneys.
18	THIAMINE	THIAMIN-2ml	Injection	<u>beriberi</u> :10 mg to 20 mg of thiamine hydrochloride are given IM 3 times daily for as long as 2 weeks.  Infantile beriberi :doses of 25 mg may cautiously be given IV.Neuritis of pregnancy :5 mg to 10 mg of thiamine hydrochloride IM daily.Wernicke-Korsakoff	50.40	Pain/redness/soreness at the injection site.



		AMINTOZ-100mg-1ml (THIAMINE)		syndrome, thiamine hydrochloride has been administered IV in an initial dose of 100 mg, followed by IM doses of 50 mg to 100 mg daily. Patients with marginal thiamine status to whom dextrose is being administered should receive 100 mg thiamine hydrochloride in each of the first few liters of IV fluid to avoid precipitating heart failure.	25.20	
19	PYRIDOXINE HCL	BENADON - 40 MG	Injection & Tablet	150 mg per day	13.28	Severe peripheral neuropathies (with long-term admin of large doses).
20	METHYLCOBALAMIN	NUROKIND FAST STRIPS	Strips	1tab/once or twice daily	14.00	Headache, nausea, vomiting, dizziness, diarrhoea, back pain, arthritis, back pain, anxiety, nervousness, lack of coordination in body movement.
		NUROKIND	Injection	500 mcg/day 3 times/wk	15.00	Gastrointestinal discomfort.
21	METHYLCOBALAMIN & ALPHA LIPOIC ACID, FOLIC ACID & PYRIDOXINE	RENERVE PLUS	Capsules		117.00	
22	METHYLCOBALAMIN + NIACINAMIDE + VITAMIN B6	RE-XITE PLUS-2ml	Injection	500 mcg daily IM/IV 3 times/wk.	22.00	Unusual or unpleasant taste in mouth.
23	METHYLCOBALAMIN THIAMINE MONONITRATE PYRIDOXINE HCL, FOLIC ACID	RE-XITE-BT	Tablet	1 tab once or twice daily	65.00	Hypersensitivity, Abdominal and stomach pain, Blurred vision, Bloating, Headache, Irregular heartbeat, Hair loss, Constipation, Dry mouth.
24	MECOBALAMIN, FOLIC ACID, PYRIDOXINE HYDROCHLORIDE, VITAMIN D3, CYANOCOBALAMIN, VITAMIN B12 (CYANOCOBALAMIN)	RE-XITE-D3	Tablet	1 tab once or twice daily	13.00	Nausea/vomiting, loss of appetite, unusual weight loss, mental/mood changes, signs of kidney problems (such as change in the amount of urine), bone/muscle pain, headache, increased thirst, increased urination, weakness, tiredness.
25	MENADIONE	VITAMIN - K - 1 ML	Injection	IV Vit K Deficiency 0.03 mcg/kg/day. IM Haemorrhage or threatened haemorrhage during severe hypoprothrombinaemia 2.5-10 mg/day.	6.00	Facial flushing, sweating, chest constriction/pain, dyspnoea, cyanosis, CV collapse. IM: Severe shock-like reactions, phlebitis, inj site reactions.

26	CALCIUM ASCORBATE 75 MG+COENZYME Q10 100 MG+DOCOSAHEXAENOIC ACID 60 MG+EICOSAPENTAENOIC ACID (EPA) 90 MG+L- ARGININE 100 MG+LEVOCARNITINE 50 MG+LYCOPENE 5000 MCG+MIXED CAROTENE 10.33 MG+PIPERINE 5 MG+SELENIUM DIOXIDE 100 MCG+VITAMIN B12 5 MCG+WHEAT GERM OIL 25 MG+ ZINC SULPHATE 6.5 MG	KQ-100-CAPSULE	Capsule	once a day	425.00	Very upset stomach or throwing up. Very loose stools (diarrhea).
----	---	----------------	---------	------------	--------	--

27	CYNOCOBALAMINE	CYNOCOBALAMINE INJECTION-30ml	Injection	<p><b>Pernicious Anemia</b> :Adult Dosages  100 mcg intramuscularly or subcutaneously once daily for 6-7 days, then every other day for 7 doses, then 3-4 days for 2-3 weeks, then monthly  /Alternative parenteral dosing:  1,000 mcg intramuscularly or subcutaneously once daily for 7 days, then weekly for 1 month, then monthly  Pediatric Dosages :30-50 mcg intramuscularly or subcutaneously once daily for 2 weeks or more for total dose of 1,000 mcg to 5,000 mcg administer concomitantly with 1 mg/day of folic acid for 1 month  Maintenance: 100 mcg intramuscularly or subcutaneously monthly</p> <p><b>B12 Deficiency</b> :Adult Dosages  Initial: 30 mcg intramuscularly once daily for 5-10 days.  Maintenance: 100-200 mcg intramuscularly monthly  Pediatric Dosages :0.2 mcg/kg for 2 days; follow by 1,000 mcg/day for 2-7 days; follow by 100 mcg/day for 2-7 days; then 100 mcg/week for 1 month</p>	30.00	Pain/redness at the injection site, mild diarrhea, itching, or a feeling of swelling all over the body.
28	MULTIVITAMIN PREPARATIONS	RICONIA L.P	Capsule, Syrup, Tablet,	One capsule daily, POLYBION LC - 100ML 5ml single dose, FOURTS - B 1 tab daily ,Aquasol -A Vitamin A Deficiency Malabsorption or oral administration not feasible: 100,000 U/day IM for 3 days; then 50,000 U/day for 2 weeks; follow with oral therapy.	67.00	Increased Urination

	POLYBION LC - 100ML 5ml single dose	Drops, Injections	Oral therapy: Take oral therapeutic multivitamin containing 10,000-20,000 U/day vitamin A for 2 months. Deficiency prophylaxis: 10,000- 50,000 U PO qDay	47.50
	FOURTS - B 1 tab daily		Xerophthalmia (Off- label).Recommended dose except for females of reproductive age: 200,000 units PO qDay for 2 days; repeat dose again after 2 weeks.Females of reproductive age with night blindness or Bitot's spots: 5000- 10,000 units/day; 10,000 units/day maximum or $\leq$ 25,000 units once weekly for $\geq$ 4 weeks, Neurobion Forte - 1-2 tab /day, Limcee - 500 mg :Adults (including the elderly) For the treatment of Vitamin C deficiency: 250mg - 1000mg daily in divided doses For the prevention of Vitamin C deficiency: 25mg - 75mg daily The maximum daily dose is 1000mg. Children (aged 4 to 12 years) For the treatment of Vitamin C deficiency: 125mg - 500mg daily in divided doses For the prevention of Vitamin C deficiency: 25mg - 75mg daily Children under 4 years Ascorbic Acid Tablets are not suitable for children under 4 years of age.	108.00
	AQUASOL - A		JUSTDEE : Adult Dose for Vitamin D Insufficiency: 400 to 1000 international units orally once a day.Vitamin D Deficiency:1000 international units orally	24.55

		NEUROBION FORTE		<p>once a day. Vitamin D Deficiency: 1000 international units orally once a day. Dose for Prevention of Falls: 800 international units orally every day (with calcium). Geriatric Dose for Prevention of Fractures: For patients over 65 years of age: 100,000 international units orally every 4 months. Usual Pediatric Dose for Vitamin D Insufficiency: Neonates, infants and children: 200 international units orally once daily.</p>	7.22	
		LIMCEE - 500 MG		<p>Alternate dosing: Premature neonates: 400 to 800 international units orally once daily or 150 to 400 international units/kg/day. Breastfed neonates and infants (fully or partially breastfed): 400 international units orally once daily beginning in the first few days of life. Continue supplementation until the infant is weaned to greater than or equal to 1000 ml/day or 1 qt/day of vitamin D fortified formula or whole milk (after 12 months of age).</p>	15.34	

	JUSDEE 400 - 15 ML		Formula fed neonates and infants ingesting less than 1000 ml of vitamin D fortified formula: 400 international units orally once daily. Children ingesting less than 1000 ml of vitamin D fortified milk: 400 international units orally once daily. Adolescents without adequate intake: 400 international units orally once daily. Children with increased risk of vitamin D deficiency (chronic fat malabsorption, maintained on chronic antiseizure medications): Higher doses may be required; use laboratory testing [25(OH)D, PTH, bone mineral status] to evaluate. Usual Pediatric Dose for Vitamin D Deficiency:Treatment of Vitamin D deficiency and/or rickets:Infants 1 to 12 months: 1000 to 5000 international units/day for 2 to 3 months;	61.71	
	D-RISE--60K		once radiologic evidence of healing is observed, dose should be decreased to 400 international units/day.	142.75	
	ARACHITOL- 3 I.U -1 ML		High Dose Regimen: Patient less than 5 years: 12,000 international units twice weekly for 12 weeks. Patient 5 years or older: 50,000 international units twice weekly for 12 weeks.	27.50	
	FOURTS - B			108.00	
	BEPLEX FORTE-TABLET	Tablet	Once daily	29.22	Constipation, diarrhea, or upset stomach.
	BECOZINC-CAPSULE	Capsule	1 cap OD	24.00	Swelling of lips, Swelling of face and tongue, Difficulty in breathing, Diarrhea, Swelling, Rapid weight gain, Itching or mild rash, Nausea, Stomach upset, Heartburn.

		MULTI VITAMIN-10ml INJECTION	Injection	daily dosage volume is 10 mL. One daily dose (10 mL) is diluted by adding directly to a specified volume of an intravenous fluid.	25.00	Severe allergic reaction (anaphylaxis), abnormally high levels of vitamin A, rash, redness, itching, headache, dizziness, agitation, anxiety, and double vision.
		SUPRADYN-TABLET	tablet	1-2 tabs daily	28.05	Nausea, Vomiting, Diarrhea, Constipation, Upset stomach, Black or dark-colored stools or Temporary staining of the teeth.
29	MECOBALAMIN,ALA,INOSITOL,FOLIC ACID,CHROMIMUM,SELENIUM.ZINC	VITASURGE-TABLET	Tablet	1-2 tab/day	159.00	Sleepiness, Colic pain, Injection site related side effects, Pain in abdomen, Abdomen pain, Erosion, Headache, Nausea, Allergic reactions, Vomiting.
30	MECOBALAMIN	NUROKIND - 500MG	Injection & Tablet	Oral :Peripheral neuropathies Adult: 1500 mcg/day in 3 divided doses. Parenteral :Peripheral neuropathies Adult: 500 mcg daily IM/IV 3 times/wk. Parenteral :Megaloblastic anaemia caused by vitamin B12 deficiency Adult: 500 mcg daily IM/IV 3 times/wk. Maintenance dose: After about 2 mth of therapy, reduce dose to single admin of 500 mcg every 1-3 mth.	32.40	GI Irritation, Nausea, Vomiting.
		NUROKIND			12.00	
		RENERVE			83.50	
31	MECOBALAMIN, ZINC SULPHATE, VITAMIN B6, FOLIC ACID, NIACINAMIDE, CHROMIUM PICOLINATE	NUROSTAR-TABLET	Tablet	1 capsule a day.	4.62	Skin irritation, Headache, Dizziness Nausea, Mood changes.
32	L-AEGININE, CARNITINE, PIPERINE,OMEGA 3 LYCOPENE	K.Q-100	Capsule	once daily	390.00	Nausea, Vomiting.
33	L-METHYLFOLATE, METHYLCOBALAMIN, PYRIDOXAL 5-PHOSPHATE	FOLVITE-ACTIVE-CAPSULE	Capsule	one tablet daily	12.10	Rash, itching/swelling (especially of the face/tongue/throat), dizziness, or trouble breathing
34	LEVOCARNITINE 150 MG+VITAMIN E 200 MG	EVION-LC-TABLET	Tablet	1 to 2 tablets in a day.	35.42	Rare incidences ofgastrointestinal disturbanes such as nausea, vomiting, diarrhoea, abdominalcramps.
35	B-COMP FORTE VIT-C	BECOSULES	Capsules	1tab/BD	25.80	Itching, redness.
		BECOSULES - Z			25.96	

36	BIOTIN 10 MCG+ DL-METHIONINE	MACTOTAL-200ml SUSPENSION	Suspension	1tab/bd	106.00	Constipation, diarrhea, or upset stomach.
37	BIOTIN+CYANOCOBALAMIN+ELEMENTAL IRON+ELEMENTAL ZINC+FOLIC ACID	HEMFER-225ML SYRUP	SYRUP	5ml Once daily.	135.00	Bitter taste in the mouth Darky stool may occur Vomiting Excitement Eczema Gastrointestinal discomfort Pulmonary edema.
38	FOLIC ACID 5 MG+ METHYLCOBALAMIN 1500 MCG+ PYRIDOXINE 20 MG	HOMISURE-TABLET	Tablet	50mg three times a day.		Wheezing; chest tightness; fever; itching; bad cough; blue or gray skin color; seizures; or swelling of face, lips, tongue, or throat.
39	ALPHA LIPOIC ACID, FOLIC ACID, MECOBALAMIN, PYRIDOXINE, THIAMINE MONONITRATE	MYLAMIN-TABLET	Tablet	Adult- PO- The recommended dose range is 150 to 400mg/day.	46.75	Allergic reactions, Altered sleep patterns, Nausea, Itch, Dysfunction of nerves resulting in numbness, Vomiting, Vertigo, Anorexia, Abdominal distention, Flatulence.
40	FOLIC ACID, METHYLCOBALAMIN, VITAMIN A, VITAMIN B1, VITAMIN B2, VITAMIN B3, VITAMIN B5, VITAMIN B6, VITAMIN C, VITAMIN E	A TO Z NS-200ml SYRUP	Syrup	For adults: 3 - 4 teaspoonfuls daily; For children of 4 - 12 years: 2 - 3 teaspoonfuls daily;	121.00	Constipation, diarrhea, or upset stomach.
		A TO Z-15ml DROPS	Drops		32.71	Feeling tired or weak (fatigue), stomach pain, loose stools (diarrhea), headache, nausea, or blurred vision.
41	FOLIC ACID(VITAMIN B9), VITAMIN B12, VITAMIN D3, VITAMIN B1, NIACINAMIDE, VITAMIN B2, VITAMIN B6, CALCIUM PANTOTHENATE, VITAMIN A, VITAMIN C, VITAMIN E, ZINC	A TO Z-TABLET	Tablet	Once daily	79.45	Tooth staining, increased urination, stomach bleeding, uneven heart rate, confusion, and muscle weakness or limp feeling.
42	BIOTIN, AMINOACIDS, VITAMINS, MINERALS, NATURALEXTRACTS	FOLLIHAIR TABLET		1-2cap/day	155.00	Nausia, Vomiting, Head ache
43	L-CARNITINE L-TARTRATE, MECOBALAMIN & FOLICACID	NUROKIND LC	Tablet	1,500 mcg/day in 3 divided .	89.89	Anorexia, nausea, vomiting and diarrhoea.
44	CALCIUM CARBONATE	SHELCAL - 500 MG	Powder, Suspension	Initial: 1 mcg/day. Maintenance: 0.25-1 mcg/day.	62.00	Constipation, bloating, flatulence.
45	CALICIUM CITRATE	ROCKBON		1200 mg/day.	160.00	
		SHELCAL CT			178.00	



46	CALCITRIOL	ROCALTROL - 0.25mcg	Capsule	0.015-0.02 mcg/kg/day. Maintenance: 0.03-0.06mcg/kg/day.	184.78	Weakness; headache; somnolence; nausea; vomiting; dry mouth; constipation; muscle pain; bone pain; metallic taste; polyuria; polydipsia; anorexia.
47	CALCIUM DOBESILATE	OSIL	Capsule, Injection,	Adult: PO 0.5-1 g/day in divided doses.	130.00	Skin rash, fever, vertigo.
		CALCIUM GULCONATE - 10 ML	Suspension		20.00	
		CALCIMAX-P - 200ML		99.00		
		DOBESIL-CAPSULE		153.00	Dobesil is Allergic reaction.	
48	CALCIUM FOLINATE	NEOVORIN-50mg/5ml-INJECTION (CALCIUM FOLINATE)	Injection	200 mg/ml by intravenous infusion over two hours, followed by bolus 400 mg/ml.	81.00	Constipation, oral inflammation or cold sores, fatigue, hair loss, or loss of appetite.
49	UBIDECARENONE+L-CARNITINE+L-TARTRATE+CHOLINE BITARTRATE+ SELENIUM	ASTHOQUE-TABLET	Tablet	2 tablets a day.	59.00	Drowsiness, dizziness, headache, excitement, nausea, vomiting, and diarrhea.
50	UNDENATURATED TYPE-II COLLAGEN, CALCIUM, CHOLECALCIFEROL, VITAMIN C	OSTOSHINE-TABLET	Tablet	once a day in the morning	70.00	Rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat.
51	D CHIRO INOSITOL, FOLIC ACID & MYO INOSITOL	APCOD SOFTGEL	Softgel	once or twice a day	490.00	Excessive thirst. Skin allergic reactions, such as rashes or redness. Abdominal pain. Constipation.

52	AMINO ACID	NUTRIVEN 10% I.V-200ml INJECTION*(AMINO ACID)	Injection	<p>Total Parenteral Nutrition (Central Infusion) :In unstressed adult patients with no unusual nitrogen losses, a minimum dosage of 0.1 gram nitrogen (4.2 mL of 15% Amino Acids Injection)plus 4.4 grams (15 calories) of dextrose per kilogram of body weight per day are required to achieve nitrogen balance and weight stability. For patients stressed by surgery, trauma or sepsis, and those with unusual nitrogen losses, the dosage required for maintenance may be as high as 0.3 to 0.4 grams of nitrogen (13 to 17 mL 15% Amino Acids Injection) per kilogram of body weight per day.</p> <p>Peripheral Nutrition :Dilution of 250 mL 15% Amino Acids Injection in 750 mL of 10% dextrose will reduce the osmolarity to a level (724 mOsmol/L) which is more favorable to the maintenance of the integrity of the walls of the veins.</p>	450.00	Nausea, diarrhea, and stomach pain.
53	AMINO ACIDS & VITAMINS	OVABLESS	Tablet, Capsules,	1-2 cap/day	12.00	Nausia, Pain, Head ache
		ASTYMIN FORTE	Injection	1-2 cap/day	96.00	
		ASTYMIN - 3 - 200 ML I.V		adult:200-800mL iv once/daily;children:0.2-0.25 grams of nitrogen per kg body weight; drip drate 15-20 drops initially followed by 20-40 drops per min	416.50	
		KETOADD		200mg bd or tds/day	299.00	

54	FERROUS ASCORBATE & FOLIC ACID	OROFER-XT	Tablet	The most recommended dosage in adults is one-two tablets/ tablespoon right after a meal.	13.00	Vomiting, Nausea, Altered sleep patterns, Constipation, Flatulence, Anorexia.
55	FERRIC CARBOXYMALTOSE	FERRIUM-1K-20ML	Injection	750 mg iron / 15 mL single-dose vial.	5374.50	Nausea, Flushing, Dizziness, High blood pressure, Low levels of phosphorus in the blood, Injection site reactions.
56	CALCIUM PANTOTHENATE 10 MG+ELEMENTAL ZINC 10 MG + SILYMARIN 70MG	HEPTOGON	Tablet	Recommended dose of Silymarin is 140 mg 2-3 times/day.	159.00	Constipation, diarrhea, or upset stomach.
57	CALCIUM AND VITAMIN D3	SHELCAL-HD	Tablet	Once a day in the morning	6.27	Nausea, vomiting, constipation; increased thirst or urination; muscle weakness, bone pain
58	CALCITRIOL + CALCIUM CITRATE	OSTEOFIT-C		Initial: 0.25 mcg/day		Constipation, wind, feeling sick, stomach ache, diarrhoea.
59	BENFOTIAMINE+ BIOTIN + CALCIUM ASCORBATE + CHOLINE + CITRUS BIOFLAVONOIDS + DOCOSAHEXAENOIC ACID +EICOSAPENTAENOIC ACID + ELEMENTAL CALCIUM + ELEMENTAL ZINC + FOLIC ACID + GINGKO BILOBA + GINSENG + GRAPE SEED EXTRACT + GREEN TEA POWDER + LACTOBACILLUS 500 MILLION CELLS + LUTEIN + MIXED CAROTENE + PIPERINE + POTASSIUM + VANADIUM + VITAMIN D3 400 IU + VITAMIN K + WHEAT GERM OIL.	MACTOTAL	Tablet	1 tab/bd.	165.00	Constipation, diarrhea, or upset stomach.
60	ELEMENTAL CALCIUM	MAXICAL-500mg	Tablet	1000mg/day	71.50	Nausea/vomiting, loss of appetite, unusual weight loss, mental/mood changes, bone/muscle pain, headache, increased thirst/urination, weakness, unusual tiredness.

61	CARBONYL IRON WITH ZINC AND FOLIC ACID	FEFOL-Z	Capsule	1 cap od	78.70	GI irritation, Nausea, Vomiting.
62	FERRIC PREPARATIONS	HEMSI-PD - 15ML	Drops & Injection	IRON DEFICIENCY IN INFANTS AND TODDLERS : Breastfed preterm infants - elemental iron of 2 mg/kg per day from 1 month of age through 12 months ,Formula-fed preterm infants, 2 mg/kg per day of iron from a standard infant formula.	37.30	Constipation.
		FERIUM - 500MG			ferium dose : 5 ml bid	
		DEXORANGE-200ml	Syrup	Dexorange syrup -Adult: 15-30ml; child: 2-5 yrs: 5ml; child 5-12 yrs: 10ml. All bid after meals.	109.25	Nausea, Vomiting, Indigestion, Diarrhoea, Bitter taste in mouth, Dark coloured stool.
		DEXORANGE	Capsule	Dexorange capsule -1 cap bid after meals.	106.56	

63	FERROUS PREPARATION	LIVOGEN-XT	Injections, Syrup, Tablet	Livogen XT Adult Dose for Megaloblastic Anemia :1 mg orally, intramuscularly, subcutaneously or IV once a day. Adult Dose for Folic acid (Livogen-XT) Deficiency :400 to 800 mcg orally, intramuscularly, subcutaneously or IV once a day. Women of childbearing age, pregnant, and lactating women: 800 mcg orally, intramuscularly, subcutaneously or IV once a day. Pediatric Dose for Folic acid (Livogen-XT) Deficiency Infant: :0.1 mg orally, intramuscularly, subcutaneously or IV once a day. Child: Less than 4 years: up to 0.3 mg orally, intramuscularly, subcutaneously or IV once a day. 4 years or older: 0.4 mg orally, intramuscularly, subcutaneously or IV once a day. Pediatric Dose for Vitamin/Mineral Supplementation Recommended daily allowance (RDA): Premature neonates: 50 mcg/day (15 mcg/kg/day). Ferium XT :1 tab daily.	99.50	Upset Stomach
		FERIUM-XT		102.40		
		LIVOGEN		28.90		
		LIVOGEN-250ml		129.00		
		LIVOGEN - Z		31.00		
		FEFOL		67.00		
		FERIKIND-S -100MG - 5ML		242.00		
		IV CROSS - 100 MG - 5ML		336.00		

ZINCOLIFE-50ml		Diarrhoea – Children aged < 6 month: 2.5 ml once a day for 14 days. Children aged 6 months and above): 5 ml once a day for 14 days.	51.23	Diarrhea, Abdominal cramping, Vomit, Loss of appetite, Stomach cramps, Headache.
ZINCOVIT-200ml	Syrup	Zincovit syrup :Adult dose : 5ml.	106.00	nausea, vomiting, diarrhea, metallic taste, kidney and stomach damage
ZINCOVIT		Zincovit tablet :1 tablet daily	83.00	
Z&D-DS-20-15ml		Diarrhoea – Children aged < 6 month: 2.5 ml once a day for 14 days. Children aged 6 months and above): 5 ml once a day for 14 days.	67.00	Convulsions, Vomiting, Loss of appetite, Stomach cramps, Diarrhea, Headache.
C-PINK-150ml	Suspension	Elemental iron 30 MG + Folic acid 500 MCG /5ML	133.00	Diarrhea, nausea, vomiting, abdominal cramps/pain, or heartburn.
CPINK-15ml	Drops	Elemental iron 10 MG + Folic acid 100 MCG /5ML	74.70	
ASCAZIN	Tablet	Adults: 25 to 50mg zinc daily or 1 to 2 tablets daily. Children: 5mg/kg 1 to 3 times daily.	54.00	Nausea, vomiting, diarrhea, metallic taste,kidney and stomach damage.
LIV-52	Tablet	Drug-induced hepatitis:1 tablets b.i.d. with anti TB/statin/chemotherapy/antiretroviral treatment. Jaundice and anorexia during pregnancy:1tablet b.i.d. for 8 weeks. Chronic active hepatitis:1tablets b.i.d. for 6 months Early cirrhotic conditions :1tablet t.i.d. for 6 to 12 months.Anorexia and suboptimal growth: children 1 tds adult : 2tds.Alcohol-induced liver damage:Tablet: 2 to 3 tablets t.i.d.Acute viral hepatitis:children 1tds, adult : 2tds.	95.00	Burning of skin, Skin irritation, Irritation of skin.

64	CARICA PAPAYA	CARIPILL	Tablet	Adult :One Tablet to be taken 3 times daily for 5 days. Children aged 6 to 12 years : 10 ml 3 times daily for 5 days	410.50	Severe stomach/abdominal pain, nausea/vomiting, slow heartbeat, severe drowsiness, inability to move.
65	CHOLECALCIFEROL	DVION PEARL - 60K	Capsule	10 mcg/day.	109.75	Hyperphosphataemia or hypercalcaemia (in excessive intake). Associated effects of hypercalcaemia include hypercalciuria, ectopic calcification,& renal & CV damage.
66	CHOLECALCIFEROL	ARACHITOL-6L-I.U-1ml	Injection	0.25-1microgram/day.	44.15	Arachitol is Injection site reactions (pain, swelling, redness). Chest pain, feeling short of breath; growth problems.
		D-RISE-2000IU	Capsule	Once a Week	77.72	
		D-RISE-SACHET	Powder		36.89	
67	COLLOIDAL IRON,LYSINE, FOLIC ACID.VIT B12	TONOFERON - 15ml	Drops, Syrup	Infants less than 1 yr: 5 drops b.i.d. Children above 1 yr: 10 drops b.i.d.	63.00	GI Irritation, Vomiting, Nausea.
		TONOFERON SYRUP - (PAED )			78.00	
68	ELEMENTAL ZINC+FOLIC ACID	IFOLZIN ZZ-TABLET	Tablet	Once daily	86.00	Loose stools.

### ANTI-DIURETICS AND ITS ANALOGUES

1	FRUSEMIDE	LASIX - 40mg	Tablet, Injection	20-80 mg	7.40	Hyponatraemia, hypochloaemic alkalosis, hypokalaemia, headache, drowsiness, muscle cramps, hypotension.
		LASIX - 2ml			3.31	
		LASIX-4ml			6.72	
2	FRUSEMIDE + SPIRONOLACTONE	LASILACTONE			30.30	
3	SPIRONOLACTONE	ALDACTONE - 100mg	Tablet	100 mg/day, up to 400 mg/day	81.00	Fluid or electrolyte imbalance, GI upset, gynaecomastia, drowsiness, headache, hyponatraemia; tachycardia, hypotension, oliguria, hyperkalaemia.
		ALDACTONE - 25mg			30.50	
4	MANNITOL	MANNITOL - 100ml	Injection	iv 0.25-2 g/kg	36.75	Fluid and electrolyte imbalance; (with high doses). Nausea, vomiting, headache, dizziness, convulsions, fever; tachycardia, chest pain; blurred vision thirst;
5	TORSEMIDE	DYTOR - 10mg	Tablet	2.5-5 mg once daily. Max: 5 mg/day	55.00	Electrolyte disturbances e.g. hypokalaemia, dehydration, dry mouth, headache, dizziness, hypotension.
		DYTOR - 2 ml			13.00	
		DYTOR - 5 mg			33.75	
		TORSILONG-10mg	Tablet	2.5-5 mg once daily. Max: 10mg/day	41.00	

6	TAMSULOSIN	URONEX-0.4mg	Tablet	Once a day.	110.00	Low blood pressure (hypotension), Dizziness, Drowsiness, Reduced sex drive (libido), Inability to get an erection (impotence), Tenderness or enlargement of your breasts (in men), Sleeplessness, Runny nose.
		DYNAPRES-0.4MG	Tablet, Capsule	400 mcg once daily.	242.00	Orthostatic hypotension, syncope, priapism, headache, infection, asthenia, back pain, chest pain, dizziness, somnolence, insomnia.
		URIMAX - 0.4 MG			137.00	
7	TAMSULOSIN + FINASTEROIDE	URIMAX - F	Tablet, Capsule	400 mcg once daily.	169.00	Orthostatic hypotension, syncope, priapism, headache, infection, asthenia, back pain, chest pain, dizziness, somnolence, insomnia.
8	TAMSULOSIN + DUTASTEROIDE	URIMAX - D			328.50	
9	ALPHA KETOANALOUGE	KETOALPHA-TABLET	Tablet	4 to 8 tablets three times daily during meals.	34.00	Hypercalcemia.
10	TOLTERODINE	ROLITEN - 2 MG	Tablet, Capsule	2 mg twice daily	134.25	Dry mouth, chest pain, headache, somnolence, fatigue, dry skin, abdominal pain, constipation, dyspepsia, diarrhoea, dizziness.
		TEROL LA-4			194.50	
11	ANHYDROUS DEXTROSE 15 GM+CALCIUM CHLORIDE 0.257 GM+MAGNESIUM CHLORIDE 0.0508 GM+SODIUM CHLORIDE 5.38 GM+SODIUM LACTATE 4.48 GM /1000ML	PERITONEAL DIALYSIS SOLUTION-2000ml	Solution	Approximately two liters of dialysis solution are instilled into the peritoneal cavity of adults.	235.00	Abdominal pain, bleeding, peritonitis, subcutaneous infection.



12	ALBUMIN (HUMAN)	HUMAN ALBUMIN-20%-100ml	Injection	<p>Hypovolemia :In adults, an intravenous infusion of 25 g should be given. If adequate response (stabilization of circulation) is not achieved within 15 to 30 minutes, an additional dose may be given. In spite of limited information about the efficacy in pediatric subjects, an intravenous infusion of 2.5 to 12.5 g or 0.5 to 1 g/kg body weight may be given. If adequate response (stabilization of circulation) is not achieved within 15 to 30 minutes, an additional dose may be given. Hemodilution may follow administration of Albumin (Human) 20%. If hemorrhage has occurred, this may result in relative anemia. This condition should be controlled by the supplemental administration of compatible red blood cells or compatible whole blood.</p> <p>Hypoalbuminemia :In adults, intravenous infusion of 50 to 75 g of Albumin (Human) 20% may be used.</p> <p>In burns, therapy usually starts with the administration of large volumes of crystalloid injection</p>	4332.00	Flush, urticaria, fever and nausea.
13	OXYBUTYNIN	TROPEN-2.5mg	Tablet	Three times daily	82.25	Dry mouth, constipation, diarrhea, and headache.

